

Climate changes during the Lateglacial in South Europe: new insights based on pollen and brGDGTs of Lake Matese in Italy

Mary Robles^{1,2}, Odile Peyron², Guillemette Ménot³, Elisabetta Brugiapaglia¹, Sabine Wulf⁴, Oona Appelt⁵,
Marion Blache², Boris Vanni re^{6,7}, Lucas Dugerdil², Bruno Paura¹, Salom  Ansanay-Alex³, Amy Cromartie⁸,
Laurent Charlet⁹, Stephane Gu dron⁹, Jacques-Louis de Beaulieu¹⁰, S bastien Joannin^{2,3}

¹ Univ. Molise, Department of Agricultural, Environmental and Food Sciences, Campobasso, Italy

² Univ. Montpellier, CNRS, IRD, EPHE, UMR 5554 ISEM, Montpellier, France

³ Univ. Lyon, ENSL, UCBL, UJM, CNRS, LGL-TPE, F-69007 Lyon, France

⁴ Univ. Portsmouth, School of the Environment, Geography and Geosciences, Portsmouth, United Kingdom

⁵ Helmholtz Centre Potsdam, GFZ German Research Centre of Geosciences, Section 3.6, Telegrafenberg,
Potsdam, Germany

⁶ Chrono-Environnement, CNRS, Univ. Bourgogne Franche-Comt , Besan on, France

⁷ MSHE Ledoux, CNRS, Universit  Bourgogne Franche-Comt , Besan on, France

⁸ Cornell Univ., Department of Anthropology, Ithaca, NY, USA

⁹ Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, IRD, IFSTTAR, ISTerre, Grenoble, France

¹⁰ Aix-Marseille Univ., CNRS, IRD, UMR 7263 & 237 IMBE, Aix-en-Provence, France

E-mails : Mary Robles, robles@cerege.fr; Odile Peyron, odile.peyron@umontpellier.fr

Short Abstract

Quantitative climate reconstructions based on pollen and brGDGTs reveal, for the Lateglacial, a warm B lling–Aller d and a marked cold Younger Dryas in Italy, showing no latitudinal differences in terms of temperatures across Italy. In terms of precipitation, no latitudinal differences are recorded during the B lling–Aller d whereas the latitudes 40–42 N appear as a key junction point between wetter conditions in Southern Italy and drier conditions in Northern Italy during the Younger Dryas.

Abstract

The Lateglacial (14,700–11,700 cal BP) is a key climate period marked by rapid but contrasted changes in the Northern Hemisphere. Indeed, regional climate differences have been evidenced during the Lateglacial in Europe and the Northern Mediterranean areas. However, past climate patterns are still debated since temperature and precipitation changes are poorly investigated towards the lower European latitudes. Lake Matese in Southern Italy is a key site in the Central Mediterranean to investigate climate patterns during the Lateglacial. This study aims to reconstruct climate changes and their impacts at Matese using a multi-proxy approach including magnetic susceptibility, geochemistry (XRF core scanning), pollen data and molecular biomarkers like branched Glycerol Dialkyl Glycerol Tetraethers (brGDGTs).

40 Palaeotemperatures and -precipitation patterns are quantitatively inferred from pollen
41 assemblages (multi-method approach: Modern Analogue Technique, Weighted Averaging
42 Partial Least Squares regression, Random Forest, and Boosted Regression Trees) and brGDGTs
43 calibrations. The results are compared to a latitudinal selection of regional climate
44 reconstructions in Italy to better understand climate processes in Europe and in the circum-
45 Mediterranean region. A warm Bølling–Allerød and a marked cold Younger Dryas are revealed
46 in all climate reconstructions inferred from various proxies (chironomids, ostracods,
47 speleothems, pollen, brGDGTs), showing no latitudinal differences in terms of temperatures
48 across Italy. During the Bølling–Allerød, no significant changes in terms of precipitation are
49 recorded, however, a contrasted pattern is visible during the Younger Dryas. Slightly wetter
50 conditions are recorded south of latitude 42°N whereas dry conditions are recorded north of
51 latitude 42°N. During the Younger Dryas, cold conditions can be attributed to the southward
52 position of North Atlantic sea-ice and of the Polar Frontal JetStream whereas the increase of
53 precipitation in Southern Italy seems to be linked to relocation of Atlantic storm tracks into the
54 Mediterranean, induced by the Fennoscandian ice sheet and the North European Plain. By
55 contrast, during the Bølling–Allerød warm conditions can be linked to the northward position
56 of North Atlantic sea-ice and of the Polar Frontal JetStream.

57

58 **Keywords: Mediterranean region; Palynology; Molecular Biomarker; Paleoclimate;**
59 **Transfer functions; Tephra; Younger Dryas; Bølling–Allerød; Lateglacial**

60

61 **1. Introduction**

62

63 In the Northern Hemisphere, the Lateglacial (ca. 14,700-11,700 cal BP) is a period of
64 special climatic interest characterized by contrasted and rapid climate changes, associated with
65 the successive steps of the deglaciation and changes in atmospheric and ocean circulation
66 patterns (e.g., Walker et al., 2012; Rehfeld et al., 2018). Following the cold Oldest Dryas (OD)
67 period, the Bølling–Allerød (B/A) or Greenland Interstadial-1 (GI-1) began abruptly at 14,700
68 cal BP with warmer conditions. At 12,900–11,700 cal BP, the Younger Dryas (YD) or Greenland
69 Stadial-1 (GS-1) was the last main millennial-scale cold event in Europe during the Lateglacial
70 (Greenland ice-core records; Rasmussen et al., 2014). The YD is characterized by extreme cold,
71 relative dry and windy climate conditions in northern-central Europe (Hepp et al., 2019).
72 Climate became distinctly warmer at 11,700 cal BP with the onset of the Holocene Interglacial

73 (Rasmussen et al., 2014). These rapid and marked climate oscillations have been observed in
74 the Greenland ice core records (Rasmussen et al., 2014) and in Europe from various proxies
75 such as pollen, oxygen isotopes, molecular biomarkers, beetles, and chironomids (e.g. Ammann
76 et al., 2000; Blaga et al., 2013; Coope and Lemdahl, 1995; Duprat-Oualid et al., 2022; Heiri et
77 al., 2015; Lotter et al., 2012; Millet et al., 2012; Moreno et al., 2014; Peyron et al., 2005; Ponel
78 et al., 2022).

79 Regional climate differences have been evidenced during the Lateglacial, and
80 temperature trends in Europe and the Mediterranean region are still a matter of active research
81 and debate. The chironomid-based synthesis of Heiri et al. (2014) suggests that temperature
82 variations during the Lateglacial tend to be more pronounced in Western Europe (British Isles,
83 Norway) than in Southwestern Europe, Central and Southeastern regions. This is particularly
84 true for the Younger Dryas cooling which is not well evidenced in East and Central Southern
85 Europe (Heiri et al., 2014). These regional differences would be attributed to the changing
86 position of the North Atlantic sea-ice and the Polar Frontal JetStream (Renssen and Isarin,
87 2001).

88 Diverging temperature trends are also reconstructed from different proxies during key
89 periods of the Lateglacial. Studies suggest that (1) the OD was cooler than the YD in Southern
90 and Central Europe in comparison with Northern Europe ($\sim 1-3$ °C; Heiri et al., 2014; Moreno
91 et al., 2014); (2) the Allerød period was warmer than the Bølling in Southwestern Europe and
92 the Mediterranean area (~ 1 °C; Moreno et al., 2014); and (3) temperatures were more contrasted
93 during the B/A and YD in the Northwest of Europe in comparison to the South of Europe
94 (Renssen and Isarin, 2001; Moreno et al., 2014; Heiri et al., 2014). In contrast to temperature,
95 the precipitation signal is poorly known in Europe during the Lateglacial because few proxies
96 are available to quantitatively reconstruct precipitation change. Climate models (GCMs)
97 simulate significant hydrological changes during the B/A and contrasted North-South patterns
98 during the YD (Renssen and Isarin, 2001; Rea et al., 2020). They simulate drier conditions in
99 Northern Europe and wetter conditions in Southern Europe, i.e. in the South of Italy, the Dinaric
100 Alps, and Northern Turkey (Rea et al., 2020). Climate changes during the YD are attributed to
101 a weak Atlantic Meridional Overturning Circulation (AMOC) and a southward shift of the Polar
102 Frontal JetStream (PFJS), linked to the elevation of the ice sheet, in particular the Laurentide
103 ice sheet (Renssen and Isarin, 2001; Renssen et al., 2015; Rea et al., 2020). Rea et al. (2020)
104 also explains the regional climate patterns in Europe by a relocation of Atlantic storm tracks
105 along the western European margin and into the Mediterranean.

106 The understanding of climate processes in Europe and Mediterranean regions during the
107 Lateglacial still needs to be improved. The majority of climate reconstructions are focused on
108 temperatures, and changes in precipitation remain elusive. The “key” junction area between
109 Northern and Southern Europe and regional climatic patterns also needs to be better defined.
110 Moreover, the proxies used to reconstruct climate changes (e.g., coleoptera, chironomids,
111 pollen, ostracods, speleothems) can show differences in terms of amplitudes or patterns which
112 are not only affected by temperatures, but also by precipitation or effective moisture (Moreno
113 et al., 2014; Samartin et al., 2017). For these reasons, more reliable temperature reconstructions,
114 especially from Western Europe and the Mediterranean region are required to test diverging
115 trends during the Lateglacial. The proxies largely used to quantitatively reconstruct past climate
116 changes are often a single proxy approach (e.g. Heiri et al., 2015; Gandouin et al., 2016; Peyron
117 et al., 2017; Marchegiano et al., 2020; Duprat-Oualid et al., 2022). Multiproxy approaches on
118 the same sedimentary record, including independent climate proxies, are necessary to better
119 understand the climate processes in Europe during the Lateglacial (Lotter et al., 2012; Ponel et
120 al., 2022). Pollen-based reconstructions have the advantage of reconstructing temperatures,
121 precipitation, and seasonality, however, the climate signal can be perturbed by other factors
122 such as CO₂ changes and human impact influencing vegetation development (Peyron et al.,
123 2005). Over the last decades, novel proxies based on molecular geochemistry have been
124 developed and molecular biomarkers are being increasingly used to reconstruct temperatures
125 and represent a complementary proxy for lake sediments (Castañeda and Schouten, 2011). In
126 particular, branched Glycerol Dialkyl Glycerol Tetraethers (brGDGTs) are ubiquitous organic
127 compounds synthesized by bacteria (Weijers et al., 2006) which have been useful for
128 reconstructing environmental parameters. To date, the actual producers of brGDGTs remain
129 elusive although it is proposed they come from the phylum *Acidobacteria* (Weijers et al., 2009;
130 Sinninghe Damsté et al., 2018). The relationship, however, between brGDGT distribution and
131 environmental changes, in particular pH and temperature, are well established (Naafs et al.,
132 2017b, 2017a; Dearing Crampton-Flood et al., 2020; Martínez-Sosa et al., 2021; Raberg et al.,
133 2021). The degree of methylation of brGDGTs (MBT; methylation of branched GDGTs) varies
134 depending on the mean annual air temperature (MAAT) and higher fractional abundance of
135 hexa- (III) and penta- (II) methylated brGDGTs are recorded in colder environments (Weijers
136 et al., 2007). Branched glycerol dialkyl glycerol tetraether (brGDGT) membrane lipids are
137 increasingly used as a temperature proxy: in Europe, brGDGTs have been used to reconstruct
138 the Mid to Late Holocene temperature changes in the Carpathians (Ramos-Román et al., 2022),
139 the last 36,000 years in the Southern Iberian Peninsula (Rodrigo-Gámiz et al., 2022), the

140 Holocene temperatures in France (Martin et al., 2020), and in the Eastern Mediterranean over
141 the last deglaciation (Sanchi et al., 2014; Stockhecke et al., 2021). The association in the same
142 core between brGDGTs and other proxies such as pollen for climate reconstructions are still
143 rare (Watson et al., 2018; Panagiotopoulos et al., 2020; Martin et al., 2020; Dugerdil et al.,
144 2021a, 2021b; Ramos-Román et al., 2022; Robles et al., 2022; Rodrigo-Gámiz et al., 2022) and
145 no studies are yet available for the circum-Mediterranean region during the Lateglacial.

146 This study presents a high-resolution climate reconstruction for the Lateglacial period in
147 South Central Europe, inferred from multi-proxy data of the Lake Matese sedimentary record
148 (Southern Italy). In detail, the aims of this study are to:

149 1) establish reliable and independent quantitative climate reconstructions based on
150 molecular biomarkers (brGDGTs) and pollen data to help identify potential biases of currently
151 used proxies and thus improve the reliability of each proxy-inferred climate record;

152 2) compare these reconstructions with regional climate reconstructions and in the light of
153 other South European records;

154 3) better understand the climate processes in Europe and Mediterranean during the
155 Lateglacial period.

156

157 **2. Study site**


158

159 Lake Matese (41°24'33.3"N, 14°24'22.1"E, 1012 m a.s.l.) is located in the Caserta
160 province in the Campania region, Southern Italy, approximately 60 km north of the city of
161 Naples and the active Campanian volcanoes (Vesuvius, Campi Flegrei, Ischia) (Fig. 1). The
162 lake is situated in the Matese karst massif in the Southern Apennines, which extends over 30
163 km from the NE to the SW and is composed of Late Triassic-Miocene limestones and dolomites
164 (Fiorillo and Doglioni, 2010). The present formation of the massif was the result of an extension
165 by strike-slip faults during the Quaternary, and several strong earthquakes were recorded in the
166 massif (Ferranti et al., 2015; Ferrarini et al., 2017; Galli et al., 2017; Valente et al., 2019). Lake
167 Matese is the highest karst lake of Italy and is surrounded by the two highest peaks of the massif,
168 Mount Miletto (2050 m a.s.l.) and Mount Gallinola (1923 m a.s.l.), which feed the lake by their
169 snowmelt. Along the southern side of the lake, two sinkholes named the “Breccie” and
170 “Scennerato” are present (Fiorillo and Pagnozzi, 2015). In the 1920s, hydraulic works were
171 conducted to isolate the bottom of the lake and the main sinkholes by earthen dams (Fiorillo
172 and Pagnozzi, 2015). The water level of the lake improved from 1007-1009 m a.s.l. to 1012 m
173 a.s.l. with a volume of 15 Mm³ (Fiorillo and Pagnozzi, 2015). A part of the lake water is

174 transported to the hydroelectric power station of Piedimonte Matese at the bottom of the
175 mountain massif.

176 The Matese Mountains are characterized by a Mediterranean warm-temperate, humid
177 climate (Aucelli et al., 2013). The southeastern part of the massif, including Lake Matese, have
178 the highest precipitation with a maximum of 2167 mm at Campitello Matese (1400 m a.s.l.)
179 (Fiorillo and Pagnozzi, 2015). Lake Matese shows an annual precipitation of 1808 mm with a
180 maximum in November (~290 mm) and December (~260 mm) and a minimum in July (~50
181 mm) (Fiorillo and Pagnozzi, 2015). The annual temperatures correspond to 9.3°C with a
182 minimum in January (2°C) and a maximum in July (19°C) (Fiorillo and Pagnozzi, 2015).

183 The vegetation of the Matese massif is dominated by deciduous *Quercus* and *Ostrya*
184 *carpinifolia*, while the highest altitudes at the northern flank also show an exposure of *Fagus*
185 *sylvatica* and the lower altitudes of the southern flank includes Mediterranean taxa such as
186 *Quercus ilex* (Taffetani et al., 2012; Carranza et al., 2012; Guarino et al., 2015). The
187 hygrophilous vegetation at Lake Matese is distinguished by the presence of woody (e.g. *Salix*
188 *alba*, *S. caprea*, *S. cinerea* subsp. *cinerea*, *Populus nigra*, *P. alba*), helophytes (e.g. *Phragmites*
189 *australis*, *Schoenoplectus lacustris*, *Typha angustifolia*, *T. latifolia*) and hydrophytes species
190 (*Myriophyllum spicatum*, *Persicaria amphibia*).



191

Figure 1. Location of the Lake Matese and Lateglacial paleoclimate records : Höllloch (Li et al., 2021), Maloja Riegel (Heiri et al., 2014), Lago di Lavarone (Heiri et al., 2014), Lago Piccolo di Avigliana (Larocque and Finsinger, 2008), Lago Gemini (Samartin et al., 2017), Lago Verdarolo (Samartin et al., 2017), Corchia cave (Regattieri et al., 2014), Lake Trasimeno (Marchegiano et al., 2020), Lago Grande di Monticchio (Allen et al., 2002), MD90-917 (Combourieu-Nebout et al., 2013; Sicre et al., 2013), BS7938 (Sbaffi et al., 2004), MD04-2797 (Desprat et al., 2013; Sicre et al., 2013). Dotted line indicates latitude 42°N. Location of active Campanian volcanoes (Vesuvius, Campi Flegrei, Ischia).

192

193 3. Material and methods

194

195 3.1 Coring retrieval

196

Coring of Lake Matese was performed in July 2019 in the southwestern part of the lake
 197 (41°24'33.3"N, 14°24'22.1"E, 1012 m a.s.l.). Core occurred on a floating raft composed of *Salix*
 198 spp. and *Phragmites* spp., naturally present in the eastern part of the lake. Three parallel cores
 199 (cores A, B and C) were taken with a 1 m Russian corer with a chamber diameter of 6.3 cm.
 200 The composite core, measuring 535 cm, was constructed from sections of parallel cores and is
 201 based on the lithology and XRF data.

202

203 3.2 Chronology and age-depth model

204 Several methods have been used to build the chronology of the core including
205 radiocarbon dating, and tephrochronology. The regional pollen stratigraphy is used to validate
206 this age-depth model. Twelve accelerator mass spectrometry (AMS) ¹⁴C dates were measured
207 at Poznań Radiocarbon Laboratory and at the Radiocarbon Dating Center in Lyon. Plant
208 macrofossils (plant fibers, wood) and charcoal were selected for four samples, and bulk
209 sediment was used for eight samples according to the sediment type. Radiocarbon ages were
210 calibrated in years cal BP using the *Calib 8.2* software with the IntCal20 calibration curve
211 (Reimer et al., 2020).

212 Visible tephra layers and cryptotephra layers, detected by magnetic susceptibility and
213 XRF core scanning data, were subsampled and processed for geochemical analysis.
214 Cryptotephra was extracted using H₂O₂ and HCl to remove organic matter and carbonates,
215 sieved at 20 and 100 microns, volcanic glass shards were embedded in resin, sectioned and
216 polished for electron probe microanalysis. A JEOL-JXA8230 probe the Helmholtz Centre
217 Potsdam (Germany) was used with a 15kV accelerating voltage, 10 nA beam current, and a 15
218 micron beam size. Analytical count times were 20 seconds for all elements except for K and
219 Na, measured first at 10 s. International glass standards such as the Max Planck Institute (MPI-
220 glasses) ATHO-G, StHs6/80 and GOR-132 (Jochum et al., 2006) and the natural Lipari obsidian
221 (Hunt and Hill, 1996; Kuehn et al., 2011) were measured prior to sample analysis for data
222 quality insurance. Glass geochemical data of Matese tephtras are normalized on an anhydrous,
223 volatile-free basis and compared with published tephra glass datasets (Wulf et al., 2008; Smith
224 et al., 2011; Tomlinson et al., 2012).

225 The age-depth model based on based on one radiocarbon date and correlated tephra ages
226 was constructed using an interpolated linear curve with the R ‘Clam’ program with 95%
227 confidence intervals (Blaauw, 2010). In order to validate the age depth models, the pollen
228 stratigraphy of the regional sites was compared with pollen data of Matese. The pollen
229 stratigraphy of Pavullo di Frignano (Vescovi et al., 2010), Lakes Accesa (Drescher-Schneider
230 et al., 2007), Albano (Mercuri et al., 2002), Mezzano (Sadori, 2018), Monticchio (Allen et al.,
231 2002), and Trifoglietti (De Beaulieu et al., 2017) were used to identify the OD-B/A, B/A-YD
232 and YD-Holocene transitions. We used the median age for each transition.

233

234 3.3 Magnetic susceptibility and geochemistry

235 Magnetic susceptibility (MS) was measured with a MS2E1 surface scanning sensor
236 from Bartington Instruments on a Geotek Multi-Sensor Core logger based at the Chrono-

237 environment laboratory (UMR CNRS - University of Franche-Comté). An interval of 3 mm or
238 5 mm was applied depending on the type of sediment.

239 Geochemical analyses were performed at high resolution by X-ray Fluorescence (XRF)
240 with an AVAATECH core scanner at the EDYTEM laboratory (University Savoie Mont
241 Blanc). A continuous 5 mm step measurement was applied with a run at 10 kV and 0.1 mA for
242 15 s to detect lightweight elements, such as Al, Si, K, Ca, Ti, Mn, Fe and a second run at 30 kV
243 and 0.15 mA for 20 s to detect Br, Rb, Sr and Zr. The XRF core scanning provides an estimate
244 of the geochemical composition, and the results are semi-quantitative and expressed as peak
245 intensities counts i.e. counts per second (cps).

246

247 3.4 Pollen analyses

248 A total of 56 samples from the Matese core were collected at 4 cm or 6 cm resolution
249 for pollen analysis. For each sample, 1 cm³ of sediment was processed and 3 *Lycopodium* tablets
250 were added to estimate pollen concentration. Samples were treated following the standard
251 procedure (Faegri et al., 1989; Moore et al., 1991) including HCl, KOH, sieving, acetolysis and
252 HF. The pollen concentrates were analyzed with a Leica DM1000 LED microscope at a
253 standard magnification of 400x. Pollen taxa were identified using photo atlases (Beug, 2004;
254 Reille, 1998; Van Geel, 2002) and a modern reference collection (ISEM, University of
255 Montpellier). Each slide was counted with a minimum of 300 terrestrial pollen grains, excluding
256 aquatic plants such as Cyperaceae, aquatic taxa, and fern spores. A simplified pollen diagram
257 was constructed (Fig. 2) with the R package *Rioja* (Juggins and Juggins, 2020). This study
258 presents the main pollen taxa and is not focused on variations of individual species.

259

260 3.5 Pollen-inferred climate reconstruction

261 A multi-method approach was used to reconstruct climate parameters from pollen data
262 with greater reliability than reconstructions based on a single climate reconstruction method
263 (Peyron et al., 2013, 2011, 2005; Salonen et al., 2019). We have selected here the Modern
264 Analog Technique (MAT; Guiot, 1990), Weighted Averaging Partial Least Squares regression
265 (WAPLS; ter Braak and van Dam, 1989; ter Braak and Juggins, 1993), and the most recent
266 machine-learning methods : Random Forest (RF; Breiman, 2001; Prasad et al., 2006) and
267 Boosted Regression Trees (BRT; De'ath, 2007; Elith et al., 2008).

268 The MAT is an assemblage approach, based on the measure of the degree of
269 dissimilarity (squared chord distance) between fossil and modern pollen assemblages (Guiot,
270 1990). Fossil pollen assemblages are compared to a set of modern assemblages (modern

271 dataset), each one associated with climate estimates. The closest modern samples are retained
272 and averaged to estimate past climate conditions (annual and seasonal temperature and
273 precipitation). WAPLS is a non-linear regression technique that models the relationships
274 between the climate parameters and the pollen taxa from a modern pollen dataset, before
275 applying these relationships to fossil pollen assemblages (ter Braak and Juggins, 1993; ter Braak
276 and van Dam, 1989). WAPLS and MAT methods are applied with the R package *Rioja* (Juggins
277 and Juggins, 2020). RF and BRT, based on machine learning, utilizes regression trees
278 developed with ecological data, and has been used recently to reconstruct palaeoclimatic
279 changes (Salonen et al., 2019; Robles et al., 2022). These classification trees are used to
280 partition the data by separating the pollen assemblages based on the relative pollen percentages.
281 RF is based on a large number of regression trees, each tree being estimated from a randomized
282 ensemble of different subsets of the modern pollen dataset by bootstrapping (Breiman, 2001;
283 Prasad et al., 2006). Finally, the RF prediction is applied to the fossil pollen record. BRT is also
284 based on regression trees (De'ath, 2007; Elith et al., 2008); it differs from RF in the definition
285 of the random modern datasets. In RF, each sample gets the same probability of being selected,
286 while in BRT the samples that were insufficiently described in the previous tree get a higher
287 probability of being selected. This approach is called 'boosting' and increases the performance
288 of the model over the elements that are least well predicted (Breiman, 2001; Prasad et al., 2006;
289 De'ath, 2007; Elith et al., 2008). RF is applied with the R package *randomForest* (Liaw and
290 Wiener, 2002) and BRT with the R package *dismo* (Hijmans et al., 2021).

291 The modern pollen dataset (n = 3373 sites) used for the calibration of the methods is
292 based on the large Eurasian/Mediterranean dataset compiled by Peyron et al. (2013, 2017) and
293 completed by Dugerdil et al. (2021a) and Robles et al. (2022). In our study, we added pollen
294 data of 92 surface lake sediments from Italy (Finsinger et al., 2007) and 15 moss polsters from
295 the Matese massif (Robles, 2022). Then, a biome constraint (Guiot et al., 1993), based on the
296 pollen-Plant Functional Type method and following the biomization procedure (Peyron et al.,
297 1998; Prentice et al., 1996) was applied to modern and fossil pollen samples. The modern pollen
298 dataset finally selected for the calibration of the different methods contains 1018 samples
299 belonging to 3 biomes depicted in the fossil core: "warm mixed forest" (WAMX), "temperate
300 deciduous" (TEDE) and "cold steppe" (COST). Performance of each method and calibration
301 training was statistically tested (for more details, see Dugerdil et al., 2021a) to determining if
302 modern samples are suitable for quantitative climate reconstructions. The Root Mean Square
303 Error (RMSE) and the R^2 are presented in the Supplementary Table S1. Five climate parameters
304 were reconstructed, mean annual air temperature (MAAT), mean temperature of the warmest

305 month (MTWA), mean temperature of the coldest month (MTCO), mean annual precipitation
306 (PANN), and winter precipitation (P_{winter} = December, January, and February). For each climate
307 parameter, the methods fitting with the higher R^2 and the lower RMSE were selected.
308 Cyperaceae and ferns in the Matese record have been excluded because they are associated with
309 local dynamics.

310

311 3.6 BrGDGT analyses

312 A total of 56 samples from the Matese core (4 cm or 6 cm resolution) were used for
313 GDGT analysis (same as for pollen analysis). The samples were freeze-dried, powdered and
314 subsampled (1 g for clay and 0.4 g for gyttja). Lipids were extracted from the sediment using a
315 microwave oven (MARS 6; CEM) with dichloromethane:methanol (3:1). Then, the internal
316 standard was added (C_{46} GDGT, Huguet et al., 2006). The total lipid extracts were separated
317 into apolar and polar fractions using a silica SPE cartridge with hexane:DCM (1:1) and
318 DCM:MeOH (1:1). The polar fractions containing brGDGTs were analyzed using a High-
319 Performance Liquid Chromatography Mass Spectrometry (HPLC-APCI-MS, Agilent 1200)
320 with detection via selective ion monitoring (SIM) of m/z 1050, 1048, 1046, 1036, 1034, 1032,
321 1022, 1020, and 1018 in the LGL-TPE of ENS Lyon (Hopmans et al., 2016; Davtian et al.,
322 2018). GDGT concentrations were calculated based on the internal standard (C_{46} GDGT,
323 Huguet et al., 2006). The analytic reproducibility was assessed by regularly processing a lab-
324 internal sediment sample (Vaux Marsh; 45°57'21.1"N, 5°35'32.42"E). Analytical precision is
325 based on duplicate injections of one sample of each Matese core lithological types (n=4).
326 Respective analytical 1-sigma standard deviations are then applied to each measurement within
327 one lithology.

328

329 3.7 GDGTs annual temperature reconstruction

330 The proportion of tetra- (I), penta- (II) and hexa- (III) methylated brGDGTs includes
331 the fractional abundances of the 5-methyl (X), 6-methyl (X') and 7-methyl (X7) brGDGTs
332 (Ding et al., 2016). The CBT (cyclization ratio of branched tetraethers) and MBT indexes were
333 defined by Weijers et al. (2007) and the MBT'_{5me} , only based on the 5-methyl brGDGTs, by
334 De Jonge et al., (2014). The Mean Annual Air Temperature (MAAT) was reconstructed with
335 global (Sun et al., 2011) and East African (Russell et al., 2018) lacustrine calibrations. The
336 mean temperature of Months Above Freezing (MAF) was reconstructed with a lacustrine
337 calibration based on Bayesian statistics (Martínez-Sosa et al., 2021;
338 <https://github.com/jesstierney/BayMBT>) and a global lacustrine calibrations with revised

339 compound fractional abundances based on methylation and cyclization number and methylation
 340 position (Raberg et al., 2021). Synthesis of the formulae for the main brGDGT indices are
 341 presented in Table 1. Modern MAAT and MAF of the Lake Matese corresponds to 9.3 °C.

342 The analytic reproducibility corresponds to ± 0.040 for CBT, ± 0.0167 for MBT, ± 0.0206
 343 for MBT'_{5me} , ± 0.8566 °C for MAAT developed by Sun et al. (2011), ± 0.6672 °C for MAAT
 344 developed by Russell et al. (2018), and ± 0.5403 °C and ± 1.1258 °C for MAF_{Meth} and MAF_{Full}
 345 developed by Raberg et al. (2021).
 346

Table 1. Synthesis of the formulae for the main brGDGT indices. For acronym explanation of MAF_{Meth} and MAF_{Full} , see Raberg et al. (2021). For more information about the Bayesian statistics see Martínez-Sosa et al., 2021 and references therein.

Indice	Formula	Reference
%tetra	$\frac{Ia + Ib + Ic}{\Sigma brGDGTs}$	Ding et al., 2016
%penta	$\frac{IIa + IIa' + IIa_7 + IIb + IIb' + IIb_7 + IIc + IIc' + IIc_7}{\Sigma brGDGTs}$	Ding et al., 2016
%hexa	$\frac{IIIa + IIIa' + IIIa_7 + IIIb + IIIb' + IIIb_7 + IIIc + IIIc' + IIIc_7}{\Sigma brGDGTs}$	Ding et al., 2016
CBT	$-\log \frac{Ib + IIb}{Ia + IIa}$	Weijers et al., 2007
MBT	$\frac{Ia + Ib + Ic}{\Sigma brGDGTs}$	Weijers et al., 2007
MBT'_{5me}	$\frac{Ia + Ib + Ic}{Ia + Ib + Ic + IIa + IIb + IIc + IIIa}$	De Jonge et al., 2014a
MAAT (°C)	$3.949 - 5.593 \times CBT + 38.213 \times MBT$ ($n = 100, R^2 = 0.73, RMSE = 4.27^\circ C$)	Sun et al., 2011
MAAT (°C)	$-1.21 + 32.42 \times MBT'_{5me}$ ($n = 65, R^2 = 0.92, RMSE = 2.44^\circ C$)	Russell et al., 2018
MAF_{Meth} (°C)	$92.9 + 63.84 \times fIb_{Meth}^2 - 130.51 \times fIb_{Meth}$ $- 28.77 \times fIIa_{Meth}^2 - 72.28 \times fIIb_{Meth}^2$ $- 5.88 \times fIIc_{Meth}^2 + 20.89 \times fIIIa_{Meth}^2$ $- 40.54 \times fIIIa_{Meth} - 80.47 \times fIIIb_{Meth}$ ($n = 182, R^2 = 0.90, RMSE = 2.14^\circ C$)	Raberg et al., 2021
MAF_{Full} (°C)	$-8.06 + 37.52 \times fIa_{Full} - 266.83 \times fIb_{Full}^2$ $+ 133.42 \times fIb_{Full} + 100.85 \times fIIa'_{Full}^2$ $+ 58.15 \times fIIIa'_{Full}^2 + 12.79 \times fIIIa_{Full}$	Raberg et al., 2021

($n = 182, R^2 = 0.91, RMSE = 1.97\text{ }^\circ\text{C}$)

Equation from the Bayesian model :

$$MAF\text{ (}^\circ\text{C)} \quad MBT'_{5me} = 0.030(\pm 0.001)MAF + 0.075(\pm 0.012)$$

Martínez-Sosa et al., 2021

$$(R^2 = 0.82, RMSE = 2.9\text{ }^\circ\text{C})$$

347

348

349 4. Results

350


351 4.1 Lithology, magnetic susceptibility, XRF and pollen

352 The lithology of the Matese core (Fig. 2) is mainly composed of gray clay sediment with
353 vivianite from the base to 350 cm, interrupted by an organic layer between 477-484 cm
354 (sedimentary Unit 2) and a macroscopically visible tephra layer (Fig. 2) between 476-437 cm
355 (sedimentary Unit 3). This part contains few plant fibers, which are essentially vertically
356 oriented in the core. From 349 to 320 cm, the lithology is formed by a mix of clay sediment and
357 gyttja (sedimentary Unit 5). This part is mostly composed by roots and fine rootlets.

358 Magnetic susceptibility (MS) and Potassium (K) peaks of XRF core scanning are used
359 to detect tephra layers (Fig. 2). MS and Potassium contents show increased values at 516-502
360 cm, 482-437 cm and 366-338 cm, which correspond to the deposition of tephra material
361 (macroscopic visible tephra and cryptotephra of primary and secondary deposition). Small
362 peaks are also visible in MS between 430 and 360 cm but they are not associated with any
363 observed tephra. Potassium content is also marked by an increase between 536-526 cm which
364 corresponds to tephra of primary deposition. Titanium (Ti) content, on the other hand, is
365 representative for terrigenous input which is prevailing in sedimentary Unit 4 (Fig. 2).

366 The main pollen taxa diagram (Fig. 2) shows the dominance of herbaceous taxa
367 (Poaceae, *Artemisia*) and a small proportion of arboreal taxa at the base of the sequence. From
368 520 to 425 cm, the period is marked by three expansion phases of arboreal taxa, followed
369 between 438 to 354 cm by a large increase of *Artemisia* and a drop of AP taxa starting at 422
370 cm. Finally, from 354 to 338 cm AP and Poaceae increase, whereas *Artemisia* significantly
371 decline.

372



373


Figure 2. Sediment lithology, magnetic susceptibility, geochemical data and selected terrestrial pollen taxa of Matese. Arboreal Pollen (AP; green) and Non Arboreal Pollen (NAP; yellow-orange) are expressed in percentages of total terrestrial pollen.

374 4.2 Age-depth model

375 The age-depth model is based on ^{14}C dates and tephrochronology, and then pollen
 376 stratigraphy was used to validate the age-depth model (Fig. 3). Based on their typical phono-
 377 trachytic and bimodal tephri-phonolitic to trachytic major element glass composition Matese
 378 tephtras at 530 cm and 346 cm depth can be correlated with distal Monticchio tephtras TM-8 and
 379 TM-6-2, respectively (Fig. 4; Table 2). Tephra TM-8 has been correlated with the Neapolitan
 380 Yellow Tuff (NYT) eruption (Wulf et al., 2004) which has an age of $14,194 \pm 172$ cal BP (Bronk
 381 Ramsey et al., 2015). The tephra layer at 530 cm corresponds to the primary deposition and
 382 secondary deposition of remobilised tephtras that were identified at 510 cm and 475 cm. TM-6-
 383 2 most likely are derived from the Early Holocene Casale eruption from Campi Flegrei (Smith
 384 et al., 2011) which is varve dated in Monticchio at $11,210 \pm 224$ cal BP (Wulf et al., 2008). The
 385 tephra layer at 346 cm corresponds to a primary deposition.

386 The ages obtained with the regional pollen stratigraphy show an OD-B/A transition at
 387 $14,500 \pm 93.7$ cal BP, a B/A-YD transition at $12,800 \pm 57.7$ cal BP and a YD-Holocene
 388 transition at $11,575 \pm 103.1$ cal BP (Allen et al., 2002; Mercuri et al., 2002; Drescher-Schneider
 389 et al., 2007; Vescovi et al., 2010; De Beaulieu et al., 2017; Sadori, 2018). Pollen stratigraphy of

390 the regional sites were compared with pollen data of Matese and the ages obtained show a good
 391 correspondence with the ages of tephra samples but a poor correspondence with the ^{14}C dates.
 392 Therefore, most of the ^{14}C dates (Table 3) are not included in the age-depth model (except the
 393 date at the base of the core). The organic matter extracted from sediment was essentially
 394 composed of rootlets, that explains the rejuvenation of the ^{14}C ages.




395

Figure 3. Age-depth model is based on calibrated AMS radiocarbon dates (red points; Table 3) and tephra ages (orange points; Table 2). The grey band is the 95% confidence interval. Blue triangles are the median of ages of the vegetation transition compiled with the regional pollen stratigraphy. This pollen stratigraphy includes the sites of Pavullo di Frignano (Vescovi et al., 2010), Accesa (Drescher-Schneider et al., 2007), Albano (Mercuri et al., 2002), Mezzano (Sadori, 2018), Monticchio (Allen et al., 2002), and Trifoglietti (De Beaulieu et al., 2017). AP/Artemisia ratio (blue line) is expressed on a logarithmic scale. AP: Arboreal Pollen.

Table 2. Tephra samples from Matese cores (MC) and correlation with tephra samples from Lago Grande di Monticchio (Wulf et al., 2008) and proximal eruptive sources.

Sample ID	Depth MC (cm)	Tephra Monticchio	Eruption	Age (cal BP)	Age reference
C1 96-97	346	TM-6-2	Casale	11,210 ± 224	Wulf et al., 2008
A5 75-77	475 (reworked)	TM-8	Neapolitan Yellow Tuff (NYT)	14,194 ± 172	Bronk Ramsey et al., 2015
C3 83-84	510 (reworked)				
B5 95-96	530				

396



397

Figure 4. Bivariate plot of selected major elements (SiO_2 vs. total alkalis and SiO_2 vs. Cl) of Matese tephras and potential proximal and Monticchio tephra correlatives. Data from: TM-6-2 (Monticchio, Wulf et al., 2008; this study); TM-8 (Monticchio, Tomlinson et al., 2012; this study); Casale, Fondi di Baia (proximal; Smith et al., 2011); APP/Agnano Pomice Principali and NYT/Neapolitan Yellow Tuff (proximal; Tomlinson et al., 2012).

Table 3. AMS-radiocarbon dates (Radiocarbon Laboratory, Poznań), calibrated median ages, with 2σ range of calibration from Matese cores (MC).


Sample ID	Depth MC (cm)	Lab code	Material	AMS ^{14}C age (BP)	Age (cal BP) (2σ)	Median age (cal BP)
A4 40-41	340	Poz-128971	Bulk	3425 ± 30	3573 - 3822	3668
A4 60-61	360	Poz-138111	Bulk	7850 ± 40	8540 - 8968	8631
A4 80-81	380	Poz-138112	Bulk	7640 ± 50	8370 - 8541	8432
B4 50-51	400	Poz-128972	Bulk	7580 ± 60	8206 - 8519	8385
A5 20-21	420	Poz-138113	Bulk	7570 ± 50	8206 - 8512	8379
A5 60-61	460	Poz-128976	Bulk	10020 ± 50	11280 - 11743	11519
A6 52-53	479	Poz-119283	Plant fibers, wood fragments, charcoals	6730 ± 40	7513 - 7669	7596
A5 96-97	496	Poz-137155	Wood fragments	10870 ± 60	12728 - 12903	12799
B5 64-65	500	Poz-128973	Bulk	11000 ± 60	12769 - 13078	12925
A6 98-99	525	Poz-119284	Plant fibers	6060 ± 35	6795 - 7147	6912
B5 97-98	533	60747	Plant fibers	5430 ± 30	6190 - 6295	6236
B5 98-99	534	Poz-128975	Bulk	12650 ± 130	14331 - 15477	15027

398

399 4.3 Pollen-inferred climate reconstructions

400 Pollen-inferred climate reconstructions at Matese show similar trends for all methods
401 (Fig. 5). The MAT and the BRT methods show higher sample-to-sample variability than the
402 WAPLS, and RF appears as the less sensitive method. Statistical results of the model
403 performance (Supplementary Table S1) show the better values for R^2 and RMSE for the BRT
404 method (all climatic parameters).

405 Temperature trends show two cold periods (phases 1 and 3) and two warm periods
406 (phases 2 and 4). The reconstructed values (MAAT and MTWA) during the warm periods are
407 close to modern values whereas the values of MTCO are lower than the modern values. Annual
408 precipitation (PANN) shows few variations and the values of PANN and P_{winter} are lower than
409 modern values, with all methods. Phase 1 (535-530 cm; 14,600-14,500 cal BP) is characterized
410 by cold conditions and low precipitation during winter. Phase 2 (530-436 cm; 14,500-12,800
411 cal BP) is a warm period characterized by strong warming and punctuated by three colder events
412 at 14,000 cal BP, 13,500-13,350 cal BP and 13,000 cal BP. Mean annual precipitation shows
413 little variation whereas P_{winter} shows higher values than during the phase 1. Phase 3 (436-367
414 cm; 12,800-11,570 cal BP) is a strong event marked by cold conditions, a slight decline in P_{winter}
415 and few changes for PANN. At the transition with phase 4, a significant decrease in the
416 precipitation parameters is recorded. Phase 4 (367-338 cm; 11,570-11,000 cal BP) is
417 characterized by a well-marked temperature increase (MAAT and MTCO) associated with wet
418 conditions (hydrological parameters reach their maximum value).



422 **Figure 5.** Lake Matese pollen-inferred climate reconstruction based on four methods against age: MAT
423 (Modern Analogue Technique), WAPLS (Weighted Averaging Partial Least Squares regression), RF
424 (Random Forest) and BRT (Boosted Regression Trees). Large lines correspond to loess smoothed
425 curves, shaded areas to the 95% confidence interval and dashed lines to modern climate values of Lake
426 Matese. MAAT: mean annual air temperature. MTWA: mean temperature of the warmest month.
427 MTCO: mean temperature of the coldest month. PANN: mean annual precipitation. P_{winter} : winter
428 precipitation. OD: Oldest Dryas. B/A: Bølling–Allerød. YD: Younger Dryas. EH: Early Holocene.
429


430 4.3 BrGDGT-inferred climate reconstruction

431

432 *4.3.1 Concentration and distribution of brGDGTs*

433 The total concentration of brGDGTs ranges between 0.06 and 8.63 $\mu\text{g}\cdot\text{g}^{-1}$ dry sediment.
434 The fractional abundances of brGDGTs (Fig. 6A) show a dominance of pentamethylated
435 brGDGTs (II, 46%), especially brGDGT IIa (23%), brGDGTs IIa' (7%) and brGDGTs IIb
436 (6%). The relative abundance of tetramethylated brGDGTs (I, 33%) is mainly explained by
437 brGDGT Ia (20%) and brGDGTs Ib (9%). The relative abundance of hexamethylated brGDGTs
438 (III, 21%) is mainly explained by brGDGT IIIa (11%) and brGDGTs IIIa' (6%). The relative
439 abundances of tetra, penta- and hexamethylated brGDGTs of Matese core are compared to
440 global datasets (Fig. 6B). Sediment samples of the Matese core show a good correspondence
441 with global lake and soil samples, except for some samples from sedimentary Unit 1 and 5.
442 Samples of sedimentary Unit 5, characterized by a mix of clay and gyttja, are more similar to
443 global soil and peat samples.

444



445

Figure 6. A) Fractional abundance of tetra-, penta-, and hexamethylated brGDGTs for Matese core. B) Ternary diagram showing the fractional abundances of the tetra-, penta-, and hexamethylated brGDGTs for Matese core (black points) and global lake (blue points; Martínez-Sosa et al., 2021), peat (yellow circles; Naafs et al., 2017a), and soils (gray circles; Yang et al., 2014; Naafs et al., 2017b).

446

447 4.3.2 Indices of brGDGTs

448 The relative abundance of tetra-, penta-, and hexamethylated brGDGTs changes along
 449 Matese core (Fig. 7). The fractional abundance shows a dominance of pentamethylated
 450 brGDGTs except at 518 cm depth, and during the last phase (Phase 4). The fractional abundance
 451 of hexamethylated brGDGTs shows higher values between 535-502 cm and 490-466 cm and
 452 becomes dominant at 486 cm. The fractional abundance of tetramethylated brGDGTs shows
 453 higher values between 502-490 cm and 466-352 cm and is dominant at 518 cm and 352-338
 454 cm (Phase 4).

455 The degree of methylation (MBT, MBT'_{5Me}) and the cyclisation ratio (CBT) also shows
 456 variation along Matese core (Fig. 7). The MBT and the MBT'_{5Me} show similar trends but

457 different absolute values; they vary between 0.17 and 0.52 and between 0.20 and 0.63,
 458 respectively. The degree of methylation remains relatively stable except during two phases of
 459 decrease between 534-522 cm and 486-458 cm, and two phases with higher values at 518 cm
 460 depth and during the Phase 4. The CBT varies between 0.27 and 0.74. Phase 1 (535-530 cm) is
 461 characterized by high values of CBT following by a decline until reaching a minimum between
 462 494-482 cm. Then, the CBT slightly increases; at 382 cm a slow decline is recorded, and a
 463 strong increase marks Phase 4.

464





Figure 7. Fractional abundance of tetra-, penta-, and hexamethylated brGDGTs degree of methylation (MBT, MBT'_{5Me}), cyclisation ratio (CBT) against depth for the Matese core.

482

483 4.3.3 Temperature reconstructions based on brGDGTs

484 The brGDGT inferred reconstructed MAAT using global (Sun et al., 2011) and East
 485 African (Russell et al., 2018) lacustrine calibrations show similar trends than MAF
 486 reconstructed using a Bayesian statistical model (Martínez-Sosa et al., 2021) and global (Raberg
 487 et al., 2021) lacustrine calibrations (Fig. 8). The values are higher than modern values,
 488 especially the values for the MAF_{Full} (Raberg et al., 2021). During Phase 1 (535-530 cm;
 489 14,600-14,500 cal BP), all calibrations show cold temperatures. Phase 2 (530-436 cm; 14,500-
 490 12,800 cal BP) is marked by an abrupt temperature increase or a stabilization for MAF_{Meth} or a
 491 decline for MAF_{Full}. Between 13,700 and 13,200 cal BP, lower temperatures are recorded with
 492 all calibrations and from 13,100 cal BP, temperatures slowly decrease until 11,300 cal BP

493 although a slight increase is recorded between 11,900-11,500 cal BP. Phase 4 (367-338 cm;
 494 11,570-11,000 cal BP) is characterized by a significant increase of temperature.
 495
 496



497
Figure 8. Mean Annual Air Temperature (MAAT) based on global (Sun et al., 2011) and East African (Russell et al., 2018) lacustrine calibrations and Mean temperature of Months Above Freezing (MAF) based on Bayesian statistics (Martínez-Sosa et al., 2021) and global (Raberg et al., 2021) lacustrine calibrations against age for the Matese core. Shaded areas correspond to the error associated with calibrations and dashed lines correspond to modern climate values of Lake Matese. B/A: Bølling–Allerød. YD: Younger Dryas. EH: Early Holocene.

498 **5. Discussion**

499
 500 5.1 Validation of age-depth model

501 The compilation of ages derived from the Italian pollen stratigraphy into the Matese
 502 age-model is based on the main vegetation changes identified in the area during the Lateglacial.
 503 In summary, the OD in Italian pollen records (and in the present study, Fig. 4) is characterized
 504 by an open vegetation dominated by Poaceae, *Artemisia*, with a few arboreal pollen such as
 505 *Pinus* and *Juniperus* appearing (Allen et al., 2002; Vescovi et al., 2010; Drescher-Schneider et
 506 al., 2007; De Beaulieu et al., 2017; Sadori, 2018). During the B/A, a significant increase of
 507 arboreal pollen taxa, including deciduous *Quercus* deciduous, is recorded, and in the majority
 508 of records *Betula* appears (Allen et al., 2002; Drescher-Schneider et al., 2007; Vescovi et al.,
 509 2010; Sadori, 2018; this study). During the YD, an increase of Poaceae and *Artemisia* (Allen et
 510 al., 2002; Mercuri et al., 2002; Drescher-Schneider et al., 2007; Vescovi et al., 2010) and an

511 overall decrease of arboreal pollen taxa, except in Southern Italy, (Allen et al., 2002; Beaulieu
512 et al., 2017; this study) are documented.

513 The ages of tephra samples and ages constrained from the pollen stratigraphy are in
514 good agreement, contrasting results from the ^{14}C dates which are randomly scattered and
515 systematically too young (Fig. 2). The sediments of the Matese core are mainly composed of
516 clay with only few plant fibers. Considering the recurrence of radiocarbon dates between 7570
517 and 7850 cal BP in the core interval between 420 and 360 cm depth (see Table 1), it is
518 hypothesized that the dated organic matter may have partly originated from penetrating rootlets
519 of plants growing during sedimentary Unit 5's deposition (Fig. 4). Indeed, aquatic plants of
520 sedimentary Unit 5, identified with pollen, evidence a shallow water body and the development
521 of tree species that typically grow in wetland.

522 Therefore, the overall age-depth model of the Matese core is based on imported, well-
523 accepted tephra ages and one ^{14}C date of a bulk sediment sample from the bottom of the core
524 at 534 cm (Fig. 2).

525

526 5.2 Influence of proxies and methods on climate reconstructions

527

528 *5.2.1 Lake Matese climate signal reliability*

529 Climate reconstructions are based both on pollen and brGDGTs, and some temperature
530 discrepancies (absolute values or amplitudes) are depicted depending on the proxies (Fig. 9).
531 The temperature amplitudes and absolute values are higher for brGDGTs (5-20°C) than the
532 pollen (4-10°C) reconstructions. Pollen-inferred temperature values depend heavily on the
533 quality of the modern pollen dataset including the number of samples, the diversity of samples
534 in terms of biomes, and the similarity with the fossil samples (Chevalier et al., 2020). In our
535 study, the modern database includes several modern samples from the Matese massif, and 95
536 samples from Italy were added to complete the dataset. Moreover, the spatial autocorrelation is
537 low for MAT (Moran's $I < 0.34$, $p\text{-value} < 0.01$), and climate trends are consistent between
538 methods. Reconstructed values for temperatures are close to modern values during the warmest
539 periods, however, precipitation is largely underestimated by all methods for the recent time
540 period (Fig. 5). The same observation was made in Calabria in Southern Italy (Trifoglietti;
541 Joannin et al., 2012), a region also characterized by precipitation above 1700 mm. The
542 underestimation of precipitation is certainly linked to the lack of modern samples located in
543 very wet Mediterranean areas. Considering the brGDGT climate signal, the reconstructed
544 temperatures are overestimated in comparison with modern values (Fig. 8). For shallow

545 temperate lakes (< 20 m), like Lake Matese, our brGDGT reconstructions suggest values
546 anomalously higher than the expected temperature due to thermal variability (seasonal and
547 diurnal; Martínez-Sosa et al., 2021). Lake Matese is located at an altitude of 1012 m a.s.l. and
548 the strong seasonal variability may have influenced the brGDGT distribution. Moreover, the
549 Lake Matese climate reconstructions are based on several global lacustrine calibration datasets,
550 which may not be well adapted to reconstruct paleotemperatures in the Mediterranean region.
551 According to Dugerdil et al. (2021a), local calibrations perform better to reconstruct more
552 reliable absolute values. Unfortunately, at date, only a few global lacustrine calibrations are
553 available, and a local calibration dataset for the Mediterranean region is still missing.

554

555 *5.2.2 Regional climate signal reliability depending on the proxy*

556 Climate reconstructions inferred from Lake Matese are compared to key terrestrial and
557 marine temperature and precipitation records (Fig. 9, 10) in a latitudinal transect in central
558 Mediterranean. These reconstructions for the Mediterranean region are based on different
559 proxies. Most of those are indicators of annual temperatures, but some of them are indicators
560 of seasonal temperature changes. For example, transfer functions based on chironomid
561 assemblages provide estimates of mean July air temperatures (Larocque and Finsinger, 2008;
562 Heiri et al., 2014; Samartin et al., 2017), while ostracod assemblages allow quantitative
563 reconstruction of both January and July palaeotemperatures (Marchegiano et al., 2020).
564 Planktonic foraminifera provide estimates of spring and autumn sea surface temperatures (SST)
565 (Sicre et al., 2013). Depending on the production and deposition settings, molecular biomarkers
566 are considered as indicators of annual or seasonal temperatures like brGDGTs or alkenones
567 (Sbaffi et al., 2004; Sicre et al., 2013; Zhang et al., 2013; Max et al., 2020; Martínez-Sosa et
568 al., 2021; this study). For precipitation (Fig. 10), fewer reconstructions are available and they
569 are mainly based on records of pollen (Combourieu-Nebout et al., 2013), $\delta^{18}\text{O}$ *G. bulloides* in
570 marine sediments (Sicre et al., 2013), and $\delta^{18}\text{O}$ in speleothems (Regattieri et al., 2014). Pollen
571 enable the reconstruction of both annual and seasonal temperatures and precipitation (e.g. Allen
572 et al., 2002; Tarroso et al., 2016).

573 The comparison between climate reconstructions inferred from different proxies allows
574 us to identify reliable regional climate signals and to reduce the bias linked to each proxy.
575 Indeed, differences may appear for the timing or amplitudes of changes according to the type
576 of proxy. These differences may be amplified by the proxy provenance, either marine or
577 continental. In Figure 9, the temperature reconstructions above 42°N are mainly based on
578 chironomids, and the climate signal reconstructed is consistent between the sites. In South Italy,

579 at Monticchio, climate reconstructions are based on three pollen records from the same site and
580 differences in terms of amplitude and trend are clearly evidenced (Fig. 9I). These differences
581 are linked to the differences in the core location in the lake and the pollen sample resolution
582 (Allen et al., 2002). The closer the core to the center of the lake (dark blue, Fig. 9I), the better
583 the regional vegetation record and therewith a possible regional climate signal (Peyron et al.,
584 2005). Between latitude 41°N and 36°N, sea-surface temperatures (SSTs) were reconstructed
585 from foraminifera and/or alkenones analyzed from marine cores (Sbaffi et al., 2004; Sicre et
586 al., 2013). Alkenone-based SSTs show a low amplitude of 2-3°C between the B/A and the YD
587 periods, whereas foraminifera-based reconstruction of seasonal temperature show differences
588 of 5-10°C between the B/A and the YD. The differences are linked to their respective methods:
589 For alkenones, the estimation of SSTs are based on the molecular biomarker as the C₃₇ alkenone
590 unsaturation ($U_{37}^{K'}$), whereas, for foraminifera, they are calculated with the MAT method and
591 depend on the occurrence of modern analogues (Sicre et al., 2013).

592

593 5.3 Climate changes during the Lateglacial in Italy

594

595 *5.3.1 Bølling–Allerød warming*

596 The age of transition between the OD and the Bølling–Allerød Interstadial is estimated
597 at around 14,700 cal BP based on the NGRIP ice-core chronology (Rasmussen et al., 2014). In
598 Italy, an abrupt warming is evidenced at ca 14,700 cal BP (Fig. 9). The differences between the
599 different reconstructions seem related to the type of proxy used rather than latitude. The
600 transition is not obvious in the temperature reconstructions based on alkenones (Fig. 9MO;
601 Sbaffi et al., 2004; Sicre et al., 2013), whereas it is well marked in reconstructions based on
602 foraminifera (Fig. 9N; Sicre et al., 2013) and pollen assemblages (Desprat et al., 2013) from
603 the same cores. According to Sicre et al. (2013), alkenones-inferred SSTs could be biased
604 during the Early deglaciation due to water stratification inducing warming of the thin surface
605 water layers where small size nanophytoplankton grow. Except for temperature reconstructions
606 based on alkenones, all the records show an increase of the temperature at the transition OD-
607 B/A (Larocque and Finsinger, 2008; Sicre et al., 2013; Heiri et al., 2014; Marchegiano et al.,
608 2020). The transition, although marked, seems more progressive in the Italian records than in
609 Greenland ice-core but the low resolution of some records can favor this trend. In terms of
610 precipitation (Fig. 10), few records are available in Italy but no significant changes are recorded
611 around 14,700 cal BP by $\delta^{18}\text{O}$ *G. bulloides* (Sicre et al., 2013) and pollen transfer functions
612 (Desprat et al., 2013; this study).

613 The Bølling–Allerød interstadial is a warm interstadial period interrupted by several
614 cold climate oscillations (Rasmussen et al., 2014). According to the synthesis by Moreno et al.
615 (2014), the Bølling was cooler than the Allerød in the Southern Mediterranean compared to the
616 warmer Northern Mediterranean. In Italy, above 42°N, temperature trends are complex to
617 interpret: some records show an increase of temperature (Fig. 9B; Heiri et al., 2014) whereas
618 other records show a decline (Fig. 9CE; Larocque and Finsinger, 2008; Marchegiano et al.,
619 2020). At Matese, pollen and brGDGTs inferred temperatures decrease (Fig. 9F-H), whereas in
620 the southern part of Italy, there are no significant changes during the B/A (Fig. 9I-O; Allen et
621 al., 2002; Saffi et al., 2004; Sicre et al., 2013). Temperature reconstructions in Italy show no
622 distinct difference between the Bølling and the Allerød with respect to the latitude. In terms of
623 amplitude, several studies (Renssen and Isarin, 2001; Heiri et al., 2014; Moreno et al., 2014)
624 suggests that there were less contrasts in temperatures during the B/A in Southern Europe in
625 comparison with Northern Europe. Once again, this difference is not clear in Italy (Fig. 9). At
626 Matese, a significant decrease of brGDGTs-inferred temperature is recorded at 13,700-13,200
627 cal BP (Fig. 9H). This change could be attributed to a colder period such as the Older
628 Dryas or the Inter-Allerød cold period, two short periods characterized by colder conditions in
629 the Greenland ice-core records at 14,000 and 13,100 cal BP, respectively (Rasmussen et al.,
630 2014). However, this cooling event do not appear at the same time in the Matese climate curve
631 based on pollen, and it is only vaguely recorded in other Italian records (Fig. 9). We suggest
632 that this change could be attributed to changes of local conditions that are visible in a lithology
633 change (sedimentary Unit 2, Fig. 4). Indeed, brGDGT distribution and origin can differ
634 according to the type of wetland, water level or vegetation changes (Martínez-Sosa et al., 2021;
635 Robles et al., 2022). In terms of precipitation (Fig. 10), no significant changes occur during the
636 B/A in Italy as suggested previously by Renssen and Isarin (2001) for Southern Europe. The
637 Alpine region seems instead to record wetter conditions during the B/A (Barton et al., 2018; Li
638 et al., 2021).

639

640 *5.3.2 A marked Younger Dryas cold event throughout Italy*

641 The onset of the YD is estimated around 12,900 cal BP according to the Greenland ice-
642 core chronology (Rasmussen et al., 2014). In Italy, above 42°N, the transition between the B/A
643 and the YD is progressive in terms of temperatures except for chironomid records (Fig. 9B;
644 Heiri et al., 2014). At Matese, pollen-based reconstructions show a progressive decline of
645 temperatures with all methods except the MAT (Fig. 9FG). For this method, the transition is
646 more abrupt, but this difference can be attributed to the application of the biome constraint.

647 BrGDGT-based reconstructions record a steady decrease during the YD or no significant
648 changes according to the calibrations used (Fig. 9H). For southern Italian records, the transition
649 is more abrupt and particularly marked in the foraminifera record in contrast to alkenones-based
650 reconstructions (Fig. 9J-O; Saffi et al., 2004; Sicre et al., 2013). In terms of precipitation (Fig.
651 10), the northern Italian speleothems records show an abrupt transition (Regattieri et al., 2014;
652 Li et al., 2021) whereas the southern Italian pollen and isotopes records do not reveal significant
653 changes (Sicre et al., 2013; Combourieu-Nebout et al., 2013; Desprat et al., 2013).

654 The YD is characterized by cold conditions in the Northern Hemisphere from 12,900 to
655 11,700 cal BP (Rasmussen et al., 2014). As previously mentioned for the B/A, several studies
656 (Renssen and Isarin, 2001; Heiri et al., 2014; Moreno et al., 2014) suggest that temperatures
657 during the YD are less contrasted in the South of Europe in comparison with the North. In Italy
658 as a whole (Fig. 9), a decline in temperatures is recorded in all records.

659 At Matese, a decrease of temperatures is evidenced by the pollen-based reconstructions,
660 but it is less clear from the brGDGT-based reconstructions. The difference of climate signals
661 may be related to different sources between both proxies. Pollen record local, extra-local and
662 regional vegetation (Jacobson and Bradshaw, 1981). The basin size of the Lake Matese is larger
663 than 5 hectares, which suggest a signal of regional vegetation rather than local (Jacobson and
664 Bradshaw, 1981). Moreover, the YD is marked by a large proportion of herbaceous taxa (Fig.
665 4) and favors the catching of regional pollen (Jacobson and Bradshaw, 1981). By contrast,
666 brGDGTs are produced in the lake or in the catchment area (Russell et al., 2018; Martin et al.,
667 2019) and thus are local contributors. Moreover, the YD is characterized by high erosion rates
668 in the catchment (Fig. 4), which could favor greater soil-derived brGDGTs and induce a warm
669 bias in temperatures (Martínez-Sosa et al., 2021). Indeed, the distribution of brGDGTs differ
670 according to sample type and could differ between lake sediments and catchment soils (Loomis
671 et al., 2011, 2014; Buckles et al., 2014; Russell et al., 2018; Martin et al., 2019; Martínez-Sosa
672 et al., 2021; Raberg et al., 2022). Soil sediments generally exhibit less hexamethylated
673 brGDGTs and more tetramethylated brGDGTs than lake sediments (Loomis et al., 2011, 2014;
674 Buckles et al., 2014; Russell et al., 2018; Martin et al., 2019; Martínez-Sosa et al., 2021).
675 However, an increase of tetramethylated brGDGTs is mainly associated with an increase in
676 temperatures in soils and lake sediments (Russell et al., 2018). At Matese, the YD is
677 characterized by a decrease in hexamethylated brGDGTs and a slight increase in
678 tetramethylated brGDGTs. These differences may have affected the annual temperature
679 reconstructions by inducing a warm bias in temperatures during the YD. Furthermore, soil-
680 derived brGDGTs may also be affected by changes in pH, moisture, soil compounds and

681 vegetation in the catchment of Lake Matese (Davtian et al., 2016; Martin et al., 2019; Liang et
682 al., 2019; Dugerdil et al., 2021a). Furthermore soil samples without vegetation cover are more
683 sensitive to seasonal changes than that of soil samples with grass and forest cover (Liang et al.,
684 2019). Therefore, soils with vegetation cover allow a better reconstruction of global
685 temperatures (Liang et al., 2019). Since at Matese, the YD is characterized by an open
686 vegetation, soil-derived brGDGTs could also have been affected by seasonal temperature
687 changes due to a sparse vegetation and this effect is superimposed to changes in the sources of
688 brGDGTs in lake sediments.

689 Contrasted patterns are also recorded at Monticchio (Fig. 9I) by the three different
690 climate variables used for pollen-based temperature reconstructions: a decrease in winter
691 temperature is reconstructed for two lake cores, while a fen core external to the lake, which
692 should record the local vegetation signal, does not reveal the temperature decline during the YD
693 (Allen et al., 2002). However, the two other cores clearly show a temperature decrease, that is
694 why we consider a winter temperature decrease during the YD at Monticchio. In Southern
695 Italian records, temperature reconstructions based on alkenones, foraminifera and pollen (Sbaffi
696 et al., 2004; Desprat et al., 2013; Sicre et al., 2013) show a shorter YD than in the north. For
697 alkenones-based reconstructions, even an increase of temperatures is recorded at the end of the
698 YD. In continental records of South Italy (Allen et al., 2002), this trend is only recorded at
699 Monticchio (one core only) and does not appear at Matese. Nonetheless, this hypothesis is only
700 based on marine records and should be investigated through continental records in Southern
701 Italy.

702 In terms of precipitation, the marine sequences located south of latitude 42°N record a
703 slight increase for proxies based on pollen (Fig. 9GH; Combourieu-Nebout et al., 2013) and on
704 $\delta^{18}\text{O}$ *G. bulloides* data (Fig. 9FI; Sicre et al., 2013) during the YD. However, no significant
705 change occurs at Matese for PANN (Fig. 10C), and on the contrary a low decline is recorded
706 for P_{winter} towards the end of the YD (Fig. 10D). Above latitude 42°N, a precipitation decrease
707 during the YD is recorded by two sites at Hölloch and Corchia caves (Fig. 10AB; Regattieri et
708 al., 2014; Li et al., 2021). According to the model outputs of Rea et al. (2020), drier conditions
709 occurred in Northern Europe whereas wetter conditions prevailed in Southern Europe, mainly
710 during winter and in the South of Italy, the Dinaric Alps and Northern Turkey. This pattern is
711 consistent with our reconstruction but the limit between the North and the South is closer to
712 latitude 42°N.

713 The transition between the YD and the Holocene is recorded around 11,700 cal BP by
714 Greenland ice-core records (Rasmussen et al., 2014). In Italy, an important increase of

715 temperature is recorded in all records (Fig. 9) which appears earlier (700-400 years) in southern
716 sites (Sbaffi et al., 2004; Sicre et al., 2013). In terms of precipitation, marine records south of
717 latitude 42°N continue to record a slight increase of precipitation (Fig. 10E-H; Combourieu-
718 Nebout et al., 2013; Sicre et al., 2013), and in northern sites an increase of precipitation is
719 recorded (Fig. 10A-D; Regattieri et al., 2014; Li et al., 2021; this study).

720

721 5.4 Atmospheric processes during the Lateglacial in central Mediterranean

722 According to several studies, climate changes during the Lateglacial show differences
723 in temperatures between Southern and Central Europe (Heiri et al., 2014; Moreno et al., 2014;
724 Renssen and Isarin, 2001). In Italy (Fig. 9), climate reconstructions do not show latitudinal
725 differences in terms of temperature. The B/A is marked by warm conditions and the YD by cold
726 conditions even in Southern Italy. Climate reconstructions for East-Central Southern Europe
727 from Heiri et al., (2014) are not consistent with our results probably because while two of their
728 chironomid records are located in North Italy and one in Bulgaria none consider Southern Italy.
729 In the study of Moreno et al. (2014), only the record of Monticchio is used for the South of Italy
730 during the Lateglacial, which may explain the differences in our study. Considering
731 precipitation, several studies suggest no significant changes during the B/A but drier conditions
732 in Northern Europe and wetter conditions in Southern Europe during the YD. In Italy (Fig. 10),
733 we observe the same dynamics during the B/A and the YD.

734 Several studies (Renssen and Isarin, 2001; Moreno et al., 2014; Rea et al., 2020) explain
735 that during cold periods of the Lateglacial (OD, YD) the Polar Frontal JetStream moved
736 southward with a weak Atlantic Meridional Overturning Circulation (AMOC) (Moreno et al.,
737 2014; Rea et al., 2020; Renssen and Isarin, 2001). The incursion of cold air masses is recorded
738 until the South of Italy, however, during the YD, dry conditions are not reconstructed for this
739 region. According to Rea et al. (2020), a relocation of Atlantic storm tracks in the Mediterranean
740 is induced by the Fennoscandian ice sheet and the North European Plain which created a
741 topographic barrier and a high pressure region during the YD. The presence of Atlantic storm
742 tracks into the Mediterranean could have favored wetter conditions in the South of Italy during
743 the YD. Our study suggests a limit around latitude 42°N, with drier conditions in Northern Italy
744 and slightly wetter conditions in Southern Italy during the YD. A latitude limit at 40°N was
745 previously discussed by Magny et al. (2013) for the Holocene. These echoing limits over time
746 in Italy inevitably reinforce Italy's key position to archive proxies catching atmospheric
747 patterns.

748 By contrast, during the B/A, the North Atlantic sea-ice has a more northerly position
749 inducing a northward shift of the Polar Frontal JetStream (Renssen and Isarin, 2001). The
750 incursion of warm air masses is recorded in all of Italy, however, no significant changes in
751 annual precipitation occur. Our study does not suggest the location of Atlantic storm tracks in
752 Italy during the B/A, although at Matese winter precipitation was higher in most pollen-based
753 climate reconstructions. However, very few records and climatic models reconstructing
754 precipitation are available in Europe and the Mediterranean region for this period. Further
755 investigations are necessary to fully understand the atmospheric processes and precipitation
756 dynamic in Europe, mainly during the B/A.





Figure 9. Synthesis of temperature records inferred from different proxies in Italy from 15,000 to 11,000 cal BP and comparison with the NGRIP ice core record. MAAT: mean annual air temperature. MTWA: mean temperature of the warmest month. MTCO: mean temperature of the coldest month. OD: Oldest Dryas. B/A: Bølling–Allerød. YD: Younger Dryas. EH: Early Holocene.

758



759

Figure 10. Synthesis of precipitation records inferred from different proxies in Italy 15,000 to 11,000 cal BP. PANN: mean annual precipitation. P_{winter}: winter precipitation. OD: Oldest Dryas. B/A: Bølling–Allerød. YD: Younger Dryas. EH: Early Holocene.

760

761 **6. Conclusions**

762 This study provides a quantitative climate reconstruction for the Lateglacial period in
763 Central-Southern Europe, inferred from a multi-proxy and multi-method approach based on the
764 Lake Matese record. The comparison of the Lake Matese climate reconstructions based on
765 brGDGTs and pollen and their comparison with regional terrestrial/marine climate
766 reconstructions show the following:

- 767 • For the first time, pollen and brGDGTs were combined to reconstruct climate
768 changes in the Mediterranean region during the Lateglacial. Temperature trends
769 reconstructed with these proxies are consistent except during the YD. Both proxies
770 show a marked cold OD, an increase of temperatures during the B/A, and an abrupt
771 transition to warmer conditions for the Holocene. During the YD, pollen-based
772 reconstructions show a decrease of temperatures, whereas brGDGT-based
773 reconstructions show no significant changes.
- 774 • Comparison with regional climate records of Italy reveals that there are no
775 latitudinal differences during the B/A and the YD in terms of temperatures. The B/A
776 is marked by an increase of temperature and the YD is characterized by cold
777 conditions in all Italy. By contrast, precipitation does not show changes during the
778 B/A, and a slight increase of precipitation during the YD is recorded in Southern
779 Italy below latitude 42°N.
- 780 • Cold conditions during the YD in Italy may be linked to the southward position of
781 North Atlantic sea-ice and of the Polar Frontal JetStream. The low increase of
782 precipitation during the YD may be linked to relocation of Atlantic storm tracks into
783 the Mediterranean, induced by the Fennoscandian ice sheet and the North European
784 Plain. We identified the latitude 42°N as a limit between dry conditions in northern
785 Italy and slightly wetter conditions in Southern Italy during the YD. By contrast,
786 warm conditions during the B/A may be linked to the northward position of North
787 Atlantic sea-ice and of the Polar Frontal JetStream.

788
789 In summary, this study allowed us to document and discuss past climate changes in Italy
790 while contributing to the debate about the atmospheric processes in Southern Europe. The
791 latitudes 40-42°N appear as a key junction point between wetter conditions in Southern Italy
792 and drier conditions in Northern Italy during the YD but also during the Early-Mid Holocene
793 (Magny et al., 2013). However, further robust paleoclimate studies are needed to provide 1)
794 high-resolution reconstructions based on several proxies in Northern Italy, 2) new records for

795 central Italy (between 41-43°N), 3) new continental records for Southern Italy (below 41°N)
796 and 4) more model outputs at regional scales with transient simulations, if possible, mainly
797 during the B/A and the YD.

798

799 **Author contribution**

800 MR: Conceptualization, Field work, Laboratory work, Formal analysis, Writing draft
801 manuscript, Review, Funding acquisition. SJ, OP and EB: Conceptualization, Field work,
802 Supervision, Review, Funding acquisition. GM: Conceptualization, Supervision, Review,
803 Funding acquisition. SW: Laboratory work, Formal analysis, Review, Funding acquisition. OA
804 and MB: Laboratory work. BV: Supervision of laboratory work, Review. BP: Field work. SAA:
805 Coordination of laboratory work. LC and SG: Conceptualization, Review. J-LB, LD and AC:
806 Review.

807

808 **Declaration of competing interest**

809 The authors declare that they have no known competing financial interests or personal
810 relationships that could have appeared to influence the work reported in this paper.

811

812 **Funding**

813 This research was co-founded by the International PhD course “Agriculture
814 Technologies and Biotechnologies” (34° Cycle, Code: DOT1339335). Financial support for
815 this study was provided by EFFICACE project from EC2CO INSU CNRS (PI: Odile Peyron)
816 and ERJ ClimMatese from LabEx CeMEB (PI: Mary Robles). The travels between Italy and
817 France were financed by VINCI founding of the Università Italo Francese (UIF). The
818 conference funding was provided by the Association des Palynologues de Langue Française
819 (APLF).

820

821 **Acknowledgements**

822 The authors would like to thank Julien Didier for magnetic susceptibility measurements,
823 Laurent Bouby and Isabelle Figueral for seed and wood identifications used for radiocarbon
824 dating. The authors would like to express their appreciation to Gwenaël Magne, Thierry Pastor
825 and Benoît Brossier for logistical support during fieldwork, and Anne-Lise Develle and Claire
826 Blanchet for help during XRF analysis. We would like to thank Sandrine Canal and Sylvie

827 Rouland for support during pollen sample preparation. This is an ISEM contribution N° 2022-
828 266.

829

830 **References**

- 831 Allen, J.R.M., Watts, W.A., McGee, E., Huntley, B., 2002. Holocene environmental variability-
832 the record from Lago Grande di Monticchio, Italy. *Quaternary International* 88, 69–80.
- 833 Ammann, B., Birks, H.J.B., Brooks, S.J., Eicher, U., von Grafenstein, U., Hofmann, W.,
834 Lemdahl, G., Schwander, J., Tobolski, K., Wick, L., 2000. Quantification of biotic
835 responses to rapid climatic changes around the Younger Dryas — a synthesis.
836 *Palaeogeography, Palaeoclimatology, Palaeoecology* 159, 313–347.
837 [https://doi.org/10.1016/S0031-0182\(00\)00092-4](https://doi.org/10.1016/S0031-0182(00)00092-4)
- 838 Aucelli, P.P.C., Cesarano, M., Di Paola, G., Filocamo, F., Roszkopf, C.M., 2013.
839 Geomorphological map of the central sector of the Matese Mountains (Southern Italy):
840 an example of complex landscape evolution in a Mediterranean mountain environment.
841 *Journal of Maps* 9, 604–616. <https://doi.org/10.1080/17445647.2013.840054>
- 842 Barton, C.M., Aura Tortosa, J.E., Garcia-Puchol, O., Riel-Salvatore, J.G., Gauthier, N., Vadillo
843 Conesa, M., Pothier Bouchard, G., 2018. Risk and resilience in the late glacial: A case
844 study from the western Mediterranean. *Quaternary Science Reviews* 184, 68–84.
845 <https://doi.org/10.1016/j.quascirev.2017.09.015>
- 846 Beug, H.-J., 2004. Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete.
847 Friedrich Pfeil, München.
- 848 Blaauw, M., 2010. Methods and code for ‘classical’ age-modelling of radiocarbon sequences.
849 *Quaternary Geochronology* 5, 512–518. <https://doi.org/10.1016/j.quageo.2010.01.002>
- 850 Blaga, C.I., Reichert, G.-J., Lotter, A.F., Anselmetti, F.S., Sinninghe Damsté, J.S., 2013. A
851 TEX₈₆ lake record suggests simultaneous shifts in temperature in Central Europe and
852 Greenland during the last deglaciation: A SWISS TEX₈₆ LAKE RECORD. *Geophys.*
853 *Res. Lett.* 40, 948–953. <https://doi.org/10.1002/grl.50181>
- 854 Breiman, L., 2001. Random Forests. *Machine Learning* 45, 5–32.
855 <https://doi.org/10.1023/A:1010933404324>
- 856 Bronk Ramsey, C., Albert, P.G., Blockley, S.P.E., Hardiman, M., Housley, R.A., Lane, C.S.,
857 Lee, S., Matthews, I.P., Smith, V.C., Lowe, J.J., 2015. Improved age estimates for key
858 Late Quaternary European tephra horizons in the RESET lattice. *Quaternary Science*
859 *Reviews* 118, 18–32. <https://doi.org/10.1016/j.quascirev.2014.11.007>
- 860 Buckles, L.K., Weijers, J.W.H., Verschuren, D., Sinninghe Damsté, J.S., 2014. Sources of core
861 and intact branched tetraether membrane lipids in the lacustrine environment: Anatomy
862 of Lake Challa and its catchment, equatorial East Africa. *Geochimica et Cosmochimica*
863 *Acta* 140, 106–126. <https://doi.org/10.1016/j.gca.2014.04.042>
- 864 Carranza, M.L., Frate, L., Paura, B., 2012. Structure, ecology and plant richness patterns in
865 fragmented beech forests. *Plant Ecology & Diversity* 5, 541–551.
866 <https://doi.org/10.1080/17550874.2012.740509>
- 867 Castañeda, I.S., Schouten, S., 2011. A review of molecular organic proxies for examining
868 modern and ancient lacustrine environments. *Quaternary Science Reviews* 30, 2851–
869 2891. <https://doi.org/10.1016/j.quascirev.2011.07.009>
- 870 Chevalier, M., Davis, B.A.S., Heiri, O., Seppä, H., Chase, B.M., Gajewski, K., Lacourse, T.,
871 Telford, R.J., Finsinger, W., Guiot, J., Kühl, N., Maezumi, S.Y., Tipton, J.R., Carter,
872 V.A., Brussel, T., Phelps, L.N., Dawson, A., Zanon, M., Vallé, F., Nolan, C., Mauri, A.,
873 de Vernal, A., Izumi, K., Holmström, L., Marsicek, J., Goring, S., Sommer, P.S.,
874 Chaput, M., Kupriyanov, D., 2020. Pollen-based climate reconstruction techniques for

875 late Quaternary studies. *Earth-Science Reviews* 210, 103384.
876 <https://doi.org/10.1016/j.earscirev.2020.103384>

877 Combourieu-Nebout, N., Peyron, O., Bout-Roumazielles, V., Goring, S., Dormoy, I., Joannin,
878 S., Sadori, L., Siani, G., Magny, M., 2013. Holocene vegetation and climate changes in
879 the central Mediterranean inferred from a high-resolution marine pollen record (Adriatic
880 Sea). *Clim. Past* 9, 2023–2042. <https://doi.org/10.5194/cp-9-2023-2013>

881 Coope, G.R., Lemdahl, G., 1995. Regional differences in the Lateglacial climate of northern
882 Europe based on coleopteran analysis. *Journal of Quaternary Science* 10, 391–395.
883 <https://doi.org/10.1002/jqs.3390100409>

884 Davtian, N., Bard, E., Ménot, G., Fagault, Y., 2018. The importance of mass accuracy in
885 selected ion monitoring analysis of branched and isoprenoid tetraethers. *Organic*
886 *Geochemistry* 118, 58–62. <https://doi.org/10.1016/j.orggeochem.2018.01.007>

887 Davtian, N., Ménot, G., Bard, E., Poulenard, J., Podwojewski, P., 2016. Consideration of soil
888 types for the calibration of molecular proxies for soil pH and temperature using global
889 soil datasets and Vietnamese soil profiles. *Organic Geochemistry* 101, 140–153.
890 <https://doi.org/10.1016/j.orggeochem.2016.09.002>

891 De Beaulieu, J.-L., Brugiapaglia, E., Joannin, S., Guiter, F., Zanchetta, G., Wulf, S., Peyron,
892 O., Bernardo, L., Didier, J., Stock, A., Rius, D., Magny, M., 2017. Lateglacial-Holocene
893 abrupt vegetation changes at Lago Trifoglietti in Calabria, Southern Italy: The setting
894 of ecosystems in a refugial zone. *Quaternary Science Reviews* 158, 44–57.
895 <https://doi.org/10.1016/j.quascirev.2016.12.013>

896 De Jonge, C., Hopmans, E.C., Zell, C.I., Kim, J.-H., Schouten, S., Sinninghe Damsté, J.S.,
897 2014. Occurrence and abundance of 6-methyl branched glycerol dialkyl glycerol
898 tetraethers in soils: Implications for palaeoclimate reconstruction. *Geochimica et*
899 *Cosmochimica Acta* 141, 97–112. <https://doi.org/10.1016/j.gca.2014.06.013>

900 Dearing Crampton-Flood, E., Tierney, J.E., Peterse, F., Kirkels, F.M.S.A., Sinninghe Damsté,
901 J.S., 2020. BayMBT: A Bayesian calibration model for branched glycerol dialkyl
902 glycerol tetraethers in soils and peats. *Geochimica et Cosmochimica Acta* 268, 142–
903 159. <https://doi.org/10.1016/j.gca.2019.09.043>

904 De'ath, G., 2007. Boosted trees for ecological modeling and prediction. *Ecology* 88, 243–251.
905 [https://doi.org/10.1890/0012-9658\(2007\)88\[243:BTFEMA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2007)88[243:BTFEMA]2.0.CO;2)

906 Desprat, S., Combourieu-Nebout, N., Essallami, L., Sicre, M.A., Dormoy, I., Peyron, O., Siani,
907 G., Bout Roumazielles, V., Turon, J.L., 2013. Deglacial and Holocene vegetation and
908 climatic changes in the southern Central Mediterranean from a direct land–sea
909 correlation. *Clim. Past* 9, 767–787. <https://doi.org/10.5194/cp-9-767-2013>

910 Ding, S., Schwab, V.F., Ueberschaar, N., Roth, V.-N., Lange, M., Xu, Y., Gleixner, G., Pohnert,
911 G., 2016. Identification of novel 7-methyl and cyclopentanyl branched glycerol dialkyl
912 glycerol tetraethers in lake sediments. *Organic Geochemistry* 102, 52–58.
913 <https://doi.org/10.1016/j.orggeochem.2016.09.009>

914 Drescher-Schneider, R., de Beaulieu, J.-L., Magny, M., Walter-Simonnet, A.-V., Bossuet, G.,
915 Millet, L., Brugiapaglia, E., Drescher, A., 2007. Vegetation history, climate and human
916 impact over the last 15,000 years at Lago dell'Accesa (Tuscany, Central Italy). *Veget*
917 *Hist Archaeobot* 16, 279–299. <https://doi.org/10.1007/s00334-006-0089-z>

918 Dugerdil, L., Joannin, S., Peyron, O., Jouffroy-Bapicot, I., Vannièrè, B., Boldgiv, B.,
919 Unkelbach, J., Behling, H., Ménot, G., 2021a. Climate reconstructions based on GDGT
920 and pollen surface datasets from Mongolia and Baikal area: calibrations and
921 applicability to extremely cold–dry environments over the Late Holocene. *Clim. Past*
922 17, 1199–1226. <https://doi.org/10.5194/cp-17-1199-2021>

923 Dugerdil, L., Ménot, G., Peyron, O., Jouffroy-Bapicot, I., Ansanay-Alex, S., Antheaume, I.,
924 Behling, H., Boldgiv, B., Develle, A.-L., Grossi, V., Magail, J., Makou, M., Robles, M.,

- 925 Unkelbach, J., Vanni re, B., Joannin, S., 2021b. Late Holocene Mongolian climate and
926 environment reconstructions from brGDGTs, NPPs and pollen transfer functions for
927 Lake Ayrag: Paleoclimate implications for Arid Central Asia. *Quaternary Science*
928 *Reviews* 273, 107235. <https://doi.org/10.1016/j.quascirev.2021.107235>
- 929 Duprat-Oualid, F., B geot, C., Peyron, O., Rius, D., Millet, L., Magny, M., 2022. High-
930 frequency vegetation and climatic changes during the Lateglacial inferred from the
931 Lapsou pollen record (Cantal, southern Massif Central, France). *Quaternary*
932 *International* S1040618222001537. <https://doi.org/10.1016/j.quaint.2022.04.012>
- 933 Elith, J., Leathwick, J.R., Hastie, T., 2008. A working guide to boosted regression trees. *J Anim*
934 *Ecology* 77, 802–813. <https://doi.org/10.1111/j.1365-2656.2008.01390.x>
- 935 Faegri, K., Kaland, P.E., Krzywinski, K., 1989. *Textbook of pollen analysis*. John Wiley &
936 Sons, Chichester.
- 937 Ferranti, L., Milano, G., Burrato, P., Palano, M., Cannav , F., 2015. The seismogenic structure
938 of the 2013–2014 Matese seismic sequence, Southern Italy: implication for the
939 geometry of the Apennines active extensional belt. *Geophysical Journal International*
940 201, 823–837. <https://doi.org/10.1093/gji/ggv053>
- 941 Ferrarini, F., Boncio, P., de Nardis, R., Pappone, G., Cesarano, M., Aucelli, P.P.C., Lavecchia,
942 G., 2017. Segmentation pattern and structural complexities in seismogenic extensional
943 settings: The North Matese Fault System (Central Italy). *Journal of Structural Geology*
944 95, 93–112. <https://doi.org/10.1016/j.jsg.2016.11.006>
- 945 Finsinger, W., Heiri, O., Valsecchi, V., Tinner, W., Lotter, A.F., 2007. Modern pollen
946 assemblages as climate indicators in southern Europe. *Global Ecology and*
947 *Biogeography* 16, 567–582. <https://doi.org/10.1111/j.1466-8238.2007.00313.x>
- 948 Fiorillo, F., Doglioni, A., 2010. The relation between karst spring discharge and rainfall by
949 cross-correlation analysis (Campania, southern Italy). *Hydrogeol J* 18, 1881–1895.
950 <https://doi.org/10.1007/s10040-010-0666-1>
- 951 Fiorillo, F., Pagnozzi, M., 2015. Recharge processes of Matese karst massif (southern Italy).
952 *Environ Earth Sci* 74, 7557–7570. <https://doi.org/10.1007/s12665-015-4678-y>
- 953 Galli, P., Giaccio, B., Messina, P., Peronace, E., Amato, V., Naso, G., Nomade, S., Pereira, A.,
954 Piscitelli, S., Bellanova, J., Billi, A., Blamart, D., Galderisi, A., Giocoli, A., Stabile, T.,
955 Thil, F., 2017. Middle to Late Pleistocene activity of the northern Matese fault system
956 (southern Apennines, Italy). *Tectonophysics* 699, 61–81.
957 <https://doi.org/10.1016/j.tecto.2017.01.007>
- 958 Gandouin, E., Rioual, P., Pailles, C., Brooks, S.J., Ponel, P., Guiter, F., Djamali, M., Andrieu-
959 Ponel, V., Birks, H.J.B., Leydet, M., Belkacem, D., Haas, J.N., Van der Putten, N., de
960 Beaulieu, J.L., 2016. Environmental and climate reconstruction of the late-glacial-
961 Holocene transition from a lake sediment sequence in Aubrac, French Massif Central:
962 Chironomid and diatom evidence. *Palaeogeography, Palaeoclimatology, Palaeoecology*
963 461, 292–309. <https://doi.org/10.1016/j.palaeo.2016.08.039>
- 964 Guarino, R., Bazan, G., Paura, B., 2015. Downy-Oak Woods of Italy: Phytogeographical
965 Remarks on a Controversial Taxonomic and Ecologic Issue, in: Box, E.O., Fujiwara, K.
966 (Eds.), *Warm-Temperate Deciduous Forests around the Northern Hemisphere*,
967 *Geobotany Studies*. Springer International Publishing, Cham, pp. 139–151.
968 https://doi.org/10.1007/978-3-319-01261-2_7
- 969 Guiot, J., 1990. Methodology of the last climatic cycle reconstruction in France from pollen
970 data. *Palaeogeography, Palaeoclimatology, Palaeoecology, Methods for the Study of*
971 *Stratigraphical Records* 80, 49–69. [https://doi.org/10.1016/0031-0182\(90\)90033-4](https://doi.org/10.1016/0031-0182(90)90033-4)
- 972 Guiot, J., de Beaulieu, J.L., Cheddadi, R., David, F., Ponel, P., Reille, M., 1993. The climate in
973 Western Europe during the last Glacial/Interglacial cycle derived from pollen and insect

974 remains. *Palaeogeography, Palaeoclimatology, Palaeoecology* 103, 73–93.
975 [https://doi.org/10.1016/0031-0182\(93\)90053-L](https://doi.org/10.1016/0031-0182(93)90053-L)

976 Heiri, O., Brooks, S.J., Renssen, H., Bedford, A., Hazekamp, M., Ilyashuk, B., Jeffers, E.S.,
977 Lang, B., Kirilova, E., Kuiper, S., Millet, L., Samartin, S., Toth, M., Verbruggen, F.,
978 Watson, J.E., van Asch, N., Lammertsma, E., Amon, L., Birks, H.H., Birks, H.J.B.,
979 Mortensen, M.F., Hoek, W.Z., Magyari, E., Muñoz Sobrino, C., Seppä, H., Tinner, W.,
980 Tonkov, S., Veski, S., Lotter, A.F., 2014. Validation of climate model-inferred regional
981 temperature change for late-glacial Europe. *Nat Commun* 5, 4914.
982 <https://doi.org/10.1038/ncomms5914>

983 Heiri, O., Ilyashuk, B., Millet, L., Samartin, S., Lotter, A.F., 2015. Stacking of discontinuous
984 regional palaeoclimate records: Chironomid-based summer temperatures from the
985 Alpine region. *The Holocene* 25, 137–149. <https://doi.org/10.1177/0959683614556382>

986 Hepp, J., Wüthrich, L., Bromm, T., Bliedtner, M., Schäfer, I.K., Glaser, B., Rozanski, K.,
987 Sirocko, F., Zech, R., Zech, M., 2019. How dry was the Younger Dryas? Evidence from
988 a coupled $\delta^2\text{H}$ – $\delta^{18}\text{O}$ biomarker paleohygrometer applied to the Gemündener Maar
989 sediments, Western Eifel, Germany. *Climate of the Past* 15, 713–733.
990 <https://doi.org/10.5194/cp-15-713-2019>

991 Hijmans, R.J., Phillips, S., Elith, J.L. and J., 2021. *dismo: Species Distribution Modeling.*

992 Hopmans, E.C., Schouten, S., Sinninghe Damsté, J.S., 2016. The effect of improved
993 chromatography on GDGT-based palaeoproxies. *Organic Geochemistry* 93, 1–6.
994 <https://doi.org/10.1016/j.orggeochem.2015.12.006>

995 Huguet, C., Hopmans, E.C., Febo-Ayala, W., Thompson, D.H., Sinninghe Damsté, J.S.,
996 Schouten, S., 2006. An improved method to determine the absolute abundance of
997 glycerol dibiphytanyl glycerol tetraether lipids. *Organic Geochemistry* 37, 1036–1041.
998 <https://doi.org/10.1016/j.orggeochem.2006.05.008>

999 Hunt, J.B., Hill, P.G., 1996. An inter-laboratory comparison of the electron probe microanalysis
1000 of glass geochemistry. *Quaternary International* 34–36, 229–241.
1001 [https://doi.org/10.1016/1040-6182\(95\)00088-7](https://doi.org/10.1016/1040-6182(95)00088-7)

1002 Jacobson, G.L., Bradshaw, R.H.W., 1981. The Selection of Sites for Paleovegetational Studies.
1003 *Quat. res.* 16, 80–96. [https://doi.org/10.1016/0033-5894\(81\)90129-0](https://doi.org/10.1016/0033-5894(81)90129-0)

1004 Joannin, S., Brugiapaglia, E., Vanniere, B., 2012. Pollen-based reconstruction of Holocene
1005 vegetation and climate in southern Italy: the case of Lago Trifoglietti. *Clim. Past* 24.

1006 Jochum, K.P., Stoll, B., Herwig, K., Willbold, M., Hofmann, A.W., Amini, M., Aarburg, S.,
1007 Abouchami, W., Hellebrand, E., Mocek, B., Raczek, I., Stracke, A., Alard, O., Bouman,
1008 C., Becker, S., Dücking, M., Brätz, H., Klemd, R., de Bruin, D., Canil, D., Cornell, D.,
1009 de Hoog, C.-J., Dalpé, C., Danyushevsky, L., Eisenhauer, A., Gao, Y., Snow, J.E.,
1010 Groschopf, N., Günther, D., Latkoczy, C., Guillong, M., Hauri, E.H., Höfer, H.E.,
1011 Lahaye, Y., Horz, K., Jacob, D.E., Kasemann, S.A., Kent, A.J.R., Ludwig, T., Zack, T.,
1012 Mason, P.R.D., Meixner, A., Rosner, M., Misawa, K., Nash, B.P., Pfänder, J., Premo,
1013 W.R., Sun, W.D., Tiepolo, M., Vannucci, R., Vennemann, T., Wayne, D., Woodhead,
1014 J.D., 2006. MPI-DING reference glasses for in situ microanalysis: New reference values
1015 for element concentrations and isotope ratios: MPI-DING REFERENCE GLASSES.
1016 *Geochem. Geophys. Geosyst.* 7, n/a-n/a. <https://doi.org/10.1029/2005GC001060>

1017 Juggins, S., Juggins, M.S., 2020. Package ‘rioja.’

1018 Kuehn, S.C., Froese, D.G., Shane, P.A.R., 2011. The INTAV intercomparison of electron-beam
1019 microanalysis of glass by tephrochronology laboratories: Results and recommendations.
1020 *Quaternary International, Enhancing tephrochronology and its application (INTREPID*
1021 *Project): Hiroshi Machida commemorative volume* 246, 19–47.
1022 <https://doi.org/10.1016/j.quaint.2011.08.022>

- 1023 Larocque, I., Finsinger, W., 2008. Late-glacial chironomid-based temperature reconstructions
1024 for Lago Piccolo di Avigliana in the southwestern Alps (Italy). *Palaeogeography,*
1025 *Palaeoclimatology,* *Palaeoecology* 257, 207–223.
1026 <https://doi.org/10.1016/j.palaeo.2007.10.021>
- 1027 Li, H., Spötl, C., Cheng, H., 2021. A high-resolution speleothem proxy record of the Late
1028 Glacial in the European Alps: extending the NALPS19 record until the beginning of the
1029 Holocene. *J. Quaternary Sci* 36, 29–39. <https://doi.org/10.1002/jqs.3255>
- 1030 Liang, J., Russell, J.M., Xie, H., Lupien, R.L., Si, G., Wang, J., Hou, J., Zhang, G., 2019.
1031 Vegetation effects on temperature calibrations of branched glycerol dialkyl glycerol
1032 tetraether (brGDGTs) in soils. *Organic Geochemistry* 127, 1–11.
1033 <https://doi.org/10.1016/j.orggeochem.2018.10.010>
- 1034 Liaw, A., Wiener, M., 2002. Classification and Regression by randomForest 2, 5.
- 1035 Loomis, S.E., Russell, J.M., Heureux, A.M., D'Andrea, W.J., Sinninghe Damsté, J.S., 2014.
1036 Seasonal variability of branched glycerol dialkyl glycerol tetraethers (brGDGTs) in a
1037 temperate lake system. *Geochimica et Cosmochimica Acta* 144, 173–187.
1038 <https://doi.org/10.1016/j.gca.2014.08.027>
- 1039 Loomis, S.E., Russell, J.M., Sinninghe Damsté, J.S., 2011. Distributions of branched GDGTs
1040 in soils and lake sediments from western Uganda: Implications for a lacustrine
1041 paleothermometer. *Organic Geochemistry* 42, 739–751.
1042 <https://doi.org/10.1016/j.orggeochem.2011.06.004>
- 1043 Lotter, A.F., Heiri, O., Brooks, S., van Leeuwen, J.F.N., Eicher, U., Ammann, B., 2012. Rapid
1044 summer temperature changes during Termination 1a: high-resolution multi-proxy
1045 climate reconstructions from Gerzensee (Switzerland). *Quaternary Science Reviews,*
1046 *The INTegration of Ice core, Marine and TERrestrial records of the last termination*
1047 *(INTIMATE) 60,000 to 8000 BP* 36, 103–113.
1048 <https://doi.org/10.1016/j.quascirev.2010.06.022>
- 1049 Magny, M., Combourieu-Nebout, N., de Beaulieu, J.-L., Bout-Roumazeilles, V., Colombaroli,
1050 D., Desprat, S., Francke, A., Joannin, S., Ortu, E., Peyron, O., Revel, M., Sadori, L.,
1051 Siani, G., Sicre, M.A., Samartin, S., Simonneau, A., Tinner, W., Vanniere, B., Wagner,
1052 B., Zanchetta, G., Anselmetti, F., Brugiapaglia, E., Chapron, E., Debret, M., Didier, J.,
1053 Essallami, L., Galop, D., Gilli, A., Kallel, N., Millet, L., Stock, A., Turon, J.L., Wirth,
1054 S., 2013. North-south palaeohydrological contrasts in the central Mediterranean during
1055 the Holocene: tentative synthesis and working hypotheses. *Clim. Past* 30.
- 1056 Marchegiano, M., Horne, D.J., Gliozzi, E., Francke, A., Wagner, B., Ariztegui, D., 2020. Rapid
1057 Late Pleistocene climate change reconstructed from a lacustrine ostracod record in
1058 central Italy (Lake Trasimeno, Umbria). *Boreas* 49, 739–750.
1059 <https://doi.org/10.1111/bor.12450>
- 1060 Martin, C., Ménot, G., Thouveny, N., Davtian, N., Andrieu-Ponel, V., Reille, M., Bard, E.,
1061 2019. Impact of human activities and vegetation changes on the tetraether sources in
1062 Lake St Front (Massif Central, France). *Organic Geochemistry* 135, 38–52.
- 1063 Martin, C., Ménot, G., Thouveny, N., Peyron, O., Andrieu-Ponel, V., Montade, V., Davtian,
1064 N., Reille, M., Bard, E., 2020. Early Holocene Thermal Maximum recorded by branched
1065 tetraethers and pollen in Western Europe (Massif Central, France). *Quaternary Science*
1066 *Reviews* 228, 106109. <https://doi.org/10.1016/j.quascirev.2019.106109>
- 1067 Martínez-Sosa, P., Tierney, J.E., Stefanescu, I.C., Dearing Crampton-Flood, E., Shuman, B.N.,
1068 Routson, C., 2021. A global Bayesian temperature calibration for lacustrine brGDGTs.
1069 *Geochimica et Cosmochimica Acta* 305, 87–105.
1070 <https://doi.org/10.1016/j.gca.2021.04.038>
- 1071 Max, L., Lembke-Jene, L., Zou, J., Shi, X., Tiedemann, R., 2020. Evaluation of reconstructed
1072 sea surface temperatures based on U37k' from sediment surface samples of the North

1073 Pacific. Quaternary Science Reviews 243, 106496.
1074 <https://doi.org/10.1016/j.quascirev.2020.106496>

1075 Mercuri, A.M., Accorsi, C.A., Bandini Mazzanti, M., 2002. The long history of Cannabis and
1076 its cultivation by the Romans in central Italy, shown by pollen records from Lago
1077 Albano and Lago di Nemi. *Veget Hist Archaeobot* 11, 263–276.
1078 <https://doi.org/10.1007/s003340200039>

1079 Millet, L., Rius, D., Galop, D., Heiri, O., Brooks, S.J., 2012. Chironomid-based reconstruction
1080 of Lateglacial summer temperatures from the Ech palaeolake record (French western
1081 Pyrenees). *Palaeogeography, Palaeoclimatology, Palaeoecology* 315–316, 86–99.
1082 <https://doi.org/10.1016/j.palaeo.2011.11.014>

1083 Moore, P.D., Webb, J.A., Collinson, M.E., 1991. *Pollen Analysis*, Subsequent edition. ed.
1084 Blackwell Science Inc, Oxford.

1085 Moreno, A., Svensson, A., Brooks, S.J., Connor, S., Engels, S., Fletcher, W., Genty, D., Heiri,
1086 O., Labuhn, I., Perşoiu, A., Peyron, O., Sadori, L., Valero-Garcés, B., Wulf, S.,
1087 Zanchetta, G., 2014. A compilation of Western European terrestrial records 60–8 ka BP:
1088 towards an understanding of latitudinal climatic gradients. *Quaternary Science Reviews*
1089 106, 167–185. <https://doi.org/10.1016/j.quascirev.2014.06.030>

1090 Naafs, B.D.A., Gallego-Sala, A.V., Inglis, G.N., Pancost, R.D., 2017a. Refining the global
1091 branched glycerol dialkyl glycerol tetraether (brGDGT) soil temperature calibration.
1092 *Organic Geochemistry* 106, 48–56. <https://doi.org/10.1016/j.orggeochem.2017.01.009>

1093 Naafs, B.D.A., Inglis, G.N., Zheng, Y., Amesbury, M.J., Biester, H., Bindler, R., Blewett, J.,
1094 Burrows, M.A., del Castillo Torres, D., Chambers, F.M., Cohen, A.D., Evershed, R.P.,
1095 Feakins, S.J., Gałka, M., Gallego-Sala, A., Gandois, L., Gray, D.M., Hatcher, P.G.,
1096 Honorio Coronado, E.N., Hughes, P.D.M., Huguet, A., Könönen, M., Laggoun-
1097 Défarge, F., Lähteenoja, O., Lamentowicz, M., Marchant, R., McClymont, E.,
1098 Pontevedra-Pombal, X., Ponton, C., Pourmand, A., Rizzuti, A.M., Rochefort, L.,
1099 Schellekens, J., De Vleeschouwer, F., Pancost, R.D., 2017b. Introducing global peat-
1100 specific temperature and pH calibrations based on brGDGT bacterial lipids. *Geochimica*
1101 *et Cosmochimica Acta* 208, 285–301. <https://doi.org/10.1016/j.gca.2017.01.038>

1102 Panagiotopoulos, K., Holtvoeth, J., Kouli, K., Marinova, E., Francke, A., Cvetkoska, A.,
1103 Jovanovska, E., Lacey, J.H., Lyons, E.T., Buckel, C., Bertini, A., Donders, T., Just, J.,
1104 Leicher, N., Leng, M.J., Melles, M., Pancost, R.D., Sadori, L., Tauber, P., Vogel, H.,
1105 Wagner, B., Wilke, T., 2020. Insights into the evolution of the young Lake Ohrid
1106 ecosystem and vegetation succession from a southern European refugium during the
1107 Early Pleistocene. *Quaternary Science Reviews* 227, 106044.
1108 <https://doi.org/10.1016/j.quascirev.2019.106044>

1109 Peyron, O., Bégeot, C., Brewer, S., Heiri, O., Magny, M., Millet, L., Ruffaldi, P., Van Campo,
1110 E., Yu, G., 2005. Late-Glacial climatic changes in Eastern France (Lake Lautrey) from
1111 pollen, lake-levels, and chironomids. *Quat. res.* 64, 197–211.
1112 <https://doi.org/10.1016/j.yqres.2005.01.006>

1113 Peyron, O., Combourieu-Nebout, N., Brayshaw, D., Goring, S., Andrieu-Ponel, V., Desprat, S.,
1114 Fletcher, W., Gambin, B., Ioakim, C., Joannin, S., Kotthoff, U., Kouli, K., Montade, V.,
1115 Pross, J., Sadori, L., Magny, M., 2017. Precipitation changes in the Mediterranean basin
1116 during the Holocene from terrestrial and marine pollen records: a model–data
1117 comparison. *Clim. Past* 13, 249–265. <https://doi.org/10.5194/cp-13-249-2017>

1118 Peyron, O., Goring, S., Dormoy, I., Kotthoff, U., Pross, J., de Beaulieu, J.-L., Drescher-
1119 Schneider, R., Vannièrè, B., Magny, M., 2011. Holocene seasonality changes in the
1120 central Mediterranean region reconstructed from the pollen sequences of Lake Accesa
1121 (Italy) and Tenaghi Philippon (Greece). *The Holocene* 21, 131–146.
1122 <https://doi.org/10.1177/0959683610384162>

- 1123 Peyron, O., Guiot, J., Cheddadi, R., Tarasov, P., Reille, M., de Beaulieu, J.-L., Bottema, S.,
 1124 Andrieu, V., 1998. Climatic Reconstruction in Europe for 18,000 YR B.P. from Pollen
 1125 Data. *Quat. res.* 49, 183–196. <https://doi.org/10.1006/qres.1997.1961>
- 1126 Peyron, O., Magny, M., Goring, S., Joannin, S., de Beaulieu, J.-L., Brugiapaglia, E., Sadori, L.,
 1127 Garfi, G., Kouli, K., Ioakim, C., Combourieu-Nebout, N., 2013. Contrasting patterns of
 1128 climatic changes during the Holocene across the Italian Peninsula reconstructed from
 1129 pollen data. *Clim. Past* 9, 1233–1252. <https://doi.org/10.5194/cp-9-1233-2013>
- 1130 Ponel, P., Guiter, F., Gandouin, E., Peyron, O., de Beaulieu, J.-L., 2022. Late-Glacial
 1131 palaeotemperatures and palaeoprecipitations in the Aubrac Mountains (French Massif
 1132 Central) reconstructed from multiproxy analyses (Coleoptera, chironomids and pollen).
 1133 *Quaternary International*. <https://doi.org/10.1016/j.quaint.2022.02.005>
- 1134 Prasad, A.M., Iverson, L.R., Liaw, A., 2006. Newer Classification and Regression Tree
 1135 Techniques: Bagging and Random Forests for Ecological Prediction. *Ecosystems* 9,
 1136 181–199. <https://doi.org/10.1007/s10021-005-0054-1>
- 1137 Prentice, C., Guiot, J., Huntley, B., Jolly, D., Cheddadi, R., 1996. Reconstructing biomes from
 1138 palaeoecological data: a general method and its application to European pollen data at
 1139 0 and 6 ka. *Climate Dynamics* 12, 185–194. <https://doi.org/10.1007/BF00211617>
- 1140 Raberg, J.H., Flores, E., Crump, S.E., de Wet, G., Dildar, N., Miller, G.H., Geirsdóttir, Á.,
 1141 Sepúlveda, J., 2022. Intact Polar brGDGTs in Arctic Lake Catchments: Implications for
 1142 Lipid Sources and Paleoclimate Applications. *Journal of Geophysical Research:*
 1143 *Biogeosciences* 127, e2022JG006969. <https://doi.org/10.1029/2022JG006969>
- 1144 Raberg, J.H., Harning, D.J., Crump, S.E., de Wet, G., Blumm, A., Kopf, S., Geirsdóttir, Á.,
 1145 Miller, G.H., Sepúlveda, J., 2021. Revised fractional abundances and warm-season
 1146 temperatures substantially improve brGDGT calibrations in lake sediments.
 1147 *Biogeosciences* 18, 3579–3603. <https://doi.org/10.5194/bg-2021-16>
- 1148 Ramos-Román, M.J., De Jonge, C., Magyari, E., Veres, D., Ilvonen, L., Develle, A.-L., Seppä,
 1149 H., 2022. Lipid biomarker (brGDGT)- and pollen-based reconstruction of temperature
 1150 change during the Middle to Late Holocene transition in the Carpathians. *Global and*
 1151 *Planetary Change* 215, 103859. <https://doi.org/10.1016/j.gloplacha.2022.103859>
- 1152 Rasmussen, S.O., Bigler, M., Blockley, S.P., Blunier, T., Buchardt, S.L., Clausen, H.B.,
 1153 Cvijanovic, I., Dahl-Jensen, D., Johnsen, S.J., Fischer, H., Gkinis, V., Guillevic, M.,
 1154 Hoek, W.Z., Lowe, J.J., Pedro, J.B., Popp, T., Seierstad, I.K., Steffensen, J.P., Svensson,
 1155 A.M., Vallenga, P., Vinther, B.M., Walker, M.J.C., Wheatley, J.J., Winstrup, M.,
 1156 2014. A stratigraphic framework for abrupt climatic changes during the Last Glacial
 1157 period based on three synchronized Greenland ice-core records: refining and extending
 1158 the INTIMATE event stratigraphy. *Quaternary Science Reviews* 106, 14–28.
 1159 <https://doi.org/10.1016/j.quascirev.2014.09.007>
- 1160 Rea, B.R., Pellitero, R., Spagnolo, M., Hughes, P., Ivy-Ochs, S., Renssen, H., Ribolini, A.,
 1161 Bakke, J., Lukas, S., Braithwaite, R.J., 2020. Atmospheric circulation over Europe
 1162 during the Younger Dryas. *Sci. Adv.* 6, eaba4844.
 1163 <https://doi.org/10.1126/sciadv.aba4844>
- 1164 Regattieri, E., Zanchetta, G., Drysdale, R.N., Isola, I., Hellstrom, J.C., Dallai, L., 2014.
 1165 Lateglacial to Holocene trace element record (Ba, Mg, Sr) from Corchia Cave (Apuan
 1166 Alps, central Italy): paleoenvironmental implications: Trace element record from
 1167 Corchia Cave, central Italy. *J. Quaternary Sci.* 29, 381–392.
 1168 <https://doi.org/10.1002/jqs.2712>
- 1169 Rehfeld, K., Münch, T., Ho, S.L., Laepple, T., 2018. Global patterns of declining temperature
 1170 variability from the Last Glacial Maximum to the Holocene. *Nature* 554, 356–359.
 1171 <https://doi.org/10.1038/nature25454>

1172 Reille, M., 1998. Reille, Maurice, 1995. Pollen et spores d'Europe et d'Afrique du Nord,
1173 Supplément 1 . Éditions du Laboratoire de botanique historique et palynologie,
1174 Marseille, 327 p., 800 FF. / Reille, Maurice, 1998. Pollen et spores d'Europe et
1175 d'Afrique du Nord, Supplément 2 . Éditions du Laboratoire de botanique historique et
1176 palynologie, Marseille, 530 p., 1600 FF. gpq 52, 0–0. <https://doi.org/10.7202/004885ar>

1177 Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Ramsey, C.B., Butzin, M.,
1178 Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I.,
1179 Heaton, T.J., Hogg, A.G., Hughen, K.A., Kromer, B., Manning, S.W., Muscheler, R.,
1180 Palmer, J.G., Pearson, C., Plicht, J. van der, Reimer, R.W., Richards, D.A., Scott, E.M.,
1181 Southon, J.R., Turney, C.S.M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M.,
1182 Fahrni, S.M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F.,
1183 Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., Talamo, S., 2020. The IntCal20
1184 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP). *Radiocarbon*
1185 62, 725–757. <https://doi.org/10.1017/RDC.2020.41>

1186 Renssen, H., Isarin, R.F.B., 2001. The two major warming phases of the last deglaciation at
1187 ~14.7 and ~11.5 ka cal BP in Europe: climate reconstructions and AGCM experiments.
1188 *Global and Planetary Change* 30, 117–153. [https://doi.org/10.1016/S0921-](https://doi.org/10.1016/S0921-8181(01)00082-0)
1189 [8181\(01\)00082-0](https://doi.org/10.1016/S0921-8181(01)00082-0)

1190 Renssen, H., Mairesse, A., Goosse, H., Mathiot, P., Heiri, O., Roche, D.M., Nisancioglu, K.H.,
1191 Valdes, P.J., 2015. Multiple causes of the Younger Dryas cold period. *Nature Geosci* 8,
1192 946–949. <https://doi.org/10.1038/ngeo2557>

1193 Robles, M., 2022. Vegetation, climate, and human history of the Mediterranean basin: A Late-
1194 Glacial to Holocene reconstruction from Italy (Lake Matese) to Armenia (Lake Sevan)
1195 inferred from a multi-proxy approach (pollen, NPPs, brGDGTs, XRF) (PhD thesis).
1196 University of Molise, University of Montpellier, Campobasso, Montpellier.

1197 Robles, M., Peyron, O., Brugiapaglia, E., Ménot, G., Dugerdil, L., Ollivier, V., Ansanay-Alex,
1198 S., Develle, A.-L., Tozalakyan, P., Meliksetian, K., Sahakyan, K., Sahakyan, L., Perello,
1199 B., Badalyan, R., Colombié, C., Joannin, S., 2022. Impact of climate changes on
1200 vegetation and human societies during the Holocene in the South Caucasus (Vanevan,
1201 Armenia): A multiproxy approach including pollen, NPPs and brGDGTs. *Quaternary*
1202 *Science Reviews* 277, 107297. <https://doi.org/10.1016/j.quascirev.2021.107297>

1203 Rodrigo-Gámiz, M., García-Alix, A., Jiménez-Moreno, G., Ramos-Román, M.J., Camuera, J.,
1204 Toney, J.L., Sachse, D., Anderson, R.S., Sinninghe Damsté, J.S., 2022. Paleoclimate
1205 reconstruction of the last 36 kyr based on branched glycerol dialkyl glycerol tetraethers
1206 in the Padul palaeolake record (Sierra Nevada, southern Iberian Peninsula). *Quaternary*
1207 *Science Reviews* 281, 107434. <https://doi.org/10.1016/j.quascirev.2022.107434>

1208 Russell, J.M., Hopmans, E.C., Loomis, S.E., Liang, J., Sinninghe Damsté, J.S., 2018.
1209 Distributions of 5- and 6-methyl branched glycerol dialkyl glycerol tetraethers
1210 (brGDGTs) in East African lake sediment: Effects of temperature, pH, and new
1211 lacustrine paleotemperature calibrations. *Organic Geochemistry* 117, 56–69.
1212 <https://doi.org/10.1016/j.orggeochem.2017.12.003>

1213 Sadori, L., 2018. The Lateglacial and Holocene vegetation and climate history of Lago di
1214 Mezzano (central Italy). *Quaternary Science Reviews* 202, 30–44.
1215 <https://doi.org/10.1016/j.quascirev.2018.09.004>

1216 Salonen, J.S., Korpela, M., Williams, J.W., Luoto, M., 2019. Machine-learning based
1217 reconstructions of primary and secondary climate variables from North American and
1218 European fossil pollen data. *Sci Rep* 9, 15805. [https://doi.org/10.1038/s41598-019-](https://doi.org/10.1038/s41598-019-52293-4)
1219 [52293-4](https://doi.org/10.1038/s41598-019-52293-4)

- 1220 Samartin, S., Heiri, O., Joos, F., Renssen, H., Franke, J., Brönnimann, S., Tinner, W., 2017.
 1221 Warm Mediterranean mid-Holocene summers inferred from fossil midge assemblages.
 1222 Nature Geosci 10, 207–212. <https://doi.org/10.1038/ngeo2891>
- 1223 Sanchi, L., Ménot, G., Bard, E., 2014. Insights into continental temperatures in the northwestern
 1224 Black Sea area during the Last Glacial period using branched tetraether lipids.
 1225 Quaternary Science Reviews 84, 98–108.
 1226 <https://doi.org/10.1016/j.quascirev.2013.11.013>
- 1227 Sbaffi, L., Wezel, F.C., Curzi, G., Zoppi, U., 2004. Millennial- to centennial-scale
 1228 palaeoclimatic variations during Termination I and the Holocene in the central
 1229 Mediterranean Sea. Global and Planetary Change 40, 201–217.
 1230 [https://doi.org/10.1016/S0921-8181\(03\)00111-5](https://doi.org/10.1016/S0921-8181(03)00111-5)
- 1231 Sicre, M.-A., Siani, G., Genty, D., Kallel, N., Essallami, L., 2013. Seemingly divergent sea
 1232 surface temperature proxy records in the central Mediterranean during the last deglacial.
 1233 Climate of the Past 9, 1375–1383. <https://doi.org/10.5194/cpd-9-683-2013>
- 1234 Sinninghe Damsté, J.S., Rijpstra, W.I.C., Foesel, B.U., Huber, K.J., Overmann, J., Nakagawa,
 1235 S., Kim, J.J., Dunfield, P.F., Dedysh, S.N., Villanueva, L., 2018. An overview of the
 1236 occurrence of ether- and ester-linked iso-diabolic acid membrane lipids in microbial
 1237 cultures of the Acidobacteria: Implications for brGDGT paleoproxies for temperature
 1238 and pH. Organic Geochemistry 124, 63–76.
 1239 <https://doi.org/10.1016/j.orggeochem.2018.07.006>
- 1240 Smith, V.C., Isaia, R., Pearce, N.J.G., 2011. Tephrostratigraphy and glass compositions of post-
 1241 15 kyr Campi Flegrei eruptions: implications for eruption history and
 1242 chronostratigraphic markers. Quaternary Science Reviews 30, 3638–3660.
 1243 <https://doi.org/10.1016/j.quascirev.2011.07.012>
- 1244 Stockhecke, M., Bechtel, A., Peterse, F., Guillemot, T., Schubert, C.J., 2021. Temperature,
 1245 precipitation, and vegetation changes in the Eastern Mediterranean over the last
 1246 deglaciation and Dansgaard-Oeschger events. Palaeogeography, Palaeoclimatology,
 1247 Palaeoecology 577, 110535. <https://doi.org/10.1016/j.palaeo.2021.110535>
- 1248 Sun, Q., Chu, G., Liu, M., Xie, M., Li, S., Ling, Y., Wang, X., Shi, L., Jia, G., Lü, H., 2011.
 1249 Distributions and temperature dependence of branched glycerol dialkyl glycerol
 1250 tetraethers in recent lacustrine sediments from China and Nepal. J. Geophys. Res. 116,
 1251 G01008. <https://doi.org/10.1029/2010JG001365>
- 1252 Taffetani, F., Catorci, A., Ciaschetti, G., Cutini, M., Di Martino, L., Frattaroli, A.R., Paura, B.,
 1253 Pirone, G., Rismondo, M., Zitti, S., 2012. The Quercus cerris woods of the alliance
 1254 Carpinion orientalis Horvat 1958 in Italy. Plant Biosystems - An International Journal
 1255 Dealing with all Aspects of Plant Biology 146, 918–953.
 1256 <https://doi.org/10.1080/11263504.2012.682613>
- 1257 Tarroso, P., Carrión, J., Dorado-Valiño, M., Queiroz, P., Santos, L., Valdeolillos-Rodríguez,
 1258 A., Célio Alves, P., Brito, J.C., Cheddadi, R., 2016. Spatial climate dynamics in the
 1259 Iberian Peninsula since 15 000 yr BP. Climate of the Past 12, 1137–1149.
 1260 <https://doi.org/10.5194/cp-12-1137-2016>
- 1261 ter Braak, C.J.F., Juggins, S., 1993. Weighted averaging partial least squares regression (WA-
 1262 PLS): an improved method for reconstructing environmental variables from species
 1263 assemblages 18.
- 1264 ter Braak, C.J.F., van Dam, H., 1989. Inferring pH from diatoms: a comparison of old and new
 1265 calibration methods. Hydrobiologia 178, 209–223.
 1266 <https://doi.org/10.1007/BF00006028>
- 1267 Tomlinson, E.L., Arienzo, I., Civetta, L., Wulf, S., Smith, V.C., Hardiman, M., Lane, C.S.,
 1268 Carandente, A., Orsi, G., Rosi, M., Müller, W., Menzies, M.A., 2012. Geochemistry of
 1269 the Phlegraean Fields (Italy) proximal sources for major Mediterranean tephra:

1270 Implications for the dispersal of Plinian and co-ignimbritic components of explosive
1271 eruptions. *Geochimica et Cosmochimica Acta* 93, 102–128.
1272 <https://doi.org/10.1016/j.gca.2012.05.043>

1273 Valente, E., Buscher, J.T., Jourdan, F., Petrosino, P., Reddy, S.M., Tavani, S., Corradetti, A.,
1274 Ascione, A., 2019. Constraining mountain front tectonic activity in extensional setting
1275 from geomorphology and Quaternary stratigraphy: A case study from the Matese ridge,
1276 southern Apennines. *Quaternary Science Reviews* 219, 47–67.
1277 <https://doi.org/10.1016/j.quascirev.2019.07.001>

1278 Van Geel, B., 2002. Non-Pollen Palynomorphs, in: Smol, J.P., Birks, H.J.B., Last, W.M.,
1279 Bradley, R.S., Alverson, K. (Eds.), *Tracking Environmental Change Using Lake*
1280 *Sediments, Developments in Paleoenvironmental Research*. Springer Netherlands,
1281 Dordrecht, pp. 99–119. https://doi.org/10.1007/0-306-47668-1_6

1282 Vescovi, E., Kaltenrieder, P., Tinner, W., 2010. Late-Glacial and Holocene vegetation history
1283 of Pavullo nel Frignano (Northern Apennines, Italy). *Review of Palaeobotany and*
1284 *Palynology* 160, 32–45. <https://doi.org/10.1016/j.revpalbo.2010.01.002>

1285 Walker, M., Lowe, J., Blockley, S.P.E., Bryant, C., Coombes, P., Davies, S., Hardiman, M.,
1286 Turney, C.S.M., Watson, J., 2012. Lateglacial and early Holocene palaeoenvironmental
1287 ‘events’ in Sluggan Bog, Northern Ireland: comparisons with the Greenland NGRIP
1288 GICC05 event stratigraphy. *Quaternary Science Reviews* 36, 124–138.
1289 <https://doi.org/10.1016/j.quascirev.2011.09.008>

1290 Watson, B.I., Williams, J.W., Russell, J.M., Jackson, S.T., Shane, L., Lowell, T.V., 2018.
1291 Temperature variations in the southern Great Lakes during the last deglaciation:
1292 Comparison between pollen and GDGT proxies. *Quaternary Science Reviews* 182, 78–
1293 92. <https://doi.org/10.1016/j.quascirev.2017.12.011>

1294 Weijers, J.W.H., Panoto, E., van Bleijswijk, J., Schouten, S., Rijpstra, W.I.C., Balk, M., Stams,
1295 A.J.M., Damsté, J.S.S., 2009. Constraints on the Biological Source(s) of the Orphan
1296 Branched Tetraether Membrane Lipids. *Geomicrobiology Journal* 26, 402–414.
1297 <https://doi.org/10.1080/01490450902937293>

1298 Weijers, J.W.H., Schouten, S., Spaargaren, O.C., Sinninghe Damsté, J.S., 2006. Occurrence
1299 and distribution of tetraether membrane lipids in soils: Implications for the use of the
1300 TEX86 proxy and the BIT index. *Organic Geochemistry* 37, 1680–1693.
1301 <https://doi.org/10.1016/j.orggeochem.2006.07.018>

1302 Weijers, J.W.H., Schouten, S., van den Donker, J.C., Hopmans, E.C., Sinninghe Damsté, J.S.,
1303 2007. Environmental controls on bacterial tetraether membrane lipid distribution in
1304 soils. *Geochimica et Cosmochimica Acta* 71, 703–713.
1305 <https://doi.org/10.1016/j.gca.2006.10.003>

1306 Wulf, S., Kraml, M., Brauer, A., Keller, J., Negendank, J.F.W., 2004. Tephrochronology of the
1307 100ka lacustrine sediment record of Lago Grande di Monticchio (southern Italy).
1308 *Quaternary International* 122, 7–30. <https://doi.org/10.1016/j.quaint.2004.01.028>

1309 Wulf, S., Kraml, M., Keller, J., 2008. Towards a detailed distal tephrostratigraphy in the Central
1310 Mediterranean: The last 20,000 yrs record of Lago Grande di Monticchio. *Journal of*
1311 *Volcanology and Geothermal Research* 177, 118–132.
1312 <https://doi.org/10.1016/j.jvolgeores.2007.10.009>

1313 Yang, H., Pancost, R.D., Dang, X., Zhou, X., Evershed, R.P., Xiao, G., Tang, C., Gao, L., Guo,
1314 Z., Xie, S., 2014. Correlations between microbial tetraether lipids and environmental
1315 variables in Chinese soils: Optimizing the paleo-reconstructions in semi-arid and arid
1316 regions. *Geochimica et Cosmochimica Acta* 126, 49–69.
1317 <https://doi.org/10.1016/j.gca.2013.10.041>

1318 Zhang, J., Bai, Y., Xu, S., Lei, F., Jia, G., 2013. Alkenone and tetraether lipids reflect different
1319 seasonal seawater temperatures in the coastal northern South China Sea. *Organic*
1320 *Geochemistry* 58, 115–120. <https://doi.org/10.1016/j.orggeochem.2013.02.012>
1321