

# Technical note: A new online tool for $\delta^{18}\text{O}$ -temperature conversions

Daniel E. Gaskell<sup>1</sup>, Pincelli M. Hull<sup>1</sup>

<sup>1</sup>Department of Earth & Planetary Sciences, Yale University, New Haven, CT 06511, USA

*Correspondence to:* Daniel E. Gaskell (daniel.gaskell@yale.edu)

5 **Abstract.** The stable oxygen isotopic composition of marine carbonates ( $\delta^{18}\text{O}_c$ ) is one of the oldest and most widely-used paleothermometers. However, interpretation of these data is complicated by the necessity of knowing the  $\delta^{18}\text{O}$  of the source seawater ( $\delta^{18}\text{O}_w$ ) from which  $\text{CaCO}_3$  is precipitated. The effect of local hydrography (the “salinity effect”) is particularly difficult to correct for and may lead to errors of  $>10^\circ\text{C}$  in sea-surface temperatures if neglected. A variety of methods for calculating  $\delta^{18}\text{O}_w$  have been developed in the literature, but not all are readily accessible to workers. Likewise, temperature  
10 estimates are sensitive to a range of other calibration choices (such as calibration species and the inclusion or exclusion of carbonate ion effects) which can require significant effort to intercompare. We present an online tool for  $\delta^{18}\text{O}$ -temperature conversions which provides convenient access to a wide range of calibrations and methods from the literature. Our tool provides a convenient way for workers to examine the effects of alternate calibration and correction procedures on their  $\delta^{18}\text{O}$ -based temperature estimates.

## 15 1 Motivation

The stable oxygen isotopic composition of carbonates ( $\delta^{18}\text{O}_c$ ) is one of the oldest and most widely-used paleothermometers and undergirds a wide variety of paleoceanographic research (for recent reviews, see Pearson, 2012; Sharp, 2017). Converting  $\delta^{18}\text{O}_c$  to temperature is typically done using an empirical calibration in either a linear form such as

$$T = 16.5 - 4.80(\delta^{18}\text{O}_c - \delta^{18}\text{O}_w - 0.27), \quad (1)$$

20 (Bemis et al., 1998), or in a quadratic form such as

$$T = 16.0 - 5.17(\delta^{18}\text{O}_c - \delta^{18}\text{O}_w - 0.20) + 0.09(\delta^{18}\text{O}_c - \delta^{18}\text{O}_w - 0.20)^2, \quad (2)$$

(McCrea, 1950; as reformulated by Bemis et al., 1998), where  $T$  is temperature (in  $^\circ\text{C}$ ),  $\delta^{18}\text{O}_c$  is the oxygen isotope composition of the carbonate (as ‰ VPDB), and  $\delta^{18}\text{O}_w$  is the oxygen isotope composition of the water in which the carbonate was precipitated (as ‰ VSMOW). Much of the complexity of using  $\delta^{18}\text{O}$  as a paleothermometer arises from the need to know  
25  $\delta^{18}\text{O}_w$ , which may vary both globally as a function of ice volume and locally at the sea surface as a function of regional hydrography (Rohling, 2013). Global variation can be estimated using independent records of sea level, so the global record of deep-water  $\delta^{18}\text{O}$ -based temperatures has been relatively well-established (Zachos et al., 2001; Cramer et al., 2009; Westerhold et al., 2020; Rohling et al., 2021; etc.) However, local variations in surface  $\delta^{18}\text{O}_w$  are more difficult to predict,

rendering sea-surface temperature (SST) estimates from  $\delta^{18}\text{O}$  less reliable than deep-water temperature estimates. To address  
30 this, a variety of methods have been developed in the literature to estimate surface  $\delta^{18}\text{O}_w$ .

Since modern surface  $\delta^{18}\text{O}_w$  broadly covaries with latitude, a common approach has been to apply the modern latitudinal  
variation to a sample's paleolatitude (typically using the relationship fit from Southern Ocean data in Zachos et al., 1994 Eq.  
1; or more recently the updated method of Hollis et al., 2019). However, this approach performs particularly poorly in the  
North Atlantic and other high northern latitudes, where local  $\delta^{18}\text{O}_w$  can deviate significantly from the latitudinal mean (Fig. 1;  
35 Zachos et al., 1994; Gaskell et al., 2022; see also generally Tindall et al., 2010). It also assumes that the latitudinal gradient in  
 $\delta^{18}\text{O}_w$  has not changed through time, which is contradicted by modeling. In warmer climates with an altered hydrological cycle,  
models predict that regional salinity contrasts should change due to alterations in the local ratio of evaporation to precipitation  
(Richter and Xie, 2010; Singh et al., 2016), with an analogous effect on  $\delta^{18}\text{O}_w$  (Zhou et al., 2008; Tindall et al., 2010; Roberts  
et al., 2011; Zhu et al., 2020). In particularly extreme cases such as the Eocene, the theoretical difference between modern  
40 latitude-derived  $\delta^{18}\text{O}_w$  (after Zachos et al., 1994 Eq. 1) and modeled local  $\delta^{18}\text{O}_w$  at 6x preindustrial  $p\text{CO}_2$  (Zhu et al., 2020)  
yields a mean temperature error of 5 °C in the Southern Ocean (60–90 °S) or an astonishing mean temperature error of 41 °C  
above the Arctic Circle (66.5–90 °N; Figure 1).

An alternative approach is to obtain  $\delta^{18}\text{O}_w$  more or less directly from isotope-enabled climate models (Zhou et al., 2008;  
Roberts et al., 2011; Gaskell et al., 2022). Several approaches have been adopted: drawing local  $\delta^{18}\text{O}_w$  directly from model  
45 output (Roberts et al., 2011); using modeled zonal mean  $\delta^{18}\text{O}_w$  for a particular paleolatitude (Zhou et al., 2008); using models  
as input to fit a generalized equation for predicting  $\delta^{18}\text{O}_w$  from latitude and bottom-water temperature (Gaskell et al., 2022 Eq.  
S9); or, recently, a generalized method which uses bottom-water temperature to interpolate local  $\delta^{18}\text{O}_w$  between models run at  
different  $p\text{CO}_2$  (Gaskell et al., 2022). While some authors have avoided these approaches altogether due to the uncertainty of  
modeled  $\delta^{18}\text{O}_w$  (e.g., Hollis et al., 2012) or the possibility of introducing circularity into data-model comparisons (e.g., Hollis  
50 et al., 2019), model-derived  $\delta^{18}\text{O}_w$  clearly captures information lost by simpler approaches and is therefore appropriate for  
some use-cases (Roberts et al., 2011).

Here, a new online tool for  $\delta^{18}\text{O}$  temperature conversion is presented which automates a range of methods for  $\delta^{18}\text{O}_w$   
reconstruction and correction from the literature, improving the accessibility of advanced methods to workers generating  $\delta^{18}\text{O}_e$   
data.

## 55 2 Description

We present a new online tool for performing  $\delta^{18}\text{O}_e$ -temperature conversions which automates a range of methods from the  
literature. A preprint version of this tool is available at <https://www.danielgaskell.com/d18O> [NOTE: This URL will be updated  
to a permanent institutional URL for publication.] The general workflow for using the tool is summarized in Figure 2; details  
on the methodology and reasoning behind each option are given below.

## 60 2.1 $\delta^{18}\text{O}_c$ -temperature calibration

After manually entering or uploading a datasheet of  $\delta^{18}\text{O}_c$  measurements in .csv format, users may select from one of 62 different calibrations from the literature (Bemis et al., 1998; Böhm et al., 2000; Bouvier-Soumagnac and Duplessy, 1985; Duplessy et al., 2002; Epstein et al., 1953; Erez and Luz, 1983; Farmer et al., 2007; Geffen, 2012; Godiksen et al., 2010; Grossman and Ku, 1986; Høie et al., 2004; Juillet-Leclerc and Schmidt, 2001; Kim and O’Neil, 1997; Kim et al., 2007; Lynch-  
65 Stieglitz et al., 1999; Malevich et al., 2019; Marchitto et al., 2014; McCrea, 1950; Mulitza et al., 2004; O’Neil et al., 1969; Patterson et al., 1993; Reynaud-Vaganay et al., 1999; Rosenheim et al., 2009; Shackleton, 1974; Storm-Suke et al., 2007; Thorrold et al., 1997; Tremaine et al., 2011; White et al., 1999; Willmes et al., 2019). All data are expressed with  $\delta^{18}\text{O}_c$  in units of ‰ VPDB and  $\delta^{18}\text{O}_w$  in units of ‰ VSMOW, with any standard interconversions expected by the chosen calibration performed automatically. Standard interconversion is notably inconsistent in the literature, with many paleoceanographic  
70 papers employing the relationship  $\delta^{18}\text{O}_{\text{VPDB}} = \delta^{18}\text{O}_{\text{VSMOW}} - 0.27\text{‰}$  (Hut, 1987) while many geochemical papers employ the incompatible relationship  $\delta^{18}\text{O}_{\text{VPDB}} = 0.97001 \delta^{18}\text{O}_{\text{VSMOW}} - 29.99\text{‰}$  (Brand et al., 2014). The former is actually the isotopic offset between the related VPDB-CO<sub>2</sub> and VSMOW-CO<sub>2</sub> scales, but the difference is unimportant so long as all data are treated in the manner the calibration expects, as our tool ensures. Examples of common calibrations and their included standard conversions are given in Table 1.

75 Where applicable, we use the standardized reformulations of Bemis et al. (1998) and Willmes et al. (2019), or exact algebraic rearrangements of the original equations. For the bayfox core-top calibrations of Malevich et al. (2019), the standard bayfox tool re-fits the calibration coefficients with every run. Since this is computationally expensive, we instead use the linear calibration coefficients fit by runs of bayfoxr 0.0.1 directly in linear functions of the form of Eq. 1 (see Table 1). These yield results equivalent to the full fitting process within numerical error (mean residual =  $\pm 0.02$  °C, identical to the mean scatter  
80 between replicates of the full bayfox fit).

## 2.2 Global $\delta^{18}\text{O}_w$ estimation

Users may specify global  $\delta^{18}\text{O}_w$  manually or choose to draw  $\delta^{18}\text{O}_w$  by sample age from of 12 different timeseries of global  $\delta^{18}\text{O}_w$  from the literature (from Cramer et al., 2011; Henkes et al., 2018; Meckler et al., 2022; Miller et al., 2020; Modestou et al., 2020; Rohling et al., 2021; Veizer and Prokoph, 2015). These records are typically constructed by assuming that the benthic  
85  $\delta^{18}\text{O}$  record reflects a combination of temperature and ice volume and then subtracting out an independent record of temperature (e.g., using Mg/Ca-based bottom-water temperatures; Cramer et al., 2011) or ice volume (e.g., using a multi-proxy sea level reconstruction; Rohling et al., 2021) to determine the residual  $\delta^{18}\text{O}_w$ . Which global  $\delta^{18}\text{O}_w$  record is most realistic remains a contentious topic in the literature, with sea-level and Mg/Ca-based records (e.g., Cramer et al., 2011; Rohling et al., 2021) predicting up to  $\sim 1\text{‰}$  lower  $\delta^{18}\text{O}_w$  for much of the Cenozoic than records based on clumped isotope paleothermometry  
90 (Meckler et al., 2022; see also Agterhuis et al., 2022). We provide both classes of record here for comparison by the user.

Records are mapped to the user data's ages by linear interpolation. The  $\Delta 47$ -based  $\delta^{18}\text{O}_w$  records of Meckler et al. (2022) included in our tool were generated by interpolating the authors' original results to 0.1 Ma resolution using the Monte Carlo LOESS method and parameters described in the original publication (Meckler et al., 2022).

All built-in  $\delta^{18}\text{O}_w$  and temperature records are internally converted to four different timescales, so the user can select the timescale consistent with their data: GTS2004 (Gradstein et al., 2005), GTS2012 (Gradstein et al., 2012), GTS2016 (Ogg et al., 2016), and GTS2020 (Gradstein et al., 2020). These timescale conversions are performed by linear interpolation between magnetochron boundaries; dataset files can be found on the project GitHub.

### 2.3 Local $\delta^{18}\text{O}_w$ estimation

The user may select a method for estimating local  $\delta^{18}\text{O}_w$ . These are as follows: performing no local correction; using modern  $\delta^{18}\text{O}_w$  from each sample's location and a specified depth (after LeGrande and Schmidt, 2006); using reconstructed Late Holocene or Last Glacial Maximum surface  $\delta^{18}\text{O}_w$  from each sample's location (model output from Tierney et al., 2020); using  $\delta^{18}\text{O}_w$  estimated from latitude alone (after Zachos et al., 1994 Eq. 1; or the method of Hollis et al., 2019); using  $\delta^{18}\text{O}_w$  estimated from latitude and bottom-water temperature (after Gaskell et al., 2022 Eq. S9); or using  $\delta^{18}\text{O}_w$  estimated from isotope-enabled climate models (GCMs, after the method of Gaskell et al., 2022, presently provided using the datasets of Miocene and Eocene paleogeography used in that publication).

For methods which draw from an existing dataset of  $\delta^{18}\text{O}_w$ , the user may specify a number of degrees latitude/longitude or great-circle radius to average over in order to capture a regional mean when the exact paleocoordinates or local hydrography may not be known. To help determine site locations at the time of deposition, an option is also provided to automatically perform paleocoordinate rotations using the GPLates Web Service (Müller et al., 2018). Ages passed to GPLates are rounded to the nearest 100 ka to reduce the number of API calls.

Our tool does not currently implement any automated consideration of seasonal variation in local  $\delta^{18}\text{O}_w$ , as this is generally treated as negligible by standard methodologies or implicitly baked into the calibration by calibrating against mean annual temperatures and  $\delta^{18}\text{O}_w$  values (e.g., Malevich et al., 2019).

### 2.4 Carbonate chemistry effects

Because  $\delta^{18}\text{O}_c$  is known to vary with aqueous carbonate chemistry (the "carbonate ion effect"; Spero et al., 1997; Bijma et al., 1999; Ziveri et al., 2012), users may also specify a carbonate ion correction factor. This is performed by adjusting  $\delta^{18}\text{O}_c$  with the linear relationship

$$\delta^{18}\text{O}'_c = \delta^{18}\text{O}_c - (s[\text{CO}_3^{2-}] - 200s), \quad (3)$$

where  $\delta^{18}\text{O}_c$  is the uncorrected oxygen isotope composition of the carbonate,  $\delta^{18}\text{O}'_c$  is the corrected oxygen isotope composition of the carbonate,  $s$  is the selected slope of the effect (in ‰ VPDB per  $\mu\text{mol L}^{-1} \text{CO}_3^{2-}$ ), and  $[\text{CO}_3^{2-}]$  is the concentration of carbonate ion in solution (in  $\mu\text{mol kg}^{-1}$ ). This relationship yields no correction when  $[\text{CO}_3^{2-}] = 200 \mu\text{mol kg}^{-1}$ .

<sup>1</sup>, an approximation of the mean modern surface value (after the long-term record of Zeebe and Tyrrell, 2019). The user may specify [CO<sub>3</sub><sup>2-</sup>] manually or select a published long-term record of [CO<sub>3</sub><sup>2-</sup>] (Tyrrell and Zeebe, 2004; Zeebe and Tyrrell, 2019).

## 2.5 Tool output

- 125 On completion, the tool presents a formatted table of the resulting temperatures, along with any intermediate values (such as estimated  $\delta^{18}\text{O}_w$ ) which were required to generate them. Any rows with potential errors (e.g., paleocoordinates which do not yield a valid  $\delta^{18}\text{O}_w$  estimate or temperatures which exceed the data range of the calibration) are flagged with a warning. For reference, a short summary of methods is also generated, including relevant equations and a complete bibliography of citations in both text and BibTeX formats for the methods employed in each run.
- 130 It should be noted that, while the tool automates the process of applying a given calibration method, the user is still responsible for pre-screening their data for diagenetic alteration or other external biases. For example, use of  $\delta^{18}\text{O}$  data from foraminifera must consider factors such as diagenetic recrystallization, depth habitat, shell size, and the presence of gametogenic calcite (for a review, see Pearson, 2012).

## 3 Concluding remarks

- 135 Our tool provides a convenient way for workers to perform  $\delta^{18}\text{O}$ -temperature conversions and explore the sensitivity of their results to different calibrations, corrections, and  $\delta^{18}\text{O}_w$ -reconstruction methods by successively trying different options in the interface. By allowing data-generators to rapidly generate multiple temperature estimates for their records with different underlying assumptions, our tool allows workers to quickly understand and quantify the effects of different assumptions on the resulting temperature estimates.

140

## Code availability

An online version of the most current release of our tool is maintained at <https://www.danielgaskell.com/d18O> [*preprint URL, subject to change*]. Source code (Javascript and PHP) is available from the project's GitHub repository at <https://github.com/danielgaskell/d18Oconverter>.

## 145 References

Agterhuis, T., Ziegler, M., de Winter, N. J., and Lourens, L. J.: Warm deep-sea temperatures across Eocene Thermal Maximum 2 from clumped isotope thermometry, *Commun Earth Environ*, 3, 1–9, <https://doi.org/10.1038/s43247-022-00350-8>, 2022.

- 150 Bemis, B. E., Spero, H. J., Bijma, J., and Lea, D. W.: Reevaluation of the oxygen isotopic composition of planktonic foraminifera: Experimental results and revised paleotemperature equations, *Paleoceanography*, 13, 150–160, <https://doi.org/10.1029/98PA00070>, 1998.
- Bijma, J., Spero, H. J., and Lea, D. W.: Reassessing Foraminiferal Stable Isotope Geochemistry: Impact of the Oceanic Carbonate System (Experimental Results), in: *Use of Proxies in Paleoclimatology*, edited by: Fischer, D. G. and Wefer, P. D. G., Springer Berlin Heidelberg, 489–512, [https://doi.org/10.1007/978-3-642-58646-0\\_20](https://doi.org/10.1007/978-3-642-58646-0_20), 1999.
- 155 Böhm, F., Joachimski, M. M., Dullo, W.-C., Eisenhauer, A., Lehnert, H., Reitner, J., and Wörheide, G.: Oxygen isotope fractionation in marine aragonite of coralline sponges, *Geochimica et Cosmochimica Acta*, 64, 1695–1703, [https://doi.org/10.1016/S0016-7037\(99\)00408-1](https://doi.org/10.1016/S0016-7037(99)00408-1), 2000.
- Bouvier-Soumagnac, Y. and Duplessy, J.-C.: Carbon and oxygen isotopic composition of planktonic foraminifera from laboratory culture, plankton tows and Recent sediment; implications for the reconstruction of paleoclimatic conditions and of the global carbon cycle, *Journal of Foraminiferal Research*, 15, 302–320, <https://doi.org/10.2113/gsjfr.15.4.302>, 1985.
- 160 Brand, W. A., Coplen, T. B., Vogl, J., Rosner, M., and Prohaska, T.: Assessment of international reference materials for isotope-ratio analysis (IUPAC Technical Report), *Pure and Applied Chemistry*, 86, 425–467, 2014.
- Cramer, B. S., Toggweiler, J. R., Wright, J. D., Katz, M. E., and Miller, K. G.: Ocean overturning since the Late Cretaceous: Inferences from a new benthic foraminiferal isotope compilation, *Paleoceanography*, 24, PA4216, <https://doi.org/10.1029/2008PA001683>, 2009.
- 165 Cramer, B. S., Miller, K. G., Barrett, P. J., and Wright, J. D.: Late Cretaceous–Neogene trends in deep ocean temperature and continental ice volume: Reconciling records of benthic foraminiferal geochemistry ( $\delta^{18}\text{O}$  and Mg/Ca) with sea level history, *Journal of Geophysical Research: Oceans*, 116, <https://doi.org/10.1029/2011JC007255>, 2011.
- Duplessy, J.-C., Labeyrie, L., and Waelbroeck, C.: Constraints on the ocean oxygen isotopic enrichment between the Last Glacial Maximum and the Holocene: Paleoceanographic implications, *Quaternary Science Reviews*, 21, 315–330, [https://doi.org/10.1016/S0277-3791\(01\)00107-X](https://doi.org/10.1016/S0277-3791(01)00107-X), 2002.
- 170 Epstein, S., Buchsbaum, R., Lowenstam, H. A., and Urey, H. C.: Revised carbonate-water isotopic temperature scale, *Geological Society of America Bulletin*, 64, 1315–1326, 1953.
- Erez, J. and Luz, B.: Experimental paleotemperature equation for planktonic foraminifera, *Geochimica et Cosmochimica Acta*, 47, 1025–1031, [https://doi.org/10.1016/0016-7037\(83\)90232-6](https://doi.org/10.1016/0016-7037(83)90232-6), 1983.
- 175 Farmer, E. C., Kaplan, A., Menocal, P. B. de, and Lynch-Stieglitz, J.: Corroborating ecological depth preferences of planktonic foraminifera in the tropical Atlantic with the stable oxygen isotope ratios of core top specimens, *Paleoceanography*, 22, <https://doi.org/10.1029/2006PA001361>, 2007.
- Gaskell, D. E., Huber, M., O’Brien, C. L., Inglis, G. N., Acosta, R. P., Poulsen, C. J., and Hull, P. M.: The latitudinal temperature gradient and its climate dependence as inferred from foraminiferal  $\delta^{18}\text{O}$  over the past 95 million years, *Proceedings of the National Academy of Sciences*, 119, e2111332119, <https://doi.org/doi:10.1073/pnas.2111332119>, 2022.
- 180 Geffen, A. J.: Otolith oxygen and carbon stable isotopes in wild and laboratory-reared plaice (*Pleuronectes platessa*), *Environ Biol Fish*, 95, 419–430, <https://doi.org/10.1007/s10641-012-0033-2>, 2012.


- Godiksen, J. A., Svenning, M.-A., Dempson, J. B., Marttila, M., Storm-Suke, A., and Power, M.: Development of a species-specific fractionation equation for Arctic charr (*Salvelinus alpinus* (L.)): an experimental approach, *Hydrobiologia*, 650, 67–77, <https://doi.org/10.1007/s10750-009-0056-7>, 2010.
- 185 Gradstein, F. M., Ogg, J. G., and Smith, A. G. (Eds.): *A Geologic Time Scale 2004*, Cambridge University Press, Cambridge, <https://doi.org/10.1017/CBO9780511536045>, 2005.
- Gradstein, F. M., Ogg, J. G., Schmitz, M. D., and Ogg, G. M. (Eds.): *The Geologic Time Scale*, Elsevier, <https://doi.org/10.1016/C2011-1-08249-8>, 2012.
- 190 Gradstein, F. M., Ogg, J. G., Schmitz, M. D., and Ogg, G. M.: *Geologic Time Scale 2020*, Elsevier, <https://doi.org/10.1016/C2020-1-02369-3>, 2020.
- Grossman, E. L.: Chapter 10 - Oxygen Isotope Stratigraphy, in: *The Geologic Time Scale*, edited by: Gradstein, F. M., Ogg, J. G., Schmitz, M. D., and Ogg, G. M., Elsevier, Boston, 181–206, <https://doi.org/10.1016/B978-0-444-59425-9.00010-X>, 2012.
- 195 Grossman, E. L. and Ku, T.-L.: Oxygen and carbon isotope fractionation in biogenic aragonite: Temperature effects, *Chemical Geology: Isotope Geoscience section*, 59, 59–74, [https://doi.org/10.1016/0168-9622\(86\)90057-6](https://doi.org/10.1016/0168-9622(86)90057-6), 1986.
- Henkes, G. A., Passey, B. H., Grossman, E. L., Shenton, B. J., Yancey, T. E., and Pérez-Huerta, A.: Temperature evolution and the oxygen isotope composition of Phanerozoic oceans from carbonate clumped isotope thermometry, *Earth and Planetary Science Letters*, 490, 40–50, <https://doi.org/10.1016/j.epsl.2018.02.001>, 2018.
- 200 Høie, H., Otterlei, E., and Folkvord, A.: Temperature-dependent fractionation of stable oxygen isotopes in otoliths of juvenile cod (*Gadus morhua* L.), *ICES Journal of Marine Science*, 61, 243–251, <https://doi.org/10.1016/j.icesjms.2003.11.006>, 2004.
- Hollis, C. J., Taylor, K. W. R., Handley, L., Pancost, R. D., Huber, M., Creech, J. B., Hines, B. R., Crouch, E. M., Morgans, H. E. G., Crampton, J. S., Gibbs, S., Pearson, P. N., and Zachos, J. C.: Early Paleogene temperature history of the Southwest Pacific Ocean: Reconciling proxies and models, *Earth and Planetary Science Letters*, 349–350, 53–66, <https://doi.org/10.1016/j.epsl.2012.06.024>, 2012.
- 205 Hollis, C. J., Dunkley Jones, T., Anagnostou, E., Bijl, P. K., Cramwinckel, M. J., Cui, Y., Dickens, G. R., Edgar, K. M., Eley, Y., Evans, D., Foster, G. L., Frieling, J., Inglis, G. N., Kennedy, E. M., Kozdon, R., Laetano, V., Lear, C. H., Littler, K., Lourens, L., Meckler, A. N., Naafs, B. D. A., Pälike, H., Pancost, R. D., Pearson, P. N., Röhl, U., Royer, D. L., Salzmann, U., Schubert, B. A., Seebeck, H., Sluijs, A., Speijer, R. P., Stassen, P., Tierney, J., Tripathi, A., Wade, B., Westerhold, T., Witkowski, C., Zachos, J. C., Zhang, Y. G., Huber, M., and Lunt, D. J.: The DeepMIP contribution to PMIP4: methodologies for selection, compilation and analysis of latest Paleocene and early Eocene climate proxy data, incorporating version 0.1 of the DeepMIP database, *Geoscientific Model Development*, 12, 3149–3206, <https://doi.org/10.5194/gmd-12-3149-2019>, 2019.
- Hut, G.: Consultants’ group meeting on stable isotope reference samples for geochemical and hydrological investigations, International Atomic Energy Agency, Vienna (Austria), 1987.
- 215 Juillet-Leclerc, A. and Schmidt, G.: A calibration of the oxygen isotope paleothermometer of coral aragonite from Porites, *Geophysical Research Letters*, 28, <https://doi.org/10.1029/2000GL012538>, 2001.
- Kim, S.-T. and O’Neil, J. R.: Equilibrium and nonequilibrium oxygen isotope effects in synthetic carbonates, *Geochimica et Cosmochimica Acta*, 61, 3461–3475, [https://doi.org/10.1016/S0016-7037\(97\)00169-5](https://doi.org/10.1016/S0016-7037(97)00169-5), 1997.

- 220 Kim, S.-T., O'Neil, J. R., Hillaire-Marcel, C., and Mucci, A.: Oxygen isotope fractionation between synthetic aragonite and water: Influence of temperature and  $Mg^{2+}$  concentration, *Geochimica et Cosmochimica Acta*, 71, 4704–4715, <https://doi.org/10.1016/j.gca.2007.04.019>, 2007.
- LeGrande, A. N. and Schmidt, G. A.: Global gridded data set of the oxygen isotopic composition in seawater, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2006GL026011>, 2006.
- 225 Lynch-Stieglitz, J., Curry, W. B., and Slowey, N.: A geostrophic transport estimate for the Florida Current from the oxygen isotope composition of benthic foraminifera, *Paleoceanography*, 14, 360–373, <https://doi.org/10.1029/1999PA900001>, 1999.
- Malevich, S. B., Vetter, L., and Tierney, J. E.: Global Core Top Calibration of  $\delta^{18}O$  in Planktic Foraminifera to Sea Surface Temperature, *Paleoceanography and Paleoclimatology*, 34, 1292–1315, <https://doi.org/10.1029/2019PA003576>, 2019.
- 230 Marchitto, T. M., Curry, W. B., Lynch-Stieglitz, J., Bryan, S. P., Cobb, K. M., and Lund, D. C.: Improved oxygen isotope temperature calibrations for cosmopolitan benthic foraminifera, *Geochimica et Cosmochimica Acta*, 130, 1–11, <https://doi.org/10.1016/j.gca.2013.12.034>, 2014.
- McCrea, J. M.: On the Isotopic Chemistry of Carbonates and a Paleotemperature Scale, *J. Chem. Phys.*, 18, 849–857, <https://doi.org/10.1063/1.1747785>, 1950.
- 235 Meckler, A. N., Sexton, P. F., Piasecki, A. M., Leutert, T. J., Marquardt, J., Ziegler, M., Agterhuis, T., Lourens, L. J., Rae, J. W. B., Barnet, J., Tripathi, A., and Bernasconi, S. M.: Cenozoic evolution of deep ocean temperature from clumped isotope thermometry, *Science*, 377, 86–90, <https://doi.org/10.1126/science.abk0604>, 2022.
- Miller, K. G., Browning, J. V., Schmelz, W. J., Kopp, R. E., Mountain, G. S., and Wright, J. D.: Cenozoic sea-level and cryospheric evolution from deep-sea geochemical and continental margin records, *Science Advances*, 6, eaaz1346, <https://doi.org/10.1126/sciadv.aaz1346>, 2020.
- 240 Modestou, S. E., Leutert, T. J., Fernandez, A., Lear, C. H., and Meckler, A. N.: Warm Middle Miocene Indian Ocean Bottom Water Temperatures: Comparison of Clumped Isotope and Mg/Ca-Based Estimates, *Paleoceanography and Paleoclimatology*, 35, e2020PA003927, <https://doi.org/10.1029/2020PA003927>, 2020.
- 245 Mulitza, S., Donner, B., Fischer, G., Paul, A., Pätzold, J., Rühlemann, C., and Segl, M.: The South Atlantic Oxygen Isotope Record of Planktic Foraminifera, in: *The South Atlantic in the Late Quaternary: Reconstruction of Material Budgets and Current Systems*, edited by: Wefer, G., Mulitza, S., and Ratmeyer, V., Springer, Berlin, Heidelberg, 121–142, [https://doi.org/10.1007/978-3-642-18917-3\\_7](https://doi.org/10.1007/978-3-642-18917-3_7), 2004.
- Müller, R. D., Cannon, J., Qin, X., Watson, R. J., Gurnis, M., Williams, S., Pfaffelmoser, T., Seton, M., Russell, S. H. J., and Zahirovic, S.: GPlates: Building a Virtual Earth Through Deep Time, *Geochemistry, Geophysics, Geosystems*, 19, 2243–2261, <https://doi.org/10.1029/2018GC007584>, 2018.
- 250 Ogg, J. G., Ogg, G. M., and Gradstein, F. M.: *A Concise Geologic Time Scale*, Elsevier, <https://doi.org/10.1016/C2009-0-64442-1>, 2016.
- O'Neil, J. R., Clayton, R. N., and Mayeda, T. K.: Oxygen Isotope Fractionation in Divalent Metal Carbonates, *J. Chem. Phys.*, 51, 5547–5558, <https://doi.org/10.1063/1.1671982>, 1969.
- 255 Patterson, W. P., Smith, G. R., and Lohmann, K. C.: Continental Paleothermometry and Seasonality Using the Isotopic Composition of Aragonitic Otoliths of Freshwater Fishes, in: *Climate Change in Continental Isotopic Records*, American Geophysical Union (AGU), 191–202, <https://doi.org/10.1029/GM078p0191>, 1993.



- Pearson, P. N.: Oxygen Isotopes in Foraminifera: Overview and Historical Review, in: *Reconstructing Earth's Deep-Time Climate—The State of the Art in 2012*, Paleontological Society Short Course, vol. 18, The Paleontological Society, 1–38, 2012.
- 260 Reynaud-Vaganay, S., Gattuso, J.-P., Cuif, J.-P., Jaubert, J., and Juillet-Leclerc, A.: A novel culture technique for scleractinian corals: application to investigate changes in skeletal  $\delta^{18}\text{O}$  as a function of temperature, *Marine Ecology Progress Series*, 180, 121–130, <https://doi.org/10.3354/meps180121>, 1999.
- Richter, I. and Xie, S.-P.: Moisture transport from the Atlantic to the Pacific basin and its response to North Atlantic cooling and global warming, *Clim Dyn*, 35, 551–566, <https://doi.org/10.1007/s00382-009-0708-3>, 2010.
- 265 Roberts, C. D., LeGrande, A. N., and Tripathi, A. K.: Sensitivity of seawater oxygen isotopes to climatic and tectonic boundary conditions in an early Paleogene simulation with GISS ModelE-R, *Paleoceanography*, 26, <https://doi.org/10.1029/2010PA002025>, 2011.
- Rohling, E. J.: Oxygen isotope composition of seawater, in: *The Encyclopedia of Quaternary Science*, Elsevier, Amsterdam, 915–922, 2013.
- 270 Rohling, E. J., Yu, J., Heslop, D., Foster, G. L., Opdyke, B., and Roberts, A. P.: Sea level and deep-sea temperature reconstructions suggest quasi-stable states and critical transitions over the past 40 million years, *Science Advances*, 7, eabf5326, <https://doi.org/10.1126/sciadv.abf5326>, 2021.
- Rosenheim, B. E., Swart, P. K., and Willenz, P.: Calibration of sclerosponge oxygen isotope records to temperature using high-resolution  $\delta^{18}\text{O}$  data, *Geochimica et Cosmochimica Acta*, 73, 5308–5319, <https://doi.org/10.1016/j.gca.2009.05.047>, 2009.
- 275 Shackleton, N. J.: Attainment of isotopic equilibrium between ocean water and the benthonic foraminifera genus *Unigerina*: isotopic changes in the ocean during the last glacial., *Centre Natl. Rech. Sci. Coll. Inter.*, 219, 203–209, 1974.
- Sharp, Z.: *Principles of Stable Isotope Geochemistry*, 2nd ed., University of New Mexico Open Textbooks, 416 pp., 2017.
- Singh, H. K. A., Donohoe, A., Bitz, C. M., Nusbaumer, J., and Noone, D. C.: Greater aerial moisture transport distances with warming amplify interbasin salinity contrasts, *Geophysical Research Letters*, 43, 8677–8684, <https://doi.org/10.1002/2016GL069796>, 2016.
- 280 Spero, H. J., Bijma, J., Lea, D. W., and Bemis, B. E.: Effect of seawater carbonate concentration on foraminiferal carbon and oxygen isotopes, *Nature*, 390, 497–500, <https://doi.org/10.1038/37333>, 1997.
- Storm-Suke, A., Dempson, J. B., Reist, J. D., and Power, M.: A field-derived oxygen isotope fractionation equation for *Salvelinus* species, *Rapid Communications in Mass Spectrometry*, 21, 4109–4116, <https://doi.org/10.1002/rcm.3320>, 2007.
- 285 Thorrold, S. R., Campana, S. E., Jones, C. M., and Swart, P. K.: Factors determining  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  fractionation in aragonitic otoliths of marine fish, *Geochimica et Cosmochimica Acta*, 61, 2909–2919, [https://doi.org/10.1016/S0016-7037\(97\)00141-5](https://doi.org/10.1016/S0016-7037(97)00141-5), 1997.
- Tierney, J. E., Zhu, J., King, J., Malevich, S. B., Hakim, G. J., and Poulsen, C. J.: Glacial cooling and climate sensitivity revisited, *Nature*, 584, 569–573, <https://doi.org/10.1038/s41586-020-2617-x>, 2020.
- 290 Tindall, J., Flecker, R., Valdes, P., Schmidt, D. N., Markwick, P., and Harris, J.: Modelling the oxygen isotope distribution of ancient seawater using a coupled ocean–atmosphere GCM: Implications for reconstructing early Eocene climate, *Earth and Planetary Science Letters*, 292, 265–273, <https://doi.org/10.1016/j.epsl.2009.12.049>, 2010.

- Tremaine, D. M., Froelich, P. N., and Wang, Y.: Speleothem calcite farmed in situ: Modern calibration of  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  paleoclimate proxies in a continuously-monitored natural cave system, *Geochimica et Cosmochimica Acta*, 75, 4929–4950, <https://doi.org/10.1016/j.gca.2011.06.005>, 2011.
- 295 Tyrrell, T. and Zeebe, R. E.: History of carbonate ion concentration over the last 100 million years, *Geochimica et Cosmochimica Acta*, 68, 3521–3530, <https://doi.org/10.1016/j.gca.2004.02.018>, 2004.
- Veizer, J. and Prokoph, A.: Temperatures and oxygen isotopic composition of Phanerozoic oceans, *Earth-Science Reviews*, 146, 92–104, <https://doi.org/10.1016/j.earscirev.2015.03.008>, 2015.
- 300 Westerhold, T., Marwan, N., Drury, A. J., Liebrand, D., Agnini, C., Anagnostou, E., Barnet, J. S. K., Bohaty, S. M., Vleeschouwer, D. D., Florindo, F., Frederichs, T., Hodell, D. A., Holbourn, A. E., Kroon, D., Lauretano, V., Littler, K., Lourens, L. J., Lyle, M., Pälike, H., Röhl, U., Tian, J., Wilkens, R. H., Wilson, P. A., and Zachos, J. C.: An astronomically dated record of Earth’s climate and its predictability over the last 66 million years, *Science*, 369, 1383–1387, <https://doi.org/10.1126/science.aba6853>, 2020.
- 305 White, R. M. P., Dennis, P. F., and Atkinson, T. C.: Experimental calibration and field investigation of the oxygen isotopic fractionation between biogenic aragonite and water, *Rapid Communications in Mass Spectrometry*, 13, 1242–1247, [https://doi.org/10.1002/\(SICI\)1097-0231\(19990715\)13:13<1242::AID-RCM627>3.0.CO;2-F](https://doi.org/10.1002/(SICI)1097-0231(19990715)13:13<1242::AID-RCM627>3.0.CO;2-F), 1999.
- Willmes, M., Lewis, L. S., Davis, B. E., Loisel, L., James, H. F., Denny, C., Baxter, R., Conrad, J. L., Fangué, N. A., Hung, T.-C., Armstrong, R. A., Williams, I. S., Holden, P., and Hobbs, J. A.: Calibrating temperature reconstructions from fish otolith oxygen isotope analysis for California’s critically endangered Delta Smelt, *Rapid Communications in Mass Spectrometry*, 33, 1207–1220, <https://doi.org/10.1002/rcm.8464>, 2019.
- 310 Zachos, J., Pagani, M., Sloan, L., Thomas, E., and Billups, K.: Trends, Rhythms, and Aberrations in Global Climate 65 Ma to Present, *Science*, 292, 686–693, <https://doi.org/10.1126/science.1059412>, 2001.
- Zachos, J. C., Stott, L. D., and Lohmann, K. C.: Evolution of Early Cenozoic marine temperatures, *Paleoceanography*, 9, 353–387, <https://doi.org/10.1029/93PA03266>, 1994.
- 315 Zeebe, R. E. and Tyrrell, T.: History of carbonate ion concentration over the last 100 million years II: Revised calculations and new data, *Geochimica et Cosmochimica Acta*, <https://doi.org/10.1016/j.gca.2019.02.041>, 2019.
- Zhou, J., Poulsen, C. J., Pollard, D., and White, T. S.: Simulation of modern and middle Cretaceous marine  $\delta^{18}\text{O}$  with an ocean-atmosphere general circulation model, *Paleoceanography*, 23, <https://doi.org/10.1029/2008PA001596>, 2008.
- 320 Zhu, J., Poulsen, C. J., Otto-Bliesner, B. L., Liu, Z., Brady, E. C., and Noone, D. C.: Simulation of early Eocene water isotopes using an Earth system model and its implication for past climate reconstruction, *Earth and Planetary Science Letters*, 537, 116164, <https://doi.org/10.1016/j.epsl.2020.116164>, 2020.
- Ziveri, P., Thoms, S., Probert, I., Geisen, M., and Langer, G.: A universal carbonate ion effect on stable oxygen isotope ratios in unicellular planktonic calcifying organisms, *Biogeosciences*, 9, 1025–1032, <https://doi.org/10.5194/bg-9-1025-2012>, 2012.



330 **Figure 1: Effect of estimating SST using measured/modelled local  $\delta^{18}\text{O}_w$  rather than the latitude-based approximation of Zachos et al. (1994) Eq. 1. Modern: comparison with mean annual  $\delta^{18}\text{O}_w$  <50 m depth (after LeGrande and Schmidt, 2006). Last Glacial Maximum (LGM): comparison with inferred annual surface  $\delta^{18}\text{O}_w$  at the LGM (Tierney et al., 2020). Miocene: comparison with CESMv1.2\_CAM5 model run at 400 ppm  $\text{CO}_2$  with Miocene paleogeography (Gaskell et al., 2022). Eocene: comparison with CESM\_1.2\_CAM5 model run at 6x preindustrial  $\text{CO}_2$  with Eocene paleogeography (Zhu et al., 2020). Temperatures are calculated assuming a slope of  $4.80 \text{ }^\circ\text{C } \text{‰}^{-1}$  (Bemis et al., 1998).**




Figure 2: General workflow for using the tool. (Each box may reflect multiple sub-options.)

**Table 1: Commonly applied  $\delta^{18}\text{O}$ :temperature calibrations from the literature. The tool includes a total of 62 calibrations, which are cited above and listed in the tool interface. Full algebraic rearrangements used by the tool are available in the source code on GitHub.  $\Delta^{18}\text{O}_c = \delta^{18}\text{O}_c - \delta^{18}\text{O}_w$ , where the given VSMOW conversion factor is first added to  $\delta^{18}\text{O}_w$  to convert VSMOW into the format expected by the calibration.**

Reference	Material	Method	Equation	VSMOW to VPDB	Bounds (°C)	Typical application
(Kim and O'Neil, 1997)	Inorganic calcite	Laboratory precipitation	$T = 16.1 - 4.64 \Delta^{18}\text{O}_c + 0.09 (\Delta^{18}\text{O}_c)^2$	-0.27‰*	0–40	“Equilibrium calcite” temperatures, e.g., inferring calcification depth from $\delta^{18}\text{O}_c$
(Kim et al., 2007)	Inorganic aragonite	Laboratory precipitation	$T = \frac{17.88 \times 10^3}{1000 \ln \left( \frac{\delta^{18}\text{O}_c + 29.99}{\frac{0.97001}{\delta^{18}\text{O}_w + 1000} + 1000} \right)} - 273.15$	0‰†	0–40	“Equilibrium aragonite” temperatures
(Malevich et al., 2019)	Planktonic foraminifera	Core-top regression	Several Bayesian fits; 50% quantile temperatures for the annual multi-species pooled method follow the relationship: $T = 11.8790 - 4.0562 \Delta^{18}\text{O}_c$	0‰‡	0–29.5	Sea-surface temperatures (SSTs)
(Bemis et al., 1998)	<i>Orbulina universa</i>	Culture experiment	Mean of high-light and low-light calibrations: $T = 15.7 - 4.80 \Delta^{18}\text{O}_c$	-0.27‰§	15–25	Mixed-layer temperatures experienced by photosymbiotic planktonic foraminifera
(Marchitto et al., 2014)	<i>Cibicides</i> and <i>Planulina</i>	Core-top regression	$T = \frac{0.245 - \sqrt{0.045461 + 0.0044 \Delta^{18}\text{O}_c}}{0.0022}$	0‰	-0.6–25.6	Bottom-water temperatures experienced by epifaunal benthic foraminifera
(Marchitto et al., 2014)	<i>Uvigerina peregrina</i>	Core-top regression	Recommended method: subtract 0.47‰ from $\delta^{18}\text{O}_c$ and use <i>Cibicides</i> eq. above	n/a	n/a	Bottom-water temperatures experienced by infaunal benthic foraminifera
<b>Other notable historical calibrations</b>						
(Epstein et al., 1953)	Biogenic carbonates	Modern specimens	$T = 16.5 - 4.30 \Delta^{18}\text{O}_c + 0.14 (\Delta^{18}\text{O}_c)^2$	-0.27‰ <sup>l</sup>	7–30	Temperatures experienced by mollusks and other generic biocalcifiers
(Shackleton, 1974)	<i>Uvigerina</i> spp.	Core-top regression	$T = 16.9 - 4.0 \Delta^{18}\text{O}_c$	-0.20‰*	0.8–7	Bottom-water temperatures experienced by infaunal benthic foraminifera
(Erez and Luz, 1983)	<i>Trilobatus sacculifer</i>	Culture experiment	$T = 17.0 - 4.52 \Delta^{18}\text{O}_c + 0.03 (\Delta^{18}\text{O}_c)^2$	-0.22‰*	14–30	Mixed-layer temperatures experienced by photosymbiotic planktonic foraminifera

340 \*Reformulated by Bemis et al. (1998)

†Rearranged by this work; the relationship  $\delta^{18}\text{O}_{\text{VPDB}} = 0.97001 \delta^{18}\text{O}_{\text{VSMOW}} - 29.99$  (Brand et al., 2014) is included in this equation to convert  $\delta^{18}\text{O}_c$  from VPDB to VSMOW, as required by the calibration, and temperature has been converted from Kelvin to °C

‡Reformulated in this work by extracting the linear coefficients from the Bayesian posterior values

§Rearranged by this work

345 <sup>l</sup>Reformulated by Bemis et al. (1998) (with VSMOW correction after Grossman, 2012)