

# Earliest meteorological readings in San Fernando (Cádiz, Spain; 1799-1813)

Nieves Bravo-Paredes<sup>1</sup>, María Cruz Gallego<sup>1</sup>, Ricardo M. Trigo<sup>2,3</sup> and José Manuel Vaquero<sup>1</sup>

<sup>1</sup>Departamento de Física, Universidad de Extremadura, Badajoz, Spain

5 <sup>2</sup>Instituto Dom Luiz (IDL), Faculdade de Ciências da Universidade de Lisboa, 1749-016 Lisbon, Portugal

<sup>3</sup>Departamento de Meteorologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, 21941-919, Brazil

*Correspondence to:* José Manuel Vaquero (jvaquero@unex.com)

**Abstract.** Cádiz and San Fernando are two nearby towns with a wealth of meteorological records due to their connection with the Spanish Royal Navy officers and enlightened merchants. Several previous works have already recovered a significant amount of meteorological records of interest in these localities. However, unnoticed previously are more than 10 40,000 daily meteorological observations recorded at the Royal Observatory of the Spanish Navy (located in San Fernando) during the period 1799-1813 remained neither digitized nor studied. Here, we have carried out this important task describing the different steps undertaken to achieve it as well as the results obtained. The dataset is composed of different meteorological variables such as atmospheric pressure, air temperature, precipitation, or state of the sky. As a first step a 15 quality control was carried out to find possible errors in the original data or in the digitization process. Moreover, the antique units were converted to modern units. Also, the metadata and an analysis of the data are described. The dataset is freely available to the scientific community and can be downloaded at <https://doi.org/10.5281/zenodo.7104289>.

## 1 Introduction

Long-term series are very important to better understand the variability of the climate as well as to provide a more robust 20 framework to validate long-term trends and extremes. In this context, early meteorological observations are critical to extend instrumental series, particularly during the 18th and 19th centuries. Therefore, climate data rescue should be a continuous effort as part of a long term activity (Brönnimann et al., 2018). Recovering early meteorological observations represents a huge effort played by a large number of scientists all over the world. Some important examples of structured international efforts in this line of activity are the Atmospheric Circulation Reconstructions over the Earth (ACRE) ([http://www.met-](http://www.met-25 acre.org/)


acre.org/) and the International Data Rescue (I-DARE) Portal (<https://www.idare-portal.org/>). Early meteorological data have been also recovered in Latin-America and the Caribbean (Domínguez-Castro et al., 2017), in Canada (Slonosky, 2003), in South Africa (Picas et al., 2019), in Brazil (Farrona et al., 2012), and in Japan (Zaiki et al., 2006).

In Europe, different initiatives have been carried out to retrieve early meteorological data. For example, the ADVICE project reconstructed and analyzed MSLP data from the 16th century onwards (Jones et al., 1999; Luterbacher et al., 2000). There 30 have been more initiatives such as IMPROVE (Camuffo and Jones, 2002), HISTALP (Auer et al., 2007) or ERACLIM

(<http://www.era-clim.eu/>) projects. More recently initiatives include, for example, the CHIMES project that compiles pre-national weather service observations in Switzerland and makes them available in digital format (Pfister et al., 2019; Brugnara et al., 2020).

In the Iberian Peninsula (hereafter IP), different initiatives in the last two decades can be found. For example, Alcoforado et al. (2012) retrieved early Portuguese meteorological measurements from the 18th century, while project SIGN focused on 19th and early 20th century data in Portugal (Morozova and Valente, 2012; Bližňák et al., 2015). Domínguez-Castro et al. (2014) recovered more than 100000 early meteorological observations prior to 1850 in Spain, and Vaquero et al. (2021) recovered more than 750000 instrumental data from 1826 to mid-20th century in the Extremadura region (interior SW Iberia) presented in the CliPastExtrem database. In the Bay of Cádiz, there are several initiatives to rescue meteorological data that led to a wide number of publications (Barriendos et al., 2002; Gallego et al., 2007; Rodrigo, 2012; Domínguez-Castro et al., 2014; Wheeler, 1995; Rodrigo, 2019).

Figure 1 summarizes the data retrieved for the cities of Cádiz (references in black) and San Fernando (references in red). The first instrumental data were recovered by Wheeler (1995) using temperature measurements made during astronomical observations (passage of the Sun through the Cadiz meridian) in 1776. Barriendos et al. (2002) studied the long records of the Urrutia brothers in Cádiz. Gallego et al. (2007) recovered the meteorological records of the “Torre de Vigía” at Cádiz. The Royal Observatory of the Spanish Navy (ROA, from the Spanish “*Real Instituto y Observatorio de la Armada*”) (located in San Fernando) has published its meteorological records in annual volumes since 1870. In addition, other short series have been recovered (Rodrigo, 2012, 2019; Domínguez-Castro et al., 2014).




50 **Figure 1: Data retrieved for the cities of Cádiz (in black) and San Fernando (in red). The meteorological variables of each dataset are on the right: T – temperature, P – pressure, R – rain, WD – wind direction, H – humidity, RN – rainy days, W – different wind variables, ST – state of the sky.**

The main aim of this work is to retrieve the early meteorological observations measured in San Fernando (Cádiz) for the period 1799-1813. Moreover, a quality control and unit changes are carried out to make the data usable. Finally, an analysis of the data is performed. The structure of the paper is as follows: Section 2 describes the data and the instruments used; the method used for the quality control and the unit change of the data is presented in Section 3; the corresponding results are

presented in Section 4; finally, conclusions of this work are presented in Section 5. Note that most of the early instrumental records are relatively short series, often even shorter than the one presented here (Brönnimann et al., 2019). Thus, a collection of this kind of series relating to the same place or different places with similar climatic characteristics can be extremely useful for knowing the climate of the past.

## **2 Data and instruments**

The dataset of early meteorological observations here recovered were recorded in the ROA during the period 1799-1813. The ROA is located in the city of San Fernando (Cádiz) in the southwest of the IP ( $36^{\circ} 27' 42''$  N,  $6^{\circ} 12' 20''$  W). Figure 2(a) shows the location of the city of San Fernando in the IP. Figure 2(b) shows the location of San Fernando on the Bay of Cádiz. Figure 2(c) shows the location of the ROA in the city of San Fernando.



**Figure 2:** (a) The red point shows the location of the city of San Fernando in the IP; (b) The location of San Fernando in the Bay of Cádiz. The box shows the region enlarged in (c); (c) The red point shows the location of the Royal Observatory of the Spanish Navy (ROA) in the city of San Fernando. The maps were created from Instituto Geográfico Nacional ([www.ign.es](http://www.ign.es)) and the *ggspatial* package in R (Dunnington, 2020).


70

The ROA dataset used here (hereafter SF1799-1813) was obtained under the scope of a larger plan to increase astronomical observations at the Observatory starting in 1798. The plan was divided in four classes: *tiempo* (time), *planetas* (planets), *longitudes terrestres* (earth longitudes) and *física celeste* (celestial physics). Specifically, the observations of SF1799-1813 corresponded to the fourth class (*Física Celeste*) (Lafuente and Sellés, 1988). Original data were preserved at the *Real Instituto y Observatorio de la Armada* in San Fernando in a log-book (catalog number: AH1080).

75

The ROA observations were made using different instruments and different units. A total of 45862 discrete observations were made from 1799 to 1813. Figure 3 shows the observation period covered for each meteorological variable of the SF1799-1813 dataset. Pink and green bars correspond to sub-daily observations: 6 am, 12 am, 6 pm and 12 pm hours in pink;

8 am and 2 pm hours in green. Blue bars correspond to daily observations measured at 2 pm. Gray bars correspond to monthly observations. Unfortunately, there are two large gaps in the periods 1802-1804 and 1810-1812. The cause of the first gap could be that the work of the officers carried out on the sea was more valued than the scientific work in the ROA. Because of this, there were few officers working at ROA. Unfortunately, the absence of the officers often means that meteorological observations were not made (Lafuente and Sellés, 1988). The second gap is related to the movement of the observer to the Torre de Zimbrellos. The observer was assigned to watch over the French due to Napoleon's occupation during the Spanish War of Independence or Peninsular War. Since there are differences in the observation hours from one period to another, it is possible that there are biases in the daily temperature caused by this fact (Zaiki et al., 2002).



**Figure 3: Observational period covered by the new dataset for each meteorological variable. Pink, green and blue bars correspond to sub-daily observations: 6am, 12am, 6pm and 12pm in pink; 8am and 2pm in green; 2pm in blue. Gray bars correspond to monthly observations.**

Additionally, another dataset of the ROA is used for comparison purposes with the SF1799-1813 dataset. This dataset covers the recent period 1997-2021 and can be found at <https://datosclima.es/> (hereafter SF1997-2021). Finally, a monthly global paleo-reanalysis (EKF400v2) generated to cover the 1600 and 2005 time period by Valler et al. (2021) was used to make comparisons with the SF1799-1813 dataset.

The station elevation appears in the metadata but is different for each period. There is no station elevation data in the metadata for the period 1799-1801. For the period 1805-1809, the station elevation was recorded as 142.5 Burgos foot (1 Burgos foot = 0.2786 m). For the period 1812-1813, the elevation was recorded as 100 feet above sea level. This last value has been chosen for the period 1799-1801 due to the average of the pressure values for the period 1799-1801 is quite similar to the average of the pressure values for the period 1812-1813. In addition, the observations were registered with the same instruments and could be the same location. A brief description of each variable is given below:

- **Pressure:**

Pressure observations were recorded with two different mercury barometers and in different units:

- 1799-1801 and 1812-1813: pressure observations were recorded subdaily in English inches with a George Adams barometer.
- 1805-1809: pressure observations were recorded subdaily in French inches with a Megnié barometer.

- **Temperature (thermometer attached to the barometer):**

Mercury barometers typically had a thermometer attached to do the temperature correction for pressure observations. Temperature observations of the thermometers attached to the barometers were recorded subdaily with different thermometers in different units:

110 - 1799-1801 and 1812-1813: the thermometer attached to the George Adams barometer registered subdaily observations in Fahrenheit degrees.

- 1805-1809: the thermometer attached to the Megnié barometer recorded subdaily observations in Reaumur degrees.

- **Air temperature:**

Air temperature was recorded with a Dollond thermometer in Fahrenheit degrees for the period 1805-1809.

115 • **Wind direction:**

Wind direction observations were recorded subdaily by following the wind rose directions. The wind rose was divided into 32 directions: N, NxNE, NNE, NExN, NE, NExE, ENE, ExNE, E, ExSE, ESE, SExE, SE, SExS, SSE, SxSE, S, SxSW, SSW, SWxS, SW, SWxW, WSW, WxSW, W, WxNW, WNW, NWxW, NW, NWxN, NNW, NxNW. The term “x” between directions means “by” in English. In the original data, these wind directions were recorded as, for example, “NE1/4E”. In the digitization process, the term “NE1/4E” was replaced by “NExE”.

120 The prevailing wind direction in the 24h was recorded for the period 1812-1813, although there is no further indication in the original manuscript.

- **Wind force:**

125 Wind force observations were recorded subdaily in textual form using descriptors. García-Herrera et al. (2003) wrote a dictionary to clarify the archaic wind force terms in a comprehensible format. This dictionary was used to check the wind force terms from the original document. Most of the terms for the whole period appear in the dictionary.

The prevailing wind force in the 24h were recorded for the period 1812-1813.

- **State of the sky:**

130 The observers gave a detailed description of the state of the sky subdaily during the period 1799-1801. Clear or covered sky, amount of clouds, rain or other type of precipitation are examples of what usually appears in the description.

The state of sky was recorded differently since 1805. The description was summarized in just one word, for example, “clear” to indicate that the sky was cloudless.

The prevailing state of the sky in the 24h were recorded for the period 1812-1813.

135 Moreover, the number of clear, covered, rainy and stormy days were also registered monthly in different periods. Clear, covered and rainy days were recorded for the period 1799-1800. Rainy and stormy days were recorded for the period 1812-1813.

- **Humidity:**

There is not information about the humidity in the metadata. The humidity was registered only for the period 06/01/1801-11/12/1801.

140 • **Precipitation:**

The precipitation was recorded weekly in Burgos inches (1 inch = 23.22 mm) for the period 1805-1808 and monthly for the year 1809 and the period 1812-1813.

### 3 Method

145 Data were digitized by key entry from photographs taken from the manuscript. The dataset was analyzed through a basic quality control applied to detect possible errors in the digitization process or suspicious values (Vaquero et al., 2022). Moreover, the units were changed to modern SI units. The steps will be described in the following sections.

#### 3.1 Quality control

150 Digitized data often contains errors as a result of typing an incorrect number in the digital document instead of the actual number in the original document. These errors should be detected and corrected whenever possible. Moreover, quality control could detect suspicious values. These values were digitized correctly. Therefore, it is possible that the observers wrote an incorrect number instead of the correct number (e.g.: a maximum daily temperature with lower values than the corresponding minimum daily temperature). Suspicious values detected by quality control are flagged with an asterisk and are not corrected.

155 A basic quality control was carried out in order to detect possible typos, errors or suspicious values in the digitization process. The quality control consists of the following:

1. **Visual check:** some variables are bounded by upper and/or lower limit values. For example, pressure lines cannot exceed the value of 12. If there are values that exceed this threshold, the values are checked against the original data and corrected.
- 160 2. **Compare means:** the daily and monthly means calculated by the observers were digitized and compared with the means computed from the digitized data. If the means are not equal, the values corresponding to that mean are checked against the original data and corrected whenever a mistake is detected.
3. **Extreme values:** in order to detect outlier values, the mean plus/minus three standard deviations are calculated by the authors. Values that exceed this threshold are checked against the original data and corrected if there is an obvious error.
- 165 4. **Consecutive values:** the difference of the values between consecutive days is also calculated. This difference should not exceed a predefined threshold. The threshold is estimated by the mean plus/minus three standard deviations of the differences. Values above this threshold could be due to an error, a suspicious value or an outlier. Values that exceed this threshold are checked against the original data and corrected if it is an error.

The four quality control steps mention above were applied to pressure and temperature variables (temperature related to the thermometer attached to the barometer). For the air temperature variable and for the humidity only steps 1, 3 and 4 were applied.

### 3.2 Unit change

#### 3.2.1 Temperature

Temperature was measured using the Reaumur and Fahrenheit scales in different periods. Temperatures were converted to Celsius using the two following equations where Eq. (1) converts Reaumur to Celsius, while Eq. (2) converts Fahrenheit to Celsius.

$$T = \frac{5}{4}T_R \quad (1)$$

$$T = \frac{5}{9}(T_F - 32) \quad (2)$$

#### 3.2.2 Pressure

Pressure was measured in two different length units, English and French inches, in two different periods. Pressure measurements originally registered in inches were converted to meters. Equation (3) is used to convert the English inch to meters. Equation (4) is used to convert the French inch to meters.

$$h = 0.0254h_{mercury} \quad (3)$$

$$h = 0.02707h_{mercury} \quad (4)$$

#### 185 • Reduction to 0 °C:

Mercury expands and shrinks depending on the temperature. Therefore, meteorological observations recorded with a mercury barometer should be reduced to 0 °C. The following equation gives the density of mercury as a function of temperature:

$$\rho_{Hg} = \frac{\rho_0}{1+\alpha T}$$

190 where  $\alpha=0.0001818 \text{ K}^{-1}$  is the volumetric thermal coefficient for mercury at 0 °C,  $\rho_0=13595.1 \text{ Kg m}^{-3}$  is the density of mercury at 0 °C and  $T$  is the temperature of the thermometer attached to the barometer.

The barometer scale could expand due to temperature. Therefore, the scale must be corrected. The linear thermal expansion equation can be written as:

$$\beta = \frac{1}{L} \left( \frac{\Delta L}{\Delta T} \right)$$



195 Thus, expanding on this equation, the height of the scale corrected to 0 °C ( $h_0$ ) can be calculated with Eq. (5).

$$h_0 = [1 + \beta T]h \quad (5)$$

Finally, the pressure in meters of mercury is converted to hectopascal through Eq. (6):

$$p_0 = g_n \frac{\rho_0}{1 + \alpha T} [1 + \beta T]h \quad (6)$$

where  $h$  is the height of the mercury,  $T$  is the temperature of the thermometer attached to the barometer, and  $g_n=9.80665 \text{ m s}^{-2}$  is the normalized gravity.

- **Correction for local gravity:**

Equation (7) gives the local gravity as a function of the latitude ( $\varphi$ ) and the station elevation ( $H$ ).

$$g_{\varphi,H} = 9.8062(1 - 0.0026442 \cos 2\varphi - 0.0000058 \cos^2 2\varphi) - 0.000003086H \quad (7)$$

Finally, Eq. (8) gives the pressure corrected to 0 °C and local gravity.

205 
$$p_n = p_0 \frac{g_{\varphi,H}}{g_n} \quad (8)$$

- **Reduction to mean sea-level pressure:**

Pressure values at different altitudes should be reduced to mean sea-level pressure when comparisons between them are carried out (WMO, 2018). The equation to obtain pressure values reduced to MSL is:

$$p_{MSL} = p_n \cdot \exp\left(\frac{\frac{g_n H}{R}}{T + \frac{\alpha H}{2}}\right) \quad (9)$$

210 where  $H$  is the station elevation,  $g_n$  is the normalized gravity,  $R=287.05 \text{ J Kg}^{-1} \text{ K}^{-1}$  is the gas constant of dry air,  $T$  is the air temperature, and  $\alpha=0.0065 \text{ K m}^{-1}$  is the assumed lapse-rate in the fictitious air column extending from sea level to the station elevation level.

Air temperature observations are only available for the period 1805-1809. The coefficient of determination between air temperature observations and temperatures of thermometer attached to the barometers observations for the same period is 215  $R^2=0.8638$  (3174 degrees of freedom). Thus,  $T$  in Eq. (9) is considered as the temperature of the thermometer attached to the barometer to cover the entire period 1799-1813.

## 4 Results and discussion

In total, 19618 values were analyzed by the quality control procedures and circa 300 digitized values were detected as errors in the digitization process and were corrected. Three values were detected as suspicious and were flagged with an asterisk.

220 This represents 1.54 % of the total data analyzed (0.66 % of the total data). The total errors in each variable vary substantially: 82 errors were detected in pressure observations, 183 in temperature observations (related to the thermometer attached to the barometer), 18 in air temperature and 17 in humidity observations. The number of total readings analyzed for pressure were 7652, for temperature (thermometer attached to the barometer) 7774, for air temperature 3176 and for humidity 1016. Therefore, the percentage of errors detected in each variable is, respectively: 1.07 % for pressure, 2.35 % for  
225 temperature (related to the barometer), 0.57 % for air temperature and 1.67 % for humidity.


Both pressure series (SF1799-1813 and SF1997-2021) have been compared. For pressure data, the available period of the SF1997-2021 dataset is 2006-2021. The average of the pressure values of SF1799-1813 are lower than the average of the more recent pressure values of 2006-2021. We have made an adjustment of the SF1799-1813 values to be able to compare them with modern data. The adjustment consists of making the difference between the average of the pressure values of the  
230 two series and adding it to the pressure values of the SF1799-1813 series. In fact, two adjustments are required in different periods due to the two different barometers used (George Adams and Megnié barometers). For the period 1799-1801 and 1808-1813, a value of 10.3 hPa was added to the original observations and for the period 1805-1809 a value of 4.9 hPa was added. Other studies found this same problem and have applied similar correction adjustments. For example, the pressure series for Leiden required an adjustment of 16.7 hPa (Können and Brandsma, 2005). An adjustment between 9.0 hPa and  
235 10.5 hPa was equally needed for the pressure series of London (Slonosky et al., 2001). Rodrigo (2019) found a difference between the monthly data of Cádiz for the period 1820-1822 and the mean values for the period 1901-1930 of roughly 3 hPa. According to these authors possible explanations for this problem could be related to the calibration of the instrument, imperfections in the barometers used, the existence of trapped air, or the unknown correct altitude of the barometer.

Time series plots of the monthly pressure means computed from the daily SF1799-1813 and SF1997-2021 series are shown  
240 in Fig. 4. The solid gray line represents the pressure means of SF1799-1813 series for the period 1799-1813. The solid black line shows the adjusted pressure observations of SF1799-1813, at daily (upper panel, Fig. 4) and monthly (lower panel, Fig. 4) scales. The dashed red line represents the monthly average pressure of the SF1997-2021 series for the period 2006-2021. The dashed black line shows the monthly pressure of EKF400v2 dataset for the period 1799-1813. This line corresponds to the nearest grid point of the regular grid of the EKF400v2 dataset.

245 Note that there is a monthly decrease in pressure values in the first months of 1808 (Fig. 4). Cornes et al. (2012) recovered pressure values measured with a Megnié barometer and found that the vacuum of Megnié barometer suffered a drift of a rate of 0.02 hPa per month in the Paris Observatory. It is possible that the Megnié barometer of the ROA (SF1799-1813) suffered a similar drift as the Paris Observatory. In the metadata, the observer wrote in the manuscript that the readings of the Megnié barometer in 1808 were 0.22 inches of pressure lower than the George Adams barometer. Moreover, the readings were lower  
250 than when the barometer was installed. In 1809, the observer wrote in the manuscript that the barometer was repaired in August. The difference of 0.22 inches of pressure between the Megnié and the George Adams barometer mentioned by the observer is checked here. For the period 1808-1809, the pressure observations are 0.22 inches of pressure lower than the

pressure observations for the period 1799-1801. Due to this under-reading, 0.22 inches are added to the original pressure observations (already corrected in Fig. 4) for the period 1808-1809.

255




**Figure 4: The solid gray line represents the pressure means computed from the daily SF1799-1813 series for the period 1799-1813. The solid black line shows the adjusted pressure observations of SF1799-1813, at daily (upper panel) and monthly (lower panel) scales. The dashed red line represents the monthly pressure average cycle of SF1997-2021 series for the period 2006-2021. The dashed black line shows the monthly pressure of EKF400v2 dataset for the period 1799-1813.**

260

In addition, the pressure values of the EKF400v2 dataset are higher than the pressure values of the SF1799-1813 and SF1997-2021 datasets (see Fig. 4, lower panel). The coordinates of the EKF400v2 dataset are close to those of the city of San Fernando but they are not the same. A possible explanation of the difference in pressure values between the different


series can be found in this fact. However, the average monthly range of the EKF400v2 dataset is similar to the average  
265 monthly range of the SF1799-1813 and SF1997-2021 datasets.

Figure 5 shows the daily air temperature of SF1799-1813 for the period 1805-1809 in gray. The daily average of the air  
temperature of SF1997-2021 for the period 1997-2021 is showed in dashed red line. The dashed black line represents the  
monthly air temperature of EKF400v2 dataset for the period 1805-1809. Just like the pressure data, this curve corresponds to  
the nearest grid point of the regular grid tan of the EKF400v2 dataset. As can be seen, the three curves (SF1799-1813,  
270 SF1997-



**Figure 5: The gray line represents the daily air temperature of SF1799-1813 for the period 1805-1809. The dashed red line  
represents the mean monthly air temperature of SF1997-2021 for the period 1997-2021. The dashed black line shows the monthly  
air temperature of EKF400v2 dataset for the period 1805-1809. The two vertical lines indicate the beginning and the end of the  
275 year 1809.**

2021 and EKF400v2) show a similar behavior and there is an excellent compatibility between the air temperature for the SF1799-1813 and SF1997-2021 datasets. The coefficient of determination between these both curves is  $R^2=0.9104$ . For the case of SF1799-1813 and EKF400v2 datasets, the coefficient of determination is  $R^2=0.9384$ . For the case of the deseasonalized values of SF1799-1813 and EKF400v2 datasets, the coefficient of determination is  $R^2=0.4119$ . These last three analyses were performed with monthly values. The air temperature of SF1799-1813 dataset for the period 1805-1809 shows a seasonal cycle similar to the monthly average of the SF1997-2021 dataset and to the monthly temperature of EKF400v2 dataset. The temperature of SF1799-1813 in summer months for the period 1805-1809 is slightly higher (quite similar) than the monthly average of the SF1997-2021 (EKF400v2) dataset. As can be seen in Fig. 5, the temperature of the three series in the first months of each year of the period represented is quite similar except at the beginning of 1809. The temperature values for the first months of 1809 of SF1799-



**Figure 6:** Frequency of the wind direction represented in wind compasses for the 1799-1813 period for each month and in total (upper).


1813 dataset are considerably higher than the monthly temperature of the SF1997-2021 and EKF400v2 datasets for these  
290 months. Interestingly, similarly unusual high values appear in the temperature series of the thermometer attached to the  
barometer (not shown). In order to understand the robustness of this signal we searched for all available instrumental  
temperature series that include the year 1809. We checked the available instrumental data series (not shown) to evaluate if  
this behavior is unique to San Fernando (and therefore a potential error) or if it occurs in other regions. Six different  
temperature series from Switzerland (Brugnara et al., 2020) present the same behavior for the beginning of 1809. Also, two  
295 temperature series from Madrid and Mallorca (Spain) (Domínguez-Castro et al., 2014) show a similar behavior. Therefore, it  
seems that this higher-than usual winter temperatures did occurred throughout Southwest Europe.

Individual subplots represented in Fig. 6 show the frequency of the wind direction represented in wind compasses for the  
1799-1813 period for each month and in total (upper subplot). In general, the compasses show five prevalent wind directions  
(NW, W, SW, SE, and ESE) varying with the seasonal cycle. In winter months (December, January and February), the  
300 predominant wind directions are N, NW and SW. The SE and ESE wind directions are also relevant but to a lesser extent. In  
spring months (March, April and May), the main wind directions are NW, SW and SE, while in summer months (June, July  
and August), the prevalent wind directions are W, SW and ESE. Finally, in autumn months (September, October and  
November), the predominant wind direction is SW, but to a lesser extent other relevant wind directions are NW, SE and  
ESE. The predominant wind directions are in accordance to the location of San Fernando, which is closed to the Strait of  
305 Gibraltar. In general, the frequency of westerlies is higher than the easterlies, as it is expected (Sousa, 1987).

Figure 7 shows the number of observations and the corresponding percentage of the wind force per class (at the top) and the  
state of the sky (at the bottom) for the period 1799-1813 and 1805-1813, respectively. The terms used to categorize the wind  
forces are presented with the original Spanish name. These terms could be converted to the Beaufort scale using CLIWOC  
Dictionary (García-Herrera et al., 2004). The predominant wind forces are three: *flojo* (Beaufort scale = 3), *fresco* (Beaufort  
310 scale = 6) and *fresquito* (Beaufort scale = 5). There are also two additional terms with a high number of observations:  
*bonancible* (Beaufort scale = 4) and *recio* (Beaufort scale = 7). Calvo et al. (2008) obtained similar results for Cádiz (about  
10 km northwest of San Fernando) for the longer period of 1806-1852. The predominant state of the sky is *claro* (clear sky).  
The states of sky *nubes* (cloudy) and *lluvia* (rainy) are also relevant. The term *otros* in the two figures of Fig. 6 represents  
those observations with more than one wind force or state of sky. Nevertheless, this term represents a low number of  
315 observations and is not significant.

The accumulated precipitation per month for the period 1805-1813 can be observed in Fig. 8. In the summer months there is  
no precipitation data. In the metadata there is not information about the absence of observations or the precipitation. For this  
reason, daily pressure data and periods without rainfall have been plotted together (not shown) to check that there are not  
drops in pressure values when there is not precipitation data. According to the represented, it seems that the precipitation  
320 data are consistent. Monthly values above 100 mm occur almost every autumn-winter and the highest precipitation values  
were registered in the month of December in 1812 and October in 1813. Calvo et al. (2008) showed the absolute frequency

of precipitation (days per year) of Cadiz for the same period and obtained similar results. The maximum absolute frequency was recorded in 1813, with a frequency of 62 days per year. There is no data for the year 1812 (Calvo et al., 2008).



325 **Figure 7: Top) Number of observations and the corresponding percentage of the wind force per class for the period 1799-1813 at the top. Bottom) number of observations and the percentage of the state of the sky per class for the period 1805-1813.**


The values of sea level pressure (SLP) and the total precipitation (rain and snow) anomalies of the EKF400v2 dataset are represented below for the months of November and December 1812 and the months of October to December 1813 (Fig. 9). The highest precipitation values were recorded in the months of December 1812 and October and December 1813 for the period 1805-1813 (see Fig. 8). In the months of February and March 1813 no precipitation was recorded, but the metadata shows several days of rain for February and one for March. Therefore, there could be some error in these months, either in the identification of the rainy days or in the amount of accumulated precipitation. The precipitation anomalies estimated for the month of December 1812 for southern Spain is around 200 kg m<sup>-2</sup> (Fig. 9). For the year 1813, high precipitation values were also estimated for the months of October and December and virtually zero anomaly for the month of November for southern Spain. Pressure values of around 1020 hPa are estimated for the months of November and December 1813 and

330

335




around 1018 hPa for October 1813. Overall, the large-scale atmospheric dynamics at the monthly scale is in good agreement with the precipitation values represented in Fig. 8 for these months.



**Figure 8: Accumulated precipitation per month for the period 1805-1813.**

340



**Figure 9: Monthly SLP and monthly precipitation anomalies for the months that registered the maximum values of precipitation (see Fig. 7).**

## 5 Conclusions

345 The Cádiz-San Fernando region is one of the few places on our planet with a variety of overlapping series of early meteorological data. However, to the best of our knowledge, one of the longest and most interesting series (carried out by the

staff of the Spanish Navy observatory) had not been recovered or studied until this work. In total, 45862 daily meteorological observations of the ROA dataset have been recovered for the period 1799-1813. The dataset is composed by different meteorological variables such as atmospheric pressure, air temperature, precipitation or state of the sky. After  
350 applying a number of quality control steps a small amount of values (0.66 % of the data) were detected as suspicious in the digitization process and were corrected. Also, the units have been converted to modern units and the metadata has been described.

In the comparison between SF1799-1813 and SF1997-2021 datasets, a significant difference between the pressure values of both series can be observed. Therefore, two adjustments are applied in different periods: for the period 1799-1801 and 1808-  
355 1813, a value of 10.3 hPa was added to the pressure observations of SF1799-1813 while for the period 1805-1809 a value of 4.9 hPa was added. Moreover, it has been added 0.22 inches of pressure to the original pressure observations of SF1799-1813 dataset for the period 1808-1809. The three datasets compared (SF1799-1813, SF1997-2021 and EKF400v2) show a similar behavior for the air temperature data and there is an excellent agreement between air temperature for the SF1799-1813 and SF1997-2021 datasets.

360 The wind direction frequency analysis shows, when all months of the year are considered, five prevailing wind directions (NW, W, SW, SE, and ESE) for the period 1799-1813. This is in accordance with the location of San Fernando, which is closed to the Strait of Gibraltar. In general, the frequency of westerlies is higher than the easterlies.

The wind force analysis shows three predominant wind forces (*flojo* (Beaufort scale = 3), *fresco* (Beaufort scale = 6) and *fresquito* (Beaufort scale = 5)) for the period 1799-1813. The predominant state of the sky is *claro* (clear sky) for the period  
365 1805-1813.

The recovered precipitation data show great consistency with respect to the expected climatology: a seasonal cycle with almost no rainfall in the summer months. The years 1812 and 1813 recorded the maximum accumulated precipitation for the period 1805-1813. Unfortunately, observers only recorded precipitation readings when they occurred. Thus, the non-  
370 registration of 0 mm of rain entails a certain indeterminacy in these data. In addition, we have found a pair of contradictory data between total monthly precipitation (0 mm) and number of rainy days (other than zero) in the months of February and March 1813.

We would like to see this dataset used for different applications. We are confident that providing free access to this new historical dataset will foster such additional assessments by other researchers.

### **Dataset access**

375 The SF1799-2013 dataset recovered is publicly available at the Zenodo open data repository at <https://doi.org/10.5281/zenodo.7104289>.

## Author contribution

N.B.-P. made the data analysis and wrote the first draft of the manuscript. M.C.G and J.M.V. located the original data and initiated the collaboration. R.T., J.M.V. and M.C.G. wrote and revised the manuscript.

## 380 Competing interests

The authors declare that they have no conflict of interest.

## Acknowledgments

This research was supported by the Economy and Infrastructure Counselling of the Junta of Extremadura through project IB20080, and grant GR21080 (co-financed by the European Regional Development Fund). The research of N. Bravo-  
385 Paredes has been supported by the predoctoral fellowship PRE2018-084897 from Agencia Estatal de Investigación (Ministerio de Ciencia, Innovación y Universidades) of the Spanish Government.

## References

- Alcoforado, M. J., Vaquero, J. M., Trigo, R. M., and Taborda, J. P.: Early Portuguese meteorological measurements (18th century), *Clim. Past*, 8, 353–371, <https://doi.org/10.5194/cp-8-353-2012>, 2012.
- 390 Auer, I., Böhm, R., Jurkovic, A., Lipa, W., Orlik, A., Potzmann, R., Schöner, W., Ungersböck, M., Matulla, C., Briffa, K., Jones, P., Efthymiadis, D., Brunetti, M., Nanni, T., Maugeri, M., Mercalli, L., Mestre, O., Moisselin, J.-M., Begert, M., Müller-Westermeier, G., Kveton, V., Bochnicek, O., Stastny, P., Lapin, M., Szalai, S., Szentimrey, T., Cegnar, T., Dolinar, M., Gajic-Capka, M., Zaninovic, K., Majstorovic, Z., and Nieplova, E.: HISTALP – historical instrumental climatological surface time series of the Greater Alpine Region, *Int. J. Climatol.*, 27, 17–46, <https://doi.org/10.1002/joc.1377>, 2007.
- 395 Barriendos, M., Martín-Vide, J., Peña, J. C., and Rodríguez, R.: Daily Meteorological Observations in Cádiz - San Fernando. Analysis of the Documentary Sources and the Instrumental Data Content (1786-1996), *Clim. Change*, 53, 151–170, <https://doi.org/10.1023/A:1014991430122>, 2002.
- Blišňák, V., Valente, M. A. and Bethke, J.: Homogenization of time series from Portugal and its former colonies for the period from the late 19th to the early 21st century, *Int. J. Climatol.*, 35, 2400-2418, <https://doi.org/10.1002/joc.4151>, 2015.
- 400 Brönnimann, S., Brugnara, Y., Allan, R. J., Brunet, M., Compo, G. P., Crouthamel, R. I., Jones, P. D., Jourdain, S., Luterbacher, J., Siegmund, P., Valente, M. A., and Wilkinson, C. W.: A roadmap to climate data rescue services, *Geosci. Data J.*, 5, 28–39, <https://doi.org/10.1002/gdj3.56>, 2018.
- Brönnimann, S., Allan, R., Ashcroft, L., Baer, S., Barriendos, M., Brázdil, R., Brugnara, Y., Brunet, M., Brunetti, M., Chimani, B., Cornes, R., Domínguez-Castro, F., Founda, D., Gergis, J., Grab, S., Hannak, L., García-Herrera, R., Huhtamaa,

- 405 H., Jacobsen, K.S., Jones, P., Jourdain, S., Kiss, A., Lin, K.E., Lorrey, A., Lundstad, E., Luterbacher, J., Moberg, A., Mauelshagen, F., Maugeri, M., Maughan, N., Neukom, R., Nicholson, S., Noone, S., Nordli, Ø., Ólafsdóttir, K.B., Pearce, P.R., Pfister, L., Pribyl, K., Przybylak, R., Pudmenzky, C., Rasol, D., Reichenbach, D., Řezníčková, L., Rodrigo, F.S., Rohde, R., Rohr, C., Skrynyk, O., Slonosky, V., Thorne, P., Valente, M.A., Vaquero, J.M., Westcott, N.E., Williamson, F., and Wyszyński, P.: Unlocking pre-1850 instrumental meteorological records: A global inventory, *Bull. of the Am. Meteorol.*
- 410 Soc. 100(12), ES389–ES413, <https://doi.org/10.1175/BAMS-D-19-0040.1>, 2019.
- Brugnara, Y., Pfister, L., Villiger, L., Rohr, C., Isotta, F. A., and Brönnimann, S.: Early instrumental meteorological observations in Switzerland: 1708-1873, *Earth Syst. Sci. Data*, 12, 1179–1190, <https://doi.org/10.5194/essd-12-1179-2020>, 2020.
- Calvo, N., Gallego, D., García-Herrera, R., Gullón, A., Portela, M. J., Prieto, M. R., and Ribera, P.: El clima de Cádiz en la
- 415 primera mitad del siglo XIX según los parte de la Vigía, edited by: García-Herrera, R., Servicio de Publicaciones de la Fundación Unicaja, Málaga, Spain, 2008.
- Camuffo, D. and Jones, P.: Improved Understanding of Past Climatic Variability from Early Daily European Instrumental Sources, in: Improved Understanding of Past Climatic Variability from Early Daily European Instrumental Sources, vol. 53, edited by: Camuffo, D., and Jones, P., Springer, Dordrecht, 1–4, [https://doi.org/10.1007/978-94-010-0371-1\\_1](https://doi.org/10.1007/978-94-010-0371-1_1), 2002.
- 420 Cornes, R. C., Jones, P. D., Briffa, K. R., and Osborn, T. J.: A daily series of mean sea-level pressure for Paris, 1670-2007, *Int. J. Climatol.*, 32, 1135–1150, <https://doi.org/10.1002/joc.2349>, 2012.
- Domínguez-Castro, F., Vaquero, J. M., Rodrigo, F. S., Farrona, A. M. M., Gallego, M. C., García-Herrera, R., Barriendos, M., and Sanchez-Lorenzo, A.: Early Spanish meteorological records (1780-1850), *Int. J. Climatol.*, 34, 593–603, <https://doi.org/10.1002/joc.3709>, 2014.
- 425 Domínguez-Castro, F., Vaquero, J. M., Gallego, M. C., Farrona, A. M. M., Antuña-Marrero, J. C., Cevallos, E. E., García-Herrera, R., de la Guía, C., Mejía, R. D., Naranjo, J. M., Prieto, M. R., Ramos Guadalupe, L. E., Seiner, L., Trigo, R. M., and Villacís, M.: Early meteorological records from Latin- America and the Caribbean during the 18 th and 19 th centuries, *Sci. Data*, 4, 1–10, <https://doi.org/10.1038/sdata.2017.169>, 2017.
- Dunnington, D.: Ggsatial: spatial data framework for ggplot2, R package version 1.1.1, [https://CRAN.R-](https://CRAN.R-project.org/package=ggsatial)
- 430 [project.org/package=ggsatial](https://CRAN.R-project.org/package=ggsatial), 2020.
- Farrona, A. M. M., Trigo, R. M., Gallego, M. C., and Vaquero, J. M.: The meteorological observations of Bento Sanches Dorta, Rio de Janeiro, Brazil: 1781-1788, *Clim. Change*, 115, 579–595, <https://doi.org/10.1007/s10584-012-0467-8>, 2012.
- García-Herrera, R., Prieto, L., Gallego, D., Hernández, E., Gimeno, L., Können, G., Koek, F., Wheeler, D., Wilkinson, C., Prieto, M., Báez, C., and Woodruff, S: CLIWOC multilingual meteorological dictionary, 2003.
- 435 Jones, P. D., Davies, T. D., Lister, D. H., Slonosky, V., Jónsson, T., Barring, L., Jónsson, P., Maheras, P., Kolyva-Machera, F., Barriendos, M., Martin-Vide, J., Rodriguez, R., Alcoforado, M. J., Wanner, H., Pfister, C., Luterbacher, J., Rickli, R., Schuepbach, E., Kaas, E., Schmith, T., Jacobep, J., and Beck, C.: Monthly mean pressure reconstructions for Europe for the

- 1780-1995 period, *Int. J. Climatol.*, 19, 347–364, [https://doi.org/10.1002/\(SICI\)1097-0088\(19990330\)19:4<347::AID-JOC363>3.0.CO;2-S](https://doi.org/10.1002/(SICI)1097-0088(19990330)19:4<347::AID-JOC363>3.0.CO;2-S), 1999.
- 440 Können, G. P. and Brandsma, T.: Instrumental pressure observations from the end of the 17th century: Leiden (The Netherlands), *Int. J. Climatol.*, 25, 1139–1145, <https://doi.org/10.1002/joc.1192>, 2005.
- Lafuente, A. and Sellés, M.: *El Observatorio de Cádiz (1753-1831)*, Ministerio de Defensa, Instituto de Historia y Cultura Naval, Madrid, 1988.
- Luterbacher, J., Rickli, R., Tinguely, C., Xoplaki, E., Schüpbach, E., Dietrich, D., Hüsler, J., Ambühl, M., Pfister, C., Beeli, P., Dietrich, U., Dannecker, A., Davies, T. D., Jones, P. D., Slonosky, V., Ogilvie, A. E. J., Maheras, P., Kolyva-Machera, F., Martin-Vide, J., Barriendos, M., Alcoforado, M. J., Nunes, M. F., Jónsson, T., Glaser, R., Jacobeit, J., Beck, C., Philipp, A., Beyer, U., Kaas, E., Schmith, T., Barring, L., Jönsson, P., Rácz, L., and Wanner, H.: Monthly mean pressure reconstruction for the Late Maunder Minimum Period (AD 1675-1715), *Int. J. Climatol.*, 20, 1049–1066, [https://doi.org/10.1002/1097-0088\(200008\)20:10<1049::AID-JOC521>3.0.CO;2-6](https://doi.org/10.1002/1097-0088(200008)20:10<1049::AID-JOC521>3.0.CO;2-6), 2000.
- 445 Morozova, A. L. and Valente, M. A.: Homogenization of Portuguese long-term temperature data series: Lisbon, Coimbra and Porto, *Earth Syst. Sci. Data*, 4, 187–213, <https://doi.org/10.5194/essd-4-187-2012>, 2012.
- Pfister, L., Hupfer, F., Brugnara, Y., Munz, L., Villiger, L., Meyer, L., Schwander, M., Isotta, F. A., Rohr, C., and Brönnimann, S.: Early instrumental meteorological measurements in Switzerland, *Clim. Past*, 15, 1345–1361, <https://doi.org/10.5194/cp-15-1345-2019>, 2019.
- 450 Picas, J., Grab, S., and Allan, R.: A 19th century daily surface pressure series for the Southwestern Cape region of South Africa: 1834–1899, *Int. J. Climatol.*, 39, 1404–1414, <https://doi.org/10.1002/joc.5890>, 2019.
- Pujazón, C.: *Resumen de las observaciones pluviométricas efectuadas en los años 1805 a 1899*, Instituto y Observatorio de Marina, San Fernando, Cádiz, 1899.
- Rodrigo, F. S.: Completing the early instrumental weather record from Cádiz (Southern Spain): New data from 1799 to 1803, *Clim. Change*, 111, 697–704, <https://doi.org/10.1007/s10584-011-0174-x>, 2012.
- 460 Rodrigo, F. S.: Early meteorological data in southern Spain during the Dalton Minimum, *Int. J. Climatol.*, 39, 3593–3607, <https://doi.org/10.1002/joc.6041>, 2019.
- Slonosky, V.: The Meteorological Observations of Jean-François Gaultier, Quebec, Canada: 1742–56, *J. Clim.*, 16, 2232–2247, [https://doi.org/10.1175/1520-0442\(2003\)16<2232:TMOOJG>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)16<2232:TMOOJG>2.0.CO;2), 2003.
- 465 Slonosky, V. C., Jones, P. D., and Davies, T. D.: Instrumental pressure observations and atmospheric circulation from the 17th and 18th centuries: London and Paris, *Int. J. Climatol.*, 21, 285–298, <https://doi.org/10.1002/joc.611>, 2001.
- Sousa, R.: *Notas para una Climatología de Cádiz*, Ministerio de Transportes, Turismo y Comunicaciones, Madrid, 1987.
- Valler, V., Franke, J., Brugnara, Y., and Brönnimann, S.: An updated global atmospheric paleo-reanalysis covering the last 400 years, *Geosci. Data J.*, 1–19, <https://doi.org/10.1002/gdj3.121>, 2021.
- 470 Vaquero, J. M., Bravo-Paredes, N., Obregón, M. A., Carrasco, V. M. S., Valente, M. A., Trigo, R. M., Domínguez-Castro, F., Montero-Martín, J., Vaquero-Martínez, J., Antón, M., García, J. A., and Gallego, M. C.: Recovery of early

meteorological records from Extremadura region (SW Iberia): The ‘CliPastExtrem’ (v1.0) database, *Geosci. Data J.*, 1–14, <https://doi.org/10.1002/gdj3.131>, 2022.

475 Wheeler, D.: Early instrumental weather data from Cadiz: A study of late eighteenth and early nineteenth century records, *Int. J. Climatol.*, 15, 801–810, <https://doi.org/10.1002/joc.3370150707>, 1995.

WMO: Guide to meteorological instruments and methods of observation; WMO-No. 8: Measurement of Meteorological Variables, 525–552 pp., 2018.

Zaiki, M., Kimura, K., and Mikami, T.: A statistical estimate of daily mean temperature derived from a limited number of daily observations, *Geophys. Res. Lett.*, 29(18), 39.1-39.4, <https://doi.org/10.1029/2001GL014378>, 2002.

480 Zaiki, M., Können, G. P., Tsukahara, T., Jones, P. D., Mikami, T., and Matsumoto, K.: Recovery of nineteenth-century Tokyo/Osaka meteorological data in Japan, *Int. J. Climatol.*, 26, 399–423, <https://doi.org/10.1002/joc.1253>, 2006.