

1 The weather diary of Georg Christoph Eimmart for Nuremberg, 1695-1704

2
3 Stefan Brönnimann^{1,2}

4 ¹*Oeschger Centre for Climate Change Research, University of Bern, Switzerland*

5 ²*Institute of Geography, University of Bern, Switzerland*

6 stefan.broennimann@giub.unibe.ch

7 8 **Abstract**

9 Data from weather diaries have long been used to reconstruct past climate. However, they could also
10 be used to reconstruct past weather. Weather reconstructions could help to better understand the
11 mechanisms behind, and impacts of, climatic changes. However, reconstructing the day-to-day
12 weather requires many diaries from different regions covering the same period, ideally combined with
13 instrumental measurements. In this paper, I describe the weather diary of Georg Christoph Eimmart
14 from Nuremberg, covering the period 1695 to 1704, which was particularly cold in Europe. The diary
15 was imaged from the Russian National Library in St. Petersburg and then digitized. It contains twice
16 daily weather conditions in symbolic form, wind direction, and information on precipitation and
17 temperature in text form. Symbols changed during the first two years, after which a much reduced
18 (and stable) set of symbols was used. Re-coding all days according to the later set of symbols, I find
19 no signs of inconsistency over time in symbols, wind direction, and precipitation information extracted
20 from the text. Comparisons with other sources confirm the day-to-day weather information in the
21 diary. For instance, the wind direction in Nuremberg agrees with the daily pressure gradient between
22 Jena and Paris. Three case studies further confirm the meteorological reliability of the information.
23 This is shown on behalf of an eight-day sequence of stormy weather in 1702, a study of the severe
24 winter of 1697/8, and of the summer of 1695, which was cold and wet, possibly related to tropical
25 volcanic eruptions. The examples underline the consistency of the weather diary with other
26 information and suggest that weather reconstructions as far back as the late 17th century might become
27 possible. However, the spatial information is limited, and any approach arguably needs to make good
28 use of the temporal sequence of information.

29 **1. Introduction**

30 For decades, historians have used data from weather diaries to reconstruct past climate, i.e., to
31 generate monthly or seasonal index series (e.g., Pfister, 1999, see overview in Nash et al., 2021).
32 However, they could also be used to reconstruct past weather day by day. In fact, Manley (1975)
33 described the daily weather during the cold winter 1683/4 based on instrumental data and data from
34 weather diaries. Kington (1988) and Lamb (1991) reconstructed daily weather types and drew daily
35 weather charts for periods in the 18th, 17th, and even 16th century. They combined sparse observations
36 with expert interpretation in a reproducible way (Kington 1988).

37 As weather extremes and changes in weather have come into focus of climate science, reconstructing
38 daily weather is again considered an important goal. García-Herrera et al. (2007), Wheeler et al.
39 (2009) and numerous others have demonstrated the value of carefully reconstructing past extreme
40 weather events. However, the subjective nature of these reconstructions is not without problems (e.g.,
41 Jones et al., 2014, and references therein). Furthermore, to produce complete daily data series,
42 automated and objective methods are used rather than time consuming expert interpretations. Daily

43 weather types (Schwander et al., 2017) and gridded daily weather reconstructions (Imfeld et al., 2022)
44 have been performed for Switzerland back to 1763 and for Europe for the 1780s (Pappert et al., 2022).
45 Other daily indices such as the wind direction over the English Channel reach further back, to the late
46 17th century (Wheeler et al., 2010). Cornes et al. (2012a,b) used daily data of sea-level pressure (SLP)
47 from London and Paris to analyse atmospheric circulation and storminess. There are many more
48 examples for analyses of daily weather 300 years back (e.g., Brázdil et al., 2008, Filipiak et al., 2019),
49 a review of approaches is given in Brönnimann (2022). In addition to traditional statistical and
50 numerical methods, new approaches such as machine learning (Brohan, 2022) could possibly replace
51 the expert approach pioneered by Manley, Kington, and Lamb.

52 The success of any weather reconstruction approach ultimately depends on the available weather data.
53 From the turn of the 17th to the 18th century, several weather diaries are available. The diary from
54 Johann Heinrich Fries in Zurich covering 1684-1718 (Pfister, 1977) can be downloaded from EURO-
55 CLIMHIST (Pfister et al. 2017). The diary of the Kirch family in Leipzig (and Guben) and later
56 Berlin, covers 1677-1774 (Herbst, 2022) and was imaged by the author. Further diaries such as that of
57 David Grebner in Wroclaw covering 1692-1710 (Przybylak and Pospieszyńska, 2010) and Joseph
58 Dietrich in Einsiedeln covering 1670-1704 (Rohr and Schwarz-Zanetti, 2022) are under digitization
59 (not considered here). There are also a number of instrumental records from the late 17th and early
60 18th century (see Brönnimann et al., 2019a, Lundstad et al., 2022), many of which also have weather
61 descriptions. Combining all these data sets, reconstructing daily weather over Europe back to the late
62 17th century could become possible.

63 Here I add another weather diary, namely that of Georg Christoph Eimmart, founder of the first
64 Nuremberg (Nürnberg) astronomical observatory. His weather diary covers the years 1695-1704. This
65 paper describes the diary, its digitisation and (as the diary is mostly kept in symbolic form)
66 categorization. I then compare Eimmart's observations with other sources of daily weather
67 information.

68 The turn of the 17th to the 18th century is not only interesting as a test case as to how far back daily
69 weather reconstruction can reach, but it is also climatically interesting. It fell into the so-called "Late
70 Maunder Minimum" (Luterbacher et al., 2001), with particularly low temperatures in Europe
71 coinciding with low solar activity. At the same time, several volcanic eruptions (Hekla and Serua in
72 1693, Komagatake in 1694) might have affected climate (see Burgdorf, 2022).

73 The paper is organised as follows. Section 2 provides background about the observer and the diary.
74 Section 3 then describes the digitization and the data used for comparison. Results are presented in
75 Section 4. A brief discussion then follows in Section 5 and conclusions are drawn in Section 6.

76 **2. Georg Christoph Eimmart and his weather diary**

77 Georg Christoph Eimmart (1638-1705) is known as the founder of the first Nuremberg astronomical
78 observatory (the following text is based on Gaab, 2005, 2022). Eimmart attended the "Gymnasio
79 poetico" at Regensburg and enrolled in 1655 at the University of Jena, where he studied mathematics.
80 In 1658 he returned to Regensburg. After the death of his father, he moved to Nuremberg (following
81 his sister) around 1660. He worked as an engraver and got involved in the management of the
82 Academy of Painting founded in 1662, and from 1699 until shortly before his death he was director of
83 this institution.

84 Using the money earned with his artistic activity, Eimmart set up an observatory at Vestnertor, north
85 of Nuremberg Castle (Fig. 1) in 1678. At the end of the 17th century this observatory was the most
86 well-known in Germany (Gaab, 2005). During special celestial events, the observatory was opened to

87 the public. In 1699, Eimmart was admitted to the Parisian Academy of Sciences, and at Leibniz's
88 suggestion two years later, he was also admitted as an external member of the Prussian Academy of
89 Sciences.

90 The written legacy of Georg Christoph Eimmart can be found today in the Russian National Library in
91 St. Petersburg. Vol. 40 of that legacy is the “Diarium tempestatum” (see also Gaab, 2022). It covers
92 the period 1 January 1695 to 25 November 1704 on 120 pages. The first three months are written in
93 ink, the remaining pages in pencil. The diary is structured in tables and contains only observations, no
94 measurements (although a sketch of an instrument is found on a verso page, Fig. 2; all other verso
95 pages are empty). The table is structured in columns, which on the first page are labelled as “Dies;
96 Qualitas Aeris; Temperam: cal: et frig:; Ventus”. There is an apparent gap from 19-28 February 1700
97 („Incipit Dies 1. Martii“), which is due to the change from the Julian to the Gregorian calendar.
98 Another (real) gap of four weeks occurs in January/February 1703.



99 **Fig. 1.** (left) Map of Nuremberg, 1642 (engraving by Matthäus Merian from *Germaniae, Edition Topographia*
100 *Franconiae*, 1642; Wikimedia Commons, public domain) with the location of the observatory. (right) Location of
101 other weather observation series used in this study (dark green: instrumental measurements, light green:
102 observations).
103


104 An interesting aspect of the diary is that it is in its majority symbolic. This means that the entry is
105 mostly given as symbol, sometimes in words, sometimes both. In the first two years, each column (one
106 month, August 1695, has four columns, all others have three) has a distinct set of symbols that is used
107 (see Fig. 3). Particularly in the first two years, there are often several symbols in the same cell. After
108 2-3 years, however, the distinction between columns 1 and 2 gets lost, while the wind column remains
109 unchanged until the end of the diary. Some of the symbols, particularly those used in the second
110 column (on temperature), vanish and those used in the first column also appear in the second. A much-
111 reduced set of symbols is used in both columns after ca. 1697. There are mostly just three symbols:
112 full sun, upper half of sun, and long dash.

113 Wind is given in cardinal/intercardinal direction, plus a symbol probably referring to changing winds.
114 Sometimes several wind directions are mentioned in the same cell (see example in Fig. 2), which
115 might indicate several observations per day.


116 In addition to symbols and wind direction, there are also occasional latin expressions such as „nebula“,
117 “pluvial per intervalla”, or “nix”. They describe phenomena related to precipitation (rain snow,
118 thunderstorm, lightning, rainbows), wind speed, and temperature. Occasionally other aspects are

119 mentioned (changeable weather, clarity of the sky). Words are sometimes written across both columns
 120 (sometimes even also across the wind column), which leads to the hypothesis that, from ca. 1697
 121 onward (after the same symbols are used in both columns), Eimmart wrote twice daily observations,
 122 where the first column refers to morning and the second to afternoon. This is supported by the fact that
 123 “tonitru” (thunders) appears only 6 times in the first, but 33 times in the second column. Likewise,
 124 “aestus” (heat) appears 23 times in the first and 47 times in the second columns. However, there is no
 125 text to prove this hypothesis.

126 The text entries are not independent of the symbols. In about half of the cases, text entries replace
 127 symbolic entries, i.e., these fields then do not have a symbol on the sky conditions. For the other half
 128 of cases with text entries these complement existing symbols. Sometimes the text refers specifically to
 129 precipitation that has fallen during the night. At few instances, an additional row is even added
 130 between two rows noting nocturnal rain or nocturnal storms, or this is made clear by subdividing cells.
 131 Overall, there are slightly more text entries in the second column than in the first. Text entries in the
 132 third column are rare.



133
 134 **Fig. 2.** Eimmart’s diary for November 1696 from the National Library of Russia, St. Petersburg. The left page
 135 shows sketches of scientific instruments (© National Library of Russia, St. Petersburg).



136

137 **Fig. 3.** Symbols used in the weather diary.

138 **3. Digitising and formatting the diary and comparison to other data sources**

139 Before digitizing, an inventory of all symbols appearing in the diary was compiled (Fig. 3), and a code
140 was assigned to each symbol. Then symbols were digitized as codes. For the text I use an additional
141 column. I transcribed the latin words as good as possible, some illegible words are marked with “\$”.
142 Eimmart abbreviated some words with a symbol resembling “y”, which I transcribed as “y” (e.g.,
143 “frigy”). Words written over two fields are assigned to both fields with a corresponding bracket (e.g.,
144 „pluvius[, and]pluvius“; for the special case of August 1695, with 4 columns, I kept the three column
145 structure but assigned words across several columns only to one, with a note “[2 columns]”). A pipe
146 symbol “|” is used to indicate line breaks within a cell. In case of nocturnal weather, when the diary
147 makes it clear to which night this refers, I add it to the following day (the weather diary also mostly
148 does it this way) with a note “[previous night: ...]”. Within the text, Eimmart sometimes uses a
149 symbol for the Sun, which I transcribed as “[solaris]”.

150 From this raw transcription I generated several derived variables. First, I categorized the text entries
151 related to precipitation into snow (“nix”, “nivigit” or similar), rain (“pluvia”), or rain and snow. In the
152 category rain I also included expressions such as “tonitru” (thunder) or “nebula pluviosa” (rainy fog),
153 but not “tempestas” (storm). To these I also added symbols 7 to 9 (rain) and 11 (snow) of Fig. 3,
154 column 1, as well as symbols 7 (snow), 8 and 9 (rain) of Fig. 3, column 2. Note that these almost
155 exclusively appear in the first 9 months. This variable is called precipitation.

156 Second, I formed a unified code based on the three main symbols used after 1697, namely “sunny”
157 (full sun), “partly sunny” (half sun), and “cloudy” (horizontal line). For this, I grouped symbols 1, 2
158 (column 1) and 5 (column 2) to the category “sunny”, symbols 3 to 5 (column 1) as well as any
159 combination of a symbol 1-5 (column 1) with another symbol as category “partly sunny”, and
160 categories 6 and 12 (column 1) to category “cloudy”. All days for which the diary indicates a weather
161 symbol (76% in column 1, 60% in column 2) are thus assigned one of the three categories. This
162 variable is called “WeatherSymb”.

163 Third, for most of the days with missing symbols, there is a text entry. In a further step, I also
164 generated a code for these entries. The terms “serenum” (clear) and “sunidy” (sunny) were coded as
165 “sunny”, “coelum varium” (changeable), “pluvia per intervalla” (occasional rain), “pluvia minuta”
166 (little rain), “tonitru” (thunder), and symbol 8 (column 2) were coded as “partly sunny”, and the terms
167 “pluvia”, “pluvia tota dia”, and “nebula” as well as symbol 9 (column 2) were coded as cloudy. In this
168 way, 97% (80%) of the days in columns 1 and 2, respectively, could be coded. The smaller amount in
169 column 2 is due to the fact that many of the text entries refer to temperature and winds. This variable
170 is called “WeatherSymbText”.

171 Eimmart’s diary entries were then compared with daily weather information from other sources (see
172 Fig. 1 for locations). This includes a weather diary from Zurich, pressure measurements from London
173 (Cornes, 2012a), Paris (Cornes, 2012b), Leiden, Halle, and Jena (Lundstad et al., 2022; see
174 Supplementary Material), temperature measurements from Paris (Rousseau, 2009, Pliemon et al.,
175 2022), Berlin, Halle, and Kaliningrad (Lundstad et al., 2022; see Supplementary Material), as well as
176 wind direction from ships on the Channel (Wheeler et al., 2010, Barriopedro et al., 2014). Note that,
177 first, many of the series only overlap partly with the Eimmart diary, thus limiting comparisons, and,
178 second, their quality is mostly unknown. For instance, the Halle records of both temperature and
179 pressure did not seem to be usable in their present form and were discarded. The following
180 comparisons address the internal temporal consistency, the temporal agreement of the diary with other

181 time series, the spatial consistency across Europe, and the consistency of daily wind direction with a
182 large-scale pressure gradient.


183 More comparability options would exist on a monthly scale by comparing with monthly
184 reconstructions. However, it is not straight forward to aggregate the daily weather information to a
185 monthly level, and the focus of this paper is on the daily scale. However, I compared the weather diary
186 with other documentary series (Burgdorf et al., 2022) and with monthly climate reconstructions for
187 two specific cases, namely the summer of 1695 and the winter of 1697/98. For these case studies I
188 used the ensemble mean of the reconstruction EKF400v2, which is a global, 3-dimensional climate
189 reconstruction based on data assimilation (Valler et al., 2022).

190 4. Results

191 4.1. Time consistency of symbols, wind direction, and text

192 In first step I analysed the frequency of each symbol per month, over both columns. Due to the large
193 number of symbols used, especially at the beginning, I ignored the temperature related symbols
194 (symbols 1-4, column 2), which only appear in the first 4 months, and grouped the symbols remaining
195 related to the general weather characteristics and the sky into four main categories (see Fig. 4). I then
196 counted each use of a symbol and divided it by the total number of symbols in that month. There
197 might be several symbols per day, or just one. Note also that text is ignored here.


198 The results (Fig. 4a) show what was already observed during the digitization, namely that in the
199 beginning, many different symbols appeared while later basically three symbols were used. The grey
200 category of symbols vanishes almost completely and from around 1698 onward, the use of the
201 symbols is rather consistent. Figure 4b shows the results obtained when simply ignoring the grey
202 category of symbols. Their frequency over time changes less, but many days have no category. The
203 corresponding figure for variable “WeatherSymb” (Fig. 4c) shows slightly different behaviour in the
204 first years compare to later. When also text entries are categorized (variable “WeatherSymbText” Fig.
205 4d), no obvious inhomogeneity is seen anymore. The figure resembles Fig. 4b, but now almost all days
206 have a category. Note that cloudy conditions are more frequent because most text entries concern rain.



208 **Fig. 4.** Frequency of occurrence of weather symbols per month relative to all symbols in that month. **a** all
 209 symbols (grouped into four categories), **b** only the first three categories, **c** variable “WeatherSymb”, **d** variable
 210 “WeatherSymbText”.

211 Similar as for the weather symbols, I also analysed the wind direction. I counted the letters for the
 212 cardinal directions (N, E, S, W as well as one letter for mixed), divided the counts by the number of
 213 letters per day, and then averaged the number per month. Note that in this procedure NW counts as
 214 two letters (each weighted half), N as one. This allows a first, albeit simple visualization (Fig. 5a).
 215 Results show variations in occurrence, but no obvious inhomogeneity is seen in this plot.

216 The same was also done for the variable precipitation with its categories “snow”, “rain and snow”, and
 217 “rain” (Fig. 5b). Variability in precipitation is high, but once again there is no evidence for an
 218 inhomogeneity. The figure indicates a decrease of rain days over the 10 years. However, this might be
 219 a true climatic signal. The large spike in the beginning of the series, in summer 1695, will be analysed
 220 later and is arguably real.



221 **Fig. 5.** Frequency of occurrence of **a** cardinal wind directions per month and **b** days with snow, rain and snow, or
 222 rain.
 223

224 *4.2. Analysis of the wind direction*

225 As a next step I compared the daily wind direction at Nuremberg with the daily SLP gradient between
 226 Jena and Paris in the year 1702 (the only year for which data from Jena are available). A positive
 227 gradient is expected to correlate with winds from the South or East, whereas a negative gradient is
 228 expected to correlate with winds from the North or West. I therefore grouped the winds accordingly
 229 (S, SE, E, SW vs. W, NW, N, NE). Most of the days had only one wind direction. If more than one
 230 was noted, I excluded days that would fall into both categories.

231 To calculate the pressure gradient between Jena and Paris, I reduced the Jena data to sea level. As no
 232 temperature information was available, I assumed a sinusoidal seasonal cycle of temperature varying
 233 between -4 °C (in January) and 20 °C (in July). As the thus obtained SLP data were clearly too low in


234 Jena, I added the mean difference between Jena and Paris such that both series have the same mean. I
 235 analysed the data using a contingency table and by stratifying the SLP difference according to the
 236 wind direction.

237 The contingency table (Table 1) clearly shows a very strong association between pressure gradient and
 238 wind direction in the expected sense. A statistical test (Fisher's exact t) confirms the high significance
 239 of the results. However, while there are more cases with a negative than with a positive SLP difference
 240 (note that the average difference is zero), there are clearly more cases with an easterly-to-south-
 241 westerly wind than a westerly-to-north-easterly wind. The histogram of SLP difference (Fig. 6) also
 242 clearly shows a difference in the SLP gradient depending on the wind direction. Note that deviations
 243 are expected. Apart from the measurement errors and errors in the reduction to SLP, it should be noted
 244 that the SLP gradient may not be the best proxy for wind. Furthermore, Paris is relatively far away,
 245 and the wind over land is not geostrophic. Thermotopographic winds or topographically channelled
 246 winds may overlay the large-scale flow. Wind is a variable with a high variability, and observations
 247 provide only a snapshot. Note also that the time of day of observations is not known. In light of these
 248 uncertainties, the clear presence of a signal is therefore encouraging.

249 **Table 1.** Contingency table of wind observations in Nuremberg and SLP differences between Jena and Paris. The p-value
 250 refers to Fisher's exact t.

	E/SE/S/SW	W/NW/N/NE	Sum
$\Delta\text{SLP}>0$	118	40	158
$\Delta\text{SLP}<0$	72	98	170
Sum	190	138	$p<0.0001$

251



252

253 **Fig. 6.** Histograms of SLP difference between Jena and Paris stratified by the wind direction in Eimmart's diary.


254 *4.3. Case studies*

255 *4.3.1. Weather maps for October/November 1702*

256 Next I analysed the entries in Eimmart's diary together with all other available information for an 8-
 257 day sequence in 1702. The sequence was chosen as a strong storm was noted in this period in the
 258 observations from Kaliningrad in the night from 31 October to 1 November (Anonymous, 1703). Note
 259 that the year 1702 is arguably the year with the best data coverage within the 1695-1704 period.
 260 Instrumental data are available from Jena, Paris, London, Berlin and Kaliningrad, weather
 261 observations from Jena, Kaliningrad, Zürich, Nuremberg, and the English Channel. For display
 262 purposes, I deseasonalised (by fitting the first two harmonics of the seasonal cycle) and standardized
 263 the instrumental data with respect to the year 1702 and expressed the anomalies in standard deviations.
 264 For Berlin where the observation hours change rapidly, I only used the 8 days displayed, chose the
 265 observation closest to 8 in the morning and standardized the temperatures.

266 The sequence of maps (Fig. 7) starts with mild temperatures, sunny or changeable weather and
 267 moderately low pressure across Europe. The next day pressure increased in Paris and Jena, it remained
 268 rather warm. Pressure remained low over Kaliningrad. The next three days saw a clear pressure

269 increase over London, Paris, and Jena, with westerly flow. Temperatures were high especially on 30
 270 October. Pressure and temperature then both decreased on the day of the storm. Pressure increased
 271 right after the passing of the storm, accompanied by a marked temperature decrease and mostly sunny
 272 weather.



273
 274 **Fig. 7.** Maps of pressure, wind, and temperature for an eight-day sequence in 1702. A vertical sequence of
 275 symbols at the same location indicates sub-daily data (progressing from top to bottom).

276 While this sequence is plausible, the spatial information is insufficient to draw weather maps. There
 277 could have been a quick succession of passing depressions over Northern Europe, that may have gone
 278 unnoticed. However, the maps depict the high-pressure system to the south. Also, the passage of a cold
 279 front on 1 Nov is clearly seen.

280 4.3.2. The harsh winter of 1687/8.

281 The second case study relates to the harsh winter of 1697/8. This winter is particularly well known in
 282 England (Kington, 1999), where it was the coldest among six very cold winters in the 1690s.
 283 However, the winter was also cold over Central Europe (Pfister and Wanner, 2021). A number of
 284 European weather series exist on a daily scale for this winter, including pressure in London, Paris, and
 285 Leiden, temperature in Paris, wind over the English Channel, cloud motion and cloud cover from
 286 Paris, and weather observations from Zurich and Nuremberg. All observations are shown as time
 287 series in Fig. 8.

288 Temperature measurements in Paris as well as notes on temperature in the Zurich series point to a
 289 sequence of many cold spells during the entire winter and spring. For Nuremberg, notes on
 290 temperature are sparser, but they also agree with the two other series. According to Kington (1999),
 291 the first snow in London fell on 24 November. In Nuremberg and Zurich snowfall is reported on 22
 292 November. We also see periods of higher temperature and thawing weather (or rainfall), such as in
 293 early December. A snowy and cold period follows at all sites in mid-December. January then was
 294 particularly cold in Paris, London, and Zurich. A notable pressure drop occurred in Paris on 1 February
 295 1698, arguably associated with a warm front (Kington, 1999). Temperatures increased everywhere,

296 and precipitation fell as rain in Nuremberg (note that the temperature increase here occurred ca. two
297 days later than in Paris). The first half of March was again cold at all sites. Cold spells again occurred
298 in late April and early May. In fact, May 1698 still is the coldest May on record in the Central England
299 temperature series. This brief analysis shows that the daily series are consistent with each other. Even
300 wind directions agree well for locations close to each other (e.g., English Channel and Paris).

301
302
303





Fig. 8. Daily weather in the winter 1697/8. For Nuremberg, 1 to 3 marks the columns, subdivided into text entries that were transformed into symbols (upper line) and symbolic entries (lower line).

304 As a further opportunity for comparison, I considered the chronicle by Johann Laurentz Bünti (Bünti,
 305 1973) from central Switzerland, which points to snow fall on 3 and 8 May and again on 21 May
 306 («Hierauff folget den 3. May ein Schnee [...] Den 8. ten May hat man wiederum im ganzen Boden
 307 Schnee.»). Bünti writes that precipitation was very high and that it fell as snow in the mountains, such
 308 that there was more snow in the mountains in May 1698 than in many winters.

309 To further investigate this winter, I analysed temperature and pressure anomalies in the monthly
 310 reconstruction EKF400v2 (Fig. 9). The fields are expressed as anomalies from the subsequent 30-yr
 311 period. The Dec.-Feb. patterns in sea-level pressure resembles the positive easterly index in Mellado-
 312 Cano et al. (2020), and in fact easterly wind components are soon in the English Channel winds, Paris
 313 cloud motion, and in Eimmart's observations at Nuremberg . The same figure also shows hand-drawn
 314 pressure maps by Wanner et al. (1995). These fields have no scale (and are here compared with
 315 anomalies). However, analysing the position of highs and lows (or positive and negative anomalies), I
 316 find a mostly good agreement, indicating that the data assimilation approach and an expert approach
 317 give consistent results. Into these we can now embed the weather diaries. The fields show that January
 318 and February were actually even colder in Central Europe than in Western Europe. In January,
 319 Eimmart notes mostly easterly winds and a horizontal line, arguably denoting a persistent stratus.
 320 These features stand out over the 10 year period and are also clearly seen in Fig. 4 (increased blue
 321 area) and Fig. 5 (increased green area denoting easterly winds). This is very well in line with the
 322 monthly charts shown in Fig. 9. The cold spells in May seemed to have had more pronounced effects
 323 in Zurich than in nearby Nuremberg, consistent in the reconstructed fields, the hand-drawn fields, and
 324 the diary entries.



325
 326 **Fig. 9.** Monthly temperature (colours) and pressure anomalies (contour, in hPa) relative to the period 1698/9 to
 327 1727/28 from EKF400v2 (top) and an expert reconstruction of the pressure distribution and main flow for the
 328 same months from Wanner et al. (1995).


329 4.3.3. The summer of 1695

330 Finally, I also analysed the summer of 1695, which is known as a very cold summer (July 1695 was
 331 the second coldest July in the Central England temperature series behind 1816). The summer was also

332 cold in Switzerland. Bünti (1973) writes that 1695 was a late and wet year, with frequent summer
 333 snowfall events in the Alps («Sonsten ist dissess 1695. Jahr ein spätes und nasses Jahr gesein; im
 334 Summer [wurden] oft die Alpen überschnyt»).

335 For this case, I analysed the variables “Precipitation” and “WeatherSymbText” from Eimmart’s diary
 336 as well as temperature and weather conditions and in Zurich in June to August. I compared the
 337 frequencies in 1695 to those obtained in 1696-1704 (Fig. 10). I also compared the result to
 338 documentary climate data for that summer and to EKF400v2. For comparison, I standardized both
 339 based on the 1696-1704 period (EKF400v2 is also shown as anomalies without standardizing).
 340 Although this is a small sample, it is the longest statistical analysis the Eimmart diary allows.

341 The summer of 1695 was clearly less sunny in Eimmart’s diary, while cloudy conditions were more
 342 frequent. Also, there were more days with precipitation in Jun-Aug 1695 than in the average of all
 343 other summers. A very similar behaviour is found in Zurich, where sunny conditions were less
 344 frequent and “changeable” more frequent. A clear signal is also found in the temperature notes of
 345 Zurich. Cold days were about twice as frequent in 1695 as in the reference period, mostly at the
 346 expense of very warm days. For Nuremberg, there are too few temperature notes for an analysis.



347
 348 **Fig. 10.** Frequency of weather descriptions in the diaries of Eimmart (Nuremberg) and Fries (Zurich) for the
 349 summer (Jun-Aug) of 1695 as well as for the remaining 9 summers of Eimmart’s observation period.


350 The two diaries can now be compared to two further sources on that summer: documentary data (Fig.
 351 11, top, Supplementary Material) and reconstructions (Fig. 11, bottom). Both sources show that it was
 352 cold particularly in the northeast of the domain, less so in Western France and Spain. The
 353 documentary data indicate no specifically cold period in Italy, whereas the reconstruction suggest
 354 somewhat lower temperatures (note that the documentary data and reconstructions are largely
 355 independent, only the monthly index series indicated in Fig. 11, top, with “JJA” were assimilated into
 356 EKF400v2). Zurich and Nuremberg were both in a region that was affected by the adverse weather,
 357 although not in the core region of this climatic anomaly.

358 5. Discussion

359 The weather diary by Georg Christoph Eimmart from Nuremberg, covering the period 1695 to 1704,
 360 might be a useful addition to the compilations of existing weather diaries. The diary stands out in that
 361 it is mostly symbolic for sky conditions, complemented with wind direction and text for precipitation
 362 and temperature. Although there are quite large changes in the use of symbols and of text in the first 1-
 363 2 years, derived variables that group sky conditions and precipitation each into 3 categories according

364 to both symbols and text show now sign of inhomogeneity. The same result is found for wind
 365 direction. Observations seem to have been performed in a consistent manner over 10 years.

366 The diary compares well with other sources of information, such as the diary from Fries in Zurich. It
 367 also compares well with instrumental data, as demonstrated by comparing wind direction with a large-
 368 scale SLP gradient. This confirms that the diary not only has useful information on climate, but also
 369 on the daily weather. Combining this diary with other available observation series shows that there is
 370 spatial information in the combination of all weather data. Some features, such as the passing of a cold
 371 front, can be clearly seen, but it is still difficult to draw detailed weather maps directly from this
 372 information. Several more series will be available in the near future and might further help towards
 373 that aim. However, there is also information in the time sequence of weather at each of these sites.
 374 Given the sparse information in space, it is essential to also exploit the information in the time
 375 sequence. Novel approaches such as deep learning algorithms might potentially be used for weather
 376 reconstruction but arguably would have to make use of time sequences in order to be successful. Such
 377 methods first need to be tested extensively in long data sets generated from recent products using tools
 378 such as synthetic weather diaries (Brönnimann, 2021).



379
 380 **Fig. 11.** Top: Standardised anomalies (relative to 1696-1704) of documentary data from Burgdorf et al. (2022)
 381 for temperature and precipitation. Bottom: Anomaly (left) and standardized anomaly (right, both relative to
 382 1696-1704) of temperature in EKF400v2. Note that EKF400v2 includes the four monthly series shown in the top
 383 row (three temperature indices and one precipitation index), but not the other series.

384 Finally, the diary, together with other sources of information, provides some insights into the climate
 385 processes in the 1690s, a period that was characterized by cold winters in Europe and also cold and
 386 rainy summers. The winters such as 1697/98 were likely characterized by frequent blocking and a
 387 meridionalisation of circulation. This is also seen in Eimmart's diary. The pressure difference between
 388 Paris and London (Cornes et al., 2012a,b) exhibits its lowest values (winter average) in the 1690s (the

389 winters 1695, 1694, 1692, and 1698 occupy ranks 1, 2, 4, and 11). The North Atlantic Oscillation
390 index calculated from EKF400v2 also shows low values in these years (see Brönnimann, 2022).

391 These winters fell into the so-called Late Maunder Minimum (Luterbacher et al., 2001), when sunspot
392 activity was very low. In fact, these cold winters have often been attributed to low solar activity. A
393 more meridional circulation with more frequent blocking events due to low solar activity would be in
394 line with statistical analyses of later data (Woollings et al., 2010). In addition to solar forcing, also
395 volcanic eruptions could have played a role. Eruptions occurred in 1693 (Hekla, Serua) and 1694
396 (Komagatake), and although winters following volcanic eruptions sometimes show a winter warming
397 in north-eastern Europe, this dynamical effect does not always appear and cold seasons following
398 volcanic eruptions may also be cold. The summer case (1695) shows the cooling expected following a
399 volcanic eruption (Raible et al., 2015); documentary data confirm this also on a hemispheric scale
400 (Burgdorf, 2022). Whether cold winters and cold summers in the 1690s are related remains to be
401 studied. There are several possible memory effects that might help to maintain the cooling from the
402 summer to the next winter and spring, including the oceans (Raible et al. 2015) and Eurasian snow
403 cover (Reichen et al., 2022). All factors together may have generated a decade of cold weather similar
404 to the early 19th century (Brönnimann et al., 2019b), when both summers and winters were cold
405 particularly over Eurasia. Historical weather diaries could help to further shed light on climate
406 mechanisms operating on a decadal scale related to volcanic eruptions and a solar minimum. However,
407 more weather and climate data are needed for this.

408 The first instrumental series in Nuremberg covers 1718-1730 (observer: Rost, 3-4x daily), another
409 series (observer: Doppelmayer, daily) covers 1732-1743 (see Brönnimann et al., 2019a). Both were
410 digitised and are included in the electronic supplement. Both also contain wind direction, such that a
411 1695-1743 record of sub-daily wind could be generated. However, homogeneity needs to be assessed.

412 **6. Conclusions**

413 This paper describes the digitization of the weather diary of Georg Christoph Eimmart. The diary
414 contains information on sky conditions (in symbolic form), precipitation, temperature, and wind in
415 Nuremberg, 1695-1704. It is relevant as the 1690s were a particularly cold decade in Europe. At the
416 same time, this approximately marks the period back to which daily reconstructions of weather might
417 be possible. The newly digitized diary might contribute towards this aim.

418 The diary structure changes during the first ca. two years, but afterwards (for wind throughout the
419 period), the diary is consistent. Comparisons with other series from Europe show that the diary
420 provides useful and usable information on the daily weather. For instance, the local daily wind
421 direction in Nuremberg agrees with the large-scale pressure gradient. The usefulness is further
422 demonstrated on behalf of several case studies, covering a storm passage in October/November 1702,
423 the cold winter of 1697/8, and the cold summer of 1695. These cases also show that the spatial
424 information at the daily level is inevitably sparse in the late 17th century. Any approach to reconstruct
425 the daily weather during this time arguably needs to make use of the temporal as well as the spatial
426 information contained in all available data sources.

427 **Funding Information:** The work was funded by the Swiss National Science Foundation project WeaR (188701)
428 and the European Commission through H2020 (ERC Grant PALAEO-RA 787574).

429 **Acknowledgements.** I would like to thank Andrey Martynov for his help in obtaining the diary, Juhyeong Han
430 for her help in digitizing it, and Yuri Brugnara for coordinating the digitising work. Hans Gaab and Klaus-Dieter
431 Herbst pointed me to the diary. I also thank Heinz Wanner for the hand-drawn reconstructions and Rolando

432 Garcia-Herrera for providing the wind data. The simulations underlying EKF400v2 were performed at the Swiss
433 Supercomputer Centre (CSCS).

434 **Supplement:** Supplementary files include the diary in xls, the data files for Jena, Kaliningrad, Leiden, and
435 Nuremberg (instrumental data) in the SEF format and xls, an ASCII file with the documentary data (Fig. 10), R-
436 Code and a readme file.

437 **Code Availability:** R-code to produce Figures 9 and 11 is given in the Supplement.

438 **Data availability statement:** The diary is in the Supplementary Material, EKF400v2 is available from
439 DKRZ/WDCC (doi:10.26050/WDCC/EKF400_v2.0), the Fries diary is available from EURO-CLIMHIST
440 (www.euroclimhist.unibe.ch), pressure data from Paris and London are available from the Climatic Research
441 Unit (<https://crudata.uea.ac.uk/cru/data/parislondon/>), cloud cover and direction as well as temperature from
442 Paris is available from Climate of the Past (Pliemon et al., 2022). Documentary data are available from
443 <https://boris-portal.unibe.ch/handle/20.500.12422/207> (Burgdorf et al., 2022).

444 **Conflict of interest statement:** The author declares no conflict of interests.

445 **Author contributions:** SB digitized the diary together with a student, performed all analyses and wrote the
446 paper.

447 **References**

448 Anonymous: Curiöser Wetter-Calender/ Oder Beschreibung des Gewitters/ Wie solches im Jahr 1702. in der Königl. Preuß.
449 Residenz-Stadt Königsberg/ von Tag zu Tage/ durchs ganze Jahr observiret und aufgezeichnet worden. Kaliningrad, 61
450 pp. <http://resolver.staatsbibliothek-berlin.de/SBB0000599300000000>, 1703.

451 Barriopedro, D., Gallego, D., Álvarez-Castro, M. C., García-Herrera, R., Wheeler, D., Peña-Ortiz, C., and Barbosa, S. M.:
452 Witnessing North Atlantic westerlies variability from ships' logbooks (1685-2008). *Clim. Dyn.*, 43, 939-955, 2014.

453 Brázdil, R., Kiss, A., Luterbacher, J., and Valášek, H.: Weather patterns in eastern Slovakia 1717–1730, based on records
454 from the Breslau meteorological network. *Int. J. Climatol.*, 28, 1639-1651, 2008.

455 Brohan, P.: Machine Learning for Data Assimilation. http://brohan.org/Proxy_20CR/,_2022.

456 Brönnimann, S.: Synthetic weather diaries: concept and application to Swiss weather in 1816. *Clim Past.*, 16: 1937-1952. doi:
457 10.5194/cp-16-1937-2020, 2020.

458 Brönnimann, S.: From climate to weather reconstructions. *PLOS Climate* 1, e0000034, 2022.

459 Brönnimann, S., Allan, R., Ashcroft, L., Baer, S., Barriendos, M., Brázdil, R., Brugnara, Y., Brunet, M., Brunetti, M.,
460 Chimani, B., Cornes, R., Domínguez-Castro, F., Filipiak, J., Founda, D., García Herrera, R., Gergis, J., Grab, S., Hannak,
461 L., Huhtamaa, H., Jacobsen, K. S., Jones, P., Jourdain, S., Kiss, A., Lin, K. E., Lorrey, A., Lundstad, E., Luterbacher, J.,
462 Mauelshagen, F., Maugeri, M., Maughan, N., Moberg, A., Neukom, R., Nicholson, S., Noone, S., Nordli, Ø., Ólafsdóttir,
463 K. B., Pearce, P. R., Pfister, L., Pribyl, K., Przybylak, R., Pudmenzky, C., Rasol, D., Reichenbach, D., Řezníčková, L.,
464 Rodrigo, F. S., Rohde, R., Rohr, C., Skrynyk, O., Slonosky, V., Thorne, P., Valente, M. A., Vaquero, J. M., Westcott, N.
465 E., Williamson, F., and Wyszyński, P.: Unlocking pre-1850 instrumental meteorological records: A global inventory, *B.*
466 *Am. Meteorol. Soc.*, 100, ES389–ES413, 2019a.

467 Brönnimann, S., Franke, J., Nussbaumer, S. U., Zumbühl, H. J., Steiner, D., Trachsel, M., Hegerl, G. C., Schurer, A., Worni,
468 M., Malik, A., Flückiger, J., and Raible, C. C.: Last phase of the Little Ice Age forced by volcanic eruptions. *Nature*
469 *Geosci.*, 12, 650–656, 2019b.

470 Bünti, J. L.: Chronik des Johann Laurentz Bünti, Landammann, 1661-1736. Beiträge zur Geschichte Nidwaldens 34, Stans,
471 Historischer Verein Nidwalden, 418 pp., 1973.

472 Burgdorf, A.-M.: A global inventory of quantitative documentary evidence related to climate since the 15th century. *Clim.*
473 *Past*, 18, 1407–1428, <https://doi.org/10.5194/cp-18-1407-2022>, 2022.

474 Burgdorf, A.-M., Brönnimann, S., Adamson, G., Amano, T., Aono, Y., Barriopedro, D., Bullon, T., Camenisch, C., Camuffo,
475 D., Daux, V., del Rosario Prieto, M., Dobrovolny, P., Gallego, D., Garcia-Herrera, R., Gergis, J., Grab, S., Hannaford,
476 M. J., Holopainen, J., Kelso, C., Kern, Z., Kiss, A., Lin Kuan-Hui, E., Loader, N. J., Mozny, M., Nash, D., Nicholson, S.
477 E., Pfister, C., Rodrigo, F. S., Rutishauser, T., Sharma, S., Takacs, K., Vargas, E. T., Vega, I. DOCU-CLIM: A global
478 documentary climate dataset for climate reconstructions. *Scientific Data* (submitted), data link: [https://boris-](https://boris-portal.unibe.ch/handle/20.500.12422/207)
479 [portal.unibe.ch/handle/20.500.12422/207](https://boris-portal.unibe.ch/handle/20.500.12422/207)

480 Cornes R. C., Jones, P. D., Briffa, K. R., and Osborn, T. J.: A daily series of mean sea-level pressure for London, 1692-2007.
481 *Int. J. Climatol.*, 32, 641–656, 2012a.

482 Cornes R. C., Jones, P. D., Briffa, K. R., and Osborn, T. J.: A daily series of mean sea-level pressure for Paris, 1670–2007.
483 *Int. J. Climatol.* 32(8): 1135-1150. doi: 10.1002/joc.2349, 2012b.

- 484 Filipiak, J., Przybylak, R., and Oliński, P.: The longest one-man weather chronicle (1721–1786) by Gottfried Reyger for
485 Gdańsk, Poland as a source for improved understanding of past climate variability. *Int. J. Climatol.*, 39, 828–842. doi:
486 10.1002/joc.5845, 2019.
- 487 Gaab, H.: Zur Geschichte der Eimmart-Sternwarte. „Nürnberg [...] der beste ort pro studio Astronomiae in gantz
488 Deutschland“. *Regiomontanusbote Spezial*, Vol. 18, Nuremberg, 65 pp., 2005.
- 489 Gaab, H.: Die Eimmart-Sternwarte. [https://www.astronomie-nuernberg.de/index.php?category=eimmart&page=eimmart-](https://www.astronomie-nuernberg.de/index.php?category=eimmart&page=eimmart-sternwarte)
490 [sternwarte](https://www.astronomie-nuernberg.de/index.php?category=eimmart&page=eimmart-sternwarte), last visited 15-Dec 2022, 2022.
- 491 García-Herrera, R., Gimeno, L., Ribera, P., Hernández, E., González, E., and Fernández, G.: Identification of Caribbean
492 basin hurricanes from Spanish documentary sources. *Clim. Change*, 83, 55–85, 2007.
- 493 Herbst, K.-D.: Gottfried Kirch (1639–1710) - Astronom und Kalendermacher. <https://gottfried-kirch-edition.de/>, last visited
494 15-Dec 2022, 2022.
- 495 Imfeld, N., Pfister, L., Brugnara, Y., and Brönnimann, S.: 250 years of daily weather: Temperature and precipitation fields
496 for Switzerland since 1763, *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2022-1140>, 2022.
- 497 Jones, P.D., Osborn, T.J., Harpham, C. and Briffa, K.R.: The development of Lamb weather types: from subjective analysis
498 of weather charts to objective approaches using reanalyses. *Weather*, 69, 128–132, 2014 <https://doi.org/10.1002/wea.2255>
- 499 Kington, J.: The severe winter of 1697/8. *Weather*, 54,43–49, 1999.
- 500 Kington, J. A.: *The Weather of the 1780s Over Europe*. Cambridge: Cambridge University Press, doi:
501 10.1017/CBO9780511735721, 1988.
- 502 Lamb, H. H.: British Isles daily wind and weather patterns 1588, 1781–86, 1972–91 and shorter early sequences (in 1532,
503 1570 and other years, notably 1688, 1689, 1694, 1697, 1703, 1717, 1783–4, 1791, 1792, 1795, 1822, 1825, 1829, 1845,
504 1846, 1849, 1850, 1854–5. *Climate Monitor*, 20, 47–70, 1991.
- 505 Luterbacher, J., Rickli, R., Xoplaki, E., Tinguely, C., Beck, C., Pfister, C., and Wanner, H.: The Late Maunder Minimum
506 (1675–1715) – A Key Period for Studying Decadal Scale Climatic Change in Europe. *Climatic Change*, 49 441–462. doi:
507 10.1023/A:1010667524422, 2001.
- 508 Lundstad, E., Brugnara, Y., Pappert, D., Kopp, J., Hürzeler, A., Andersson, A., Chimani, B., Cornes, R., Demarée, G.,
509 Filipiak, J., Gates, L., Ives, G. L., Jones, J. M., Jourdain, S., Kiss, A., Nicholson, S. E., Przybylak, R., Jones, P. D.,
510 Rousseau, D., Tinz, B., Rodrigo, F. S., Grab, S., Domínguez-Castro, F., Slonosky, V., Cooper, J., Brunet, N. and
511 Brönnimann, S. (2022) Global historical climate database - HCLIM. *Scientific Data* (accepted), 2022.
- 512 Manley G.: 1684: The coldest winter in the English instrumental record. *Weather*, 66, 382–388, 1975.
- 513 Mellado-Cano, J., Barriopedro, D., García-Herrera, R., and Trigo, R.: New observational insights into the atmospheric
514 circulation over the Euro-Atlantic sector since 1685. *Clim Dyn* 54, 823–841, 2020. [https://doi.org/10.1007/s00382-019-](https://doi.org/10.1007/s00382-019-05029-z)
515 [05029-z](https://doi.org/10.1007/s00382-019-05029-z)
- 516 Nash, D. J., Adamson, G. C. D., Ashcroft, L., Bauch, M., Camenisch, C., Degroot, D., Gergis, J., Jusopović, A., Labbé, T.,
517 Lin, K.-H. E., Nicholson, S. D., Pei, Q., del Rosario Prieto, M., Rack, U., Rojas, F., and White, S.: Climate indices in
518 historical climate reconstructions: a global state of the art. *Clim. Past*, 17, 1273–1314, [https://doi.org/10.5194/cp-17-](https://doi.org/10.5194/cp-17-1273-2021)
519 [1273-2021](https://doi.org/10.5194/cp-17-1273-2021), 2021.
- 520 Pappert, D., Barriendos, M., Brugnara, Y., Imfeld, N., Jourdain, S., Przybylak, R., Rohr, C. and Brönnimann, S.: Statistical
521 reconstruction of daily temperature and sea-level pressure in Europe for the severe winter 1788/9. *Clim. Past*, 18, 2545–
522 2565, <https://doi.org/10.5194/cp-18-2545-2022>, 2022.
- 523 Pfister C.: *Wetternachhersage: 500 Jahre Klimavariationen und Naturkatastrophen (1496–1995)*. Bern: Haupt Verlag, 1999.
- 524 Pfister C., and Wanner H.: *Climate and Society in Europe*. Bern: Haupt Verlag, 2021.
- 525 Pfister, C.: Zum Klima des Raumes Zürich im späten 17. und frühen 18. Jahrhundert. *Vierteljahrsschr. Naturf. Ges. Zürich*
526 122, 447–471, 1977.
- 527 Pfister, C., Rohr, C., and Jover, A. C. C.: Euro-Climhist: eine Datenplattform der Universität Bern zur Witterungs-, Klima-
528 und Katastrophengeschichte. *Wasser Energie Luft*, 109, 45–48, 2017.
- 529 Pliemon, T., Foelsche, U., Rohr, C., and Pfister, C.: Subdaily meteorological measurements of temperature, direction of the
530 movement of the clouds, and cloud cover in the Late Maunder Minimum by Louis Morin in Paris, *Clim. Past*, 18, 1685–
531 1707, <https://doi.org/10.5194/cp-18-1685-2022>, 2022.
- 532 Przybylak, R., and Pospieszńska, A.: Air temperature in Wrocław (Breslau) in the period 1710–1721 based on
533 measurements made by David von Grebner. *Acta Agrophysica*, 184, 35–43, 2010.
- 534 Raible, C. C., Brönnimann, S., Auchmann, R., Brohan, P., Frölicher, T. L., Graf, H.-F., Jones, P., Luterbacher, J., Muthers,
535 S., Neukom, R., Robock, A., Self, S., Sudrajat, A., Timmreck, C., and Wegmann, M.: Tambora 1815 as a test case for
536 high impact volcanic eruptions: Earth system effects. *WIREs Climate Change*, 7, 569–589, doi: 10.1002/wcc.407, 2016.
- 537 Reichen, L., Burgdorf, A.-M., Brönnimann, S., Rutishauser, M., Franke, J., Valler, V., Samakinwa, E., Hand, R., and
538 Brugnara, Y.: A Decade of Cold Eurasian Winters Reconstructed for the Early 19th Century. *Nature Communications* 13,
539 2116, <https://doi.org/10.1038/s41467-022-29677-8>, 2022.

- 540 Rohr, C., and Schwarz-Zanetti, G.: Das Kloster-Tagebuch des Einsiedler Paters Joseph Dietrich, 1670-1704. 1670-1704
541 http://www.dietrich-edition.unibe.ch/Ueber_das_Projekt.html, last visited 15-Dec-2022, 2022.
- 542 Rousseau, D.: Les températures mensuelles en région parisienne de 1676 à 2008, *La météorologie*, 67, 43–55, 2009.
- 543 Schwander, M., Brönnimann, S., Delaygue, G., Rohrer, M., Auchmann, R., and Brugnara, Y.: Reconstruction of Central
544 European daily weather types back to 1763. *Int. J. Climatol.* 37, 30-44. doi: 10.1002/joc.4974, 2017.
- 545 Valler, V., Franke, J., Brugnara, Y., and Brönnimann, S.: An updated global atmospheric paleo-reanalysis covering the last
546 400 years. *Geosc. Data J.*, 9, 89– 107, doi: 10.1002/gdj3.121, 2022.
- 547 Wanner, H., Pfister, C., Brázdil, R., Frich, P., Frydendahl, K., Jónsson, T., Kington, J., Lamb, H. H., Rosenørn, S., and
548 Wishman, E.: Wintertime European circulation patterns during the Late Maunder Minimum cooling period (1675–1704).
549 *Theor. Appl. Climatol.*, 51, 167-175. doi: 10.1007/BF00867443, 1995.
- 550 Wheeler, D., Garcia-Herrera, R., Vaquero, J. M., Chenoweth, M., and Mock, C. J.: Reconstructing The Trajectory of The
551 August 1680 Hurricane From Contemporary Records. *B. Amer. Meteorol. Soc.*, 90, 971-978. doi:
552 10.1175/2009BAMS2649.1, 2009.
- 553 Wheeler, D., Garcia-Herrera, R., Wilkinson, C. W., and Ward, C.: Atmospheric circulation and storminess derived from
554 Royal Navy logbooks: 1685 to 1750. *Climatic Change*, 101, 257-280. doi: 10.1007/s10584-009-9732-x, 2010.
- 555 Woollings, T., Lockwood, M., Masato, G., Bell, C., and Gray, L.: Enhanced signature of solar variability in Eurasian winter
556 climate. *Geophys. Res. Lett.*, 37, L20805, doi:[10.1029/2010GL044601](https://doi.org/10.1029/2010GL044601), 2010.