1 **The weather diaries of the Kirch family: Leipzig, Guben, and Berlin, 1677-**

2 **1774**

4 Stefan Brönnimann*, Yuri Brugnara

5 *Oeschger Centre for Climate Change Research and institute of Geography, University of Bern*

6 *corresponding author: Stefan Brönnimann, stefan.broennimann@giub.unibe.ch

7 **Abstract**

8 Astronomer and calendar maker Gottfried Kirch was a keen weather observer and made weather notes in his

9 diary starting in 1677 in Leipzig. In parallel, his second wife Maria Margaretha Winkelmann started a weather

10 diary in 1700 in Berlin. The diaries also contain instrumental measurements of temperature and later pressure.

11 After the death of Gottfried in 1710 and Maria Margaretha in 1720, observations were continued by their son

12 Christfried and then for another 44 years by their daughter Christine. The last measurements date to 1774.

13 Together, the diaries span almost a century of weather observations. The instrumental measurements constitute

14 the oldest part of Germany's longest temperature series, which was however only available as monthly means up

15 to now. Here we publish the imaged diaries, together amounting to 10445 images. Further, we present the

16 digitised instrumental series, which will serve as the starting point for a new, daily Berlin series. By comparing

17 the series to neighbouring records, we show that the pressure data are reliable in a quantitative sense, whereas

18 this is true for the temperature data only in a qualitative sense as the temperature scale was not converted.

19

20 **1. Introduction**

21 Long historical climate records are invaluable to better understand variations in climate and the

22 underlying mechanisms. For a long time, the emphasis was on monthly or seasonal averages, and long

23 meteorological series were often only available in that form. Recently, changes in weather patterns and

24 extremes came into focus. New tools such as reanalyses (e.g., Slivinski et al., 2019) or analog

25 approaches (Pappert et al., 2022) now allow the daily weather to be reconstructed, from which

26 important conclusions can be drawn about decadal to multidecadal variations in weather as well as

27 extreme weather (Brönnimann, 2022). However, many of the long series are not available at daily or

28 sub-daily resolution (i.e., the individual measurements), but only as monthly means. It is therefore

29 often required to revisit archives, image and digitise the sub-daily data and start the homogenisation

30 processes anew.

31 In this paper, we present the work of imaging, digitising, and processing for the case of the longest

32 German record, that from Berlin which goes back to 1701. Specifically we analyse the weather diary

33 of the Kirch family, covering 1677-1774, which contains some instrumental observations in 1697 and

34 then from 1701 onward. This record has been widely used since the late 18th century. Karl-Ludwig

35 Gronau compiled the measurements and calculated monthly means (Gronau, 1807), supplementing the

36 sometimes sparse measurements in an unknown way. In the 19th century, Johann Heinrich Mädler

37 continued working on a Berlin series and presented a new, extended series (Mädler, 1825). Hellmann

38 (1893) re-discovered the weather diaries of Maria Margaretha Kirch, Lenke (1964) used the Berlin

39 data in his study on the cold winter 1708/9 and Pelz (1978) examined the Kirch diaries. However,

40 most other authors did not consult the original diaries. In the German Democratic Republic, Bahr

41 (1966) worked on the history of the Berlin series in the context of her dissertation. Subsequent work

42 led to the publication of a daily temperature series back to 1766, which was recently digitized by

43 Kadow et al. (2016). In Western Germany, Pelz (1997) re-homogenised and published the monthly

44 Berlin series (Cubasch and Kadow, 2010). What is still missing is the daily or sub-daily temperature

45 series earlier than 1766. Furthermore, although pressure was also measured, it was never digitised or

46 analysed. Therefore, we revisited the original sources, imaged the diary and digitised most of the

47 instrumental measurements (both temperature and pressure) contained in the sheets.

48 The paper is organised as follows. Section 2 provides background about the Kirch family and their

49 meteorological observations. Sect. 3 describes the diary and its history. In Sect. 4 we describe the

50 digitising, Quality Control, and comparison with other sources. Results are presented in Section 5 and

51 conclusions are drawn in Sect. 6.

52 **2. The Kirch family**

53 *2.1. Life and work*

54 Gottfried Kirch was one of the leading astronomers of the late 17th century. A recent biography

55 (Herbst, 2022) gives a detailed account of his life and work, which is only briefly summarized in the

56 following. Kirch was born in Guben (Fig. 1). In the 1660s he started to publish calendars, which

57 remained an important source of income for the family even after his death. As an astronomer, Kirch

58 became famous in the early 1680s when he discovered a comet and the star cluster M11. His second

59 wife Maria Margaretha Winkelmann also was an astronomer (but she was not admitted to the

60 university of Halle, to which she applied). She discovered the comet C/1702 H1 and worked on

61 sunspots. Gottfried and Maria Margaretha Kirch had six children, many of which supported or

62 continued the astronomical and meteorological work. Two of them, Christfried and Christine, had their

63 own weather diaries. Christine was further supported by her sisters Margaretha and Dorothea. Figure 2

64 show the observers and observation locations on a time axis.

65 The Kirch family lived in Leipzig from 1676 onward, then moved to Guben in 1692 (Fig. 1). In 1700

66 Gottfried Kirch was appointed as Royal astronomer by the newly funded Prussian Academy of

67 Science in Berlin (with Gottfried Wilhelm Leibniz as president). However, the astronomical

68 observatory was not built yet, and for the next 10 years observations were performed in the family's

69 apartment (Fig. 1).

70 Also the promised apartment was not ready in 1700, and therefore the family initially stayed at

71 different places. In 1708 they eventually moved into the „Astronomenhaus" at Dorotheenstrasse 10

72 (Fig. 1). The observatory was just next door (Fig. 23), but was officially opened only in 1711,

73 although observations were made earlier.

74 Gottfried Kirch died in 1710. After the death of her husband, Maria Margaretha Kirch and her son

75 Christfried Kirch continued the observations. They decided that Christfried should note the

76 astronomical observations and Maria Margaretha Kirch the weather observations. However, the

77 financial situation became more and more precarious for Maria Margaretha Kirch. Despite her

78 qualification, she could not follow as a director of the observatory but continued to publish the

79 calendars. In 1712, she moved into the house of Baron von Krosigk at Wallstrasse 135 (Fig. 1), where

80 she observed for the next two years (however, there was no thermometer). In 1714, she moved to

81 Gdansk with the children. In 1716, Christfried Kirch became the director of the Berlin observatory,

82 and Maria Margaretha Kirch moved back to her son into the „Astronomenhaus". In 1716 she began to

83 measure again and continued almost until her death on 29 December 1720. The data presented in this

84 article start in 1721 and were presumably all taken at the "Astronomenhaus" (location B in Fig. 1).

85 Measurements were continued together by Christfried und Christine Kirch. Christfried died in 1740.

86 Although not officially a member of the Prussian Academy, Christine continued the astronomical

87 observations and was paid by the Academy. She also performed meteorological measurements and

88 continued publishing the calendars. Her house was a gathering point for scientists of the 18th century.

89 Leonhard Euler was a frequent visitor, other guests include Joseph-Nicolas Delisle and Anders

90 Celsius. The last instrumental meteorological measurements date to 30 April 1774. Christine Kirch

91 died in 1782.

92

93 **Fig. 1.** Left: Map of Berlin in 1712. A: Astronomical observatory. B: Apartment of the Kirch family until 1712

94 and from 1716 onward ("Astronomenhaus"). C: House of Baron von Krosigk, where Maria Margaretha observed

95 from 1712-1714 (from map "Grundlicher Abriß, der königl. Haupt- und Residenz Stadt Berlin", unknown

96 author, public domain, wikimedia commons). Right: Locations of other weather observations in the Kirch diaries

97 (Berlin and empty circles) and other locations, coloured according to the Pearson correlation of their pressure

records with the Kirch measurements, Berlin, 1730-1770, in the overlapping period (circle area indicates tne number of measurements *n*). For Berlin, the circle shows the correlation with the reconstruction EKF400v2.

Fig. 2. Temporal overview of observers and observation locations.

Fig. 32. The old Berlin astronomical observatory (excerpt from an etching published in the Atlas Coelestis of Johann Gabriel Doppelmayr, 1742, public domain, wikipedia).

2.2. Instruments

Gottfried and Maria Margaretha Kirch had a thermometer ("Wetterglas") since 1691. It was a Florentin-type thermometer manufactured by Gottfried Kirch: a glass bulb filled with spirit of wine with a closed tube. However, the thermometer seems to have been broken at some point. The notes clearly speak of an old and a new thermometer, although it is not fully clear when the change took place.

At that time observers made their own temperature scales as there was no agreed-upon scale. According to the description by Kirch, the thermometer had a 60 degree scale. The freezing point was at 20 degrees, and 40 degrees corresponded to a hot summer day. The temperature scale was later analysed by Lenke (1964), who converted the Berlin temperature data for the winter 1708/9 to a Celsius scale, though not without difficulties. Later, Christfried und Christine Kirch reportedly used a Fahrenheit thermometer, but the given temperatures are inconsistent with a Fahrenheit scale. Pelz (1978) mentions six different scales that were used in the Kirch diaries, of which only the first one is approximately known.

A barometer was in use since 1709. However, no details are known about the instrument. Likewise, not much is known about the siting of the instruments. According to Lenke (1964), measurements in the "Astronomenhaus" were made in a north facing window in the middle floor.

3. The weather diary

An overview of the weather diaries available for this study is given in Table 1. Gottfried Kirch's diary starts in 1677 when he worked in Leipzig. It contains mostly astronomical observations, but also sporadic weather information that were important for his astronomical observations. Noteworthy is Maria Margaeretha's diary, a specific weather diary which starts in August 1700 (plus January to June 1697, containing instrumental measurements made in Guben). The first instrumental measurements in Berlin date to 18 January 1701 (Fig. 34), the day of the coronation of the Prussian king Friedrich I. For several years, the diaries of Gottfried and Maria Margaeretha run parallel. The motivation behind the instrumental measurements most likely was checking the calendar (Herbst, 2022).

Table 1. Boxes with printouts of the Kirch diaries at Free University of Berlin, content of the boxes, original archive, number and availability of microfilms. Note that this Table also corresponds with the folder structure on the repository. More information on individual years is given in Pelz (1978).

Author	Content	Original archive	Film number	Microfilms
Gottfried Kirch	astronomical diary 1677-1685	Paris Observatory	Film 649I	not found
Gottfried Kirch	astronomical diary 1685-1689	Paris Observatory	Film 649II	available
Gottfried Kirch	astronomical diary 1696-1704	Paris Observatory	Film 650I	available
Gottfried Kirch	astronomical diary 1704-1708	Paris Observatory	Film 650II	available
Gottfried Kirch	astronomical diary 1708-1710	Paris Observatory	Film 650III	available
Maria Margaretha Kirch	weather diary 1697, 1700-1718	Royal Observatory	Rolle 1/Film 583 Rolle 2/Film 584	not found
Maria Margaretha Kirch and Christfried Kirch	weather diary 1718-1720, 1721 and 1728	Royal Observatory	Rolle 3/Film 585	not found
Christine Kirch	weather diary 1730-1734	Royal Observatory	Rolle4/Film586	available
Christine Kirch	weather diary 1734-1737	Royal Observatory	Rolle5/Film587	available
Christine Kirch	weather diary 1738-1743	Royal Observatory	Rolle6/Film588	available
Christine Kirch	weather diary 1743-1747	Royal Observatory	Rolle7/Film589	available
Christine Kirch	weather diary 1748-1756	Royal Observatory	Rolle8/Film590	available
Christine Kirch	weather diary 1757-1761	Royal Observatory	Rolle9	available
Christine Kirch	weather diary 1762-1770 and 1774	Royal Observatory	Rolle 10/Film 592 Rolle 11/Film 593	available

Both the diaries of Gottfried and Maria Margaretha end only shortly before their deaths. After 1720, weather data were contained in Christfried's diary, but we only have data for selected years. From 1730 on the weather observations are noted in Christine's diary. Note that an attribution is difficult to make since both signed their observations with "CK". Most of the observations in our digitised record are from Christine, who was assisted by her sisters. One of the last pages of her diary, from 1770, is displayed in Fig. 4-5 (there are no entries at all for 1771-1773, and in 1774 only from January to April).

142 The diary of Gottfried Kirch was already famous in the 18th century. Several copies must exist.

143 According to Lenke (1964), Joseph-Nicolas Delisle bought Gottfried Kirch's diary from Christine

144 Kirch and gave it to the "Dépôt général de la Marine". After the French Revolution, the diaries ended

145 up at the Observatoire de Paris where they remain to the present day. The diaries of Maria Margaretha,

146 Christfried and Christine Kirch are stored today at the Royal Observatory in Edinburgh. How they

147 ended up there is not known. It seems that for a long time, the location of these diaries was unknown.

148 Hellmann (1893) reports how he searched for them and how they eventually were found in Edinburgh.

149 He then published a transcription of the first years of the diary of Maria Margaretha (Hellmann, 1893).

150

151

152 **Fig. 34.** First temperature measurement from Berlin on 18 January 1701, from the weather diary of Maria

153 Margaretha Kirch (The Crawford Collection of the Royal Observatory Edinburgh, top) and Gottfried Kirch

154 (Paris Observatory, bottom). Both mention that the thermometer was at 16.5 degrees. The last lines of the top

155 excerpt read: "Das Wetterglas hing in einer ziemlich offenen Kammer und war 16 und einhalb Grad." (The

156 weather glass was in an open chamber and was 16 and one half degrees).

157 At the request of the German Weather Service, the Royal Observatory in Edinburgh filmed the Kirch

158 diaries in 1959, and in 1962 the Paris Observatory filmed diaries of Gottfried Kirch. In 1977, 16 roles

159 of film were duplicated and Xerox print-outs were made. At this occasion, Pelz (1978) analysed the

160 diaries. The print-outs and films are still stored today at the library of the Institute of Meteorology of

161 the Free University of Berlin (Fig. 56). Of the 16 roles of film, only 12 could be found (missing is film

162 649I of Gottfried Kirch's diary and roles No. 1, 2, 3, of Maria Margaretha Kirch's diary, see Table 1).

163 Print-outs of all films are available.

164 The print-outs were imaged on 6-12 Mar 2020 by the first author. The paper quality did not allow an
165 automatic scanning. Hence the paper sheets were photographed with a handheld smartphone. The
166 impending lockdown due to the COVID pandemic forced us to work quickly. It was decided not to
167 image the part from 1677-1700 since it did not contain instrumental observations, and the ca. 7500
168 pages from 1700 to 1774 were photographed rather quickly. The imaged data sheets could then be
169 transcribed during the lockdown and following period. In summer 2022, the first author returned to
170 Berlin to also photograph the remaining portion (1677-1700). No data were transcribed form from
171 these images, they were merely made to have the diaries imaged in their entirety and to keep the
172 diaries together electronically at one location.

174 **Fig. 45.** Temperature and pressure measurements in 1770 by Christine Kirch (The Crawford Collection of the
175 Royal Observatory Edinburgh).

Fig. 5~~6~~. Left: The boxes with Xerox print-outs photographed at Free University Berlin. Right: 12 Boxes with microfilms from which the print-outs were made.

4. Methods

Although first observations already start in the late 17th century, they were often not presented in a tabular form. Measurements were at times quite frequent, but they were often interspersed in the text, such as in Fig. 3~~4~~. Sometimes measurements were organised in tables, but not consistently. We digitised the data only from 1720 onward, when they are presented in a consistently tabular format. The published images accompanying this article allow digitising also the earlier measurements.

We digitised temperature and pressure measurements. The actual keying work was performed by students of the University of Bern, who received a template and instructions (see spreadsheets in Supplementary Data: Tab "Info" contains the instructions).

The pressure data were then processed as described in Brugnara et al. (2020) and converted to the station exchange format (SEF, Brunet et al., 2020), except that pressure data could not be corrected to 0°C as no temperature data are available. Temperature was digitised, but not converted, as the scale is unknown. The quality control (QC) package datarescqc (Brugnara et al., 2019) was used (see also descriptions in Brugnara et al., 2020; Brunet et al., 2020). When a quality issue was found, this was noted in the column "Meta" of the SEF file (the measurement value itself was not changed). As an example, Fig. 7 shows the digitised tabular data as well as the SEF file for the case of July 1721. The header of the SEF file provides metadata at the station level, including whether the pressure data were temperature corrected (PTC = N) or gravity corrected (PGC = Y) and the QC software used. Metadata on the level of individual measurements is indicated in the column meta. In this case, two values are flagged. The information "qc=wmo_time_consistency" means that the pressure change to the next measurement is larger than the threshold recommended by WMO (1993). In this case, the cause the low value at 11 pm local time (22:06 UTC), after which pressure jumps back to the previous value.

201 Both jumps are flagged. The conversion from local solar time to UTC does not take into account the
202 equation of time, therefore the difference with UTC is always 54 minutes.

203

205 **Fig. 7.** Example of the digitised data in the spread-sheet and the corresponding data in the SEF format.

206

207 ~~and~~The data were submitted to the GLAMOD data base of Copernicus Climate Change Service

208 (Noone et al., 2021). All data, both temperature and pressure, are also published as ~~Excel files~~spread

209 sheets as a supplement to this article.

210 For assessing the Berlin series, we used pressure series from the HCLIM database (Lundstad et al.,

211 2022). Specifically, we selected monthly data for pressure for all stations in Europe with at least 30

212 months of overlap (the shortest overlap is 45 months). In addition, daily pressure data from

213 Nuremberg were used. Further, we used Gdansk temperature (in the form monthly minima and

214 monthly maxima). Finally, we used sea-level pressure from the reconstruction EKF400v2 (Valler et

215 al., 2022) at the grid point closest to Berlin for comparison with the digitised measurements from

216 Berlin (note that Berlin pressure was not used for the reconstruction and hence is independent).

217 **5. Results**

5.1. Digitising and processing

We digitised 42065 pressure measurements and 39639 temperature measurements. Of the former, 49 have a flag "wmo_time_consistency" and two have a flag "duplicate_observation_time". An overview of the temporal coverage of pressure and temperature data is given in Fig. 68. The data cover the periods 1730 to 1751 and 1756 to 1770 very well, with typically 3 to 4 measurements per day. The 1720s and the period 1752-5 are less well covered. For the year 1726 we only know the monthly mean pressure values. No data are available for the years 1771-1773 and only few pressure measurements for 1774.

Fig. 68. Number of individual measurements digitised.

The time of day of the measurement is always indicated. A histogram of the measurement hours (Fig. 9, top) shows clear peaks, namely between 7 and 9 UTC, between 11 and 12 UTC, and 20 and 22 UTC. A smaller but distinct peak occurs between 14 and 15 UTC. Thus, the measurements were taken rather regularly at the usual observing times (local time is approximately 1 hour ahead of UTC).

A histogram of all pressure measurements is given in Fig 9 (bottom). An open question concerns the scale. Pelz (1978) assumes that pressure was not given in the official local scale, Rhineland inches (or Prussian inches, which is the same), as the numbers would otherwise be too low, but in Paris inches. However, assuming Paris inches would lead to numbers that are too high, the difference being 30 hPa. We assumed Rhineland inches, and consequently the average pressure is around 995 hPa, which is too low. We note that homogenization will be necessary to scale the pressure data. That said, the time series of monthly mean pressure (Fig. 10), does not show any obvious systematic change over time.

241

242
243

244

245

Fig. 10. Time series of monthly mean pressure.

246

247

5.2. Comparison with other sources

248

First we analysed the pressure data, which we compared with neighbouring series. Correlations on the scale of monthly mean values are shown in Fig. 1 (right). Note that only few series (Paris and London) have been homogenised. Most other series are analysed here in their original form. Nevertheless, ~~Looks great~~results show very high correlations exceeding 0.8 with Nuremberg and Zwanenburg. Also, the correlations at Berlin with EKF400v2 reaches almost 0.8 and the correlation with London pressure is in the same range. Somewhat lower, but still high correlations (>0.65) are found for Gdansk, Paris, and Basel. Lower correlations exclusively originate from comparisons with shorter records which are not homogenised and which have not yet been used much in the scientific literature, if at all. Overall

249
250
251
252
253
254
255
256

257 this clearly shows that the Berlin pressure data is of high relative quality and thus adds information to
258 the existing body of climate data.

259 *5.3. Case study: Particularly cold winter 1739/40*

260 One of the coldest European winters of the second millennium was 1739/40 (Schlaak, 1984,
261 Luterbacher and Wanner, 2002, Casty et al., 2005). It exhibits the lowest cold-season temperature of
262 the northern extratropical land areas of the last three centuries in a recent reconstruction by Reichen et
263 al. (2002). In fact, results from the spring phenology data from Europe used in that study are
264 summarized in Table 2. Clearly, the spring was extremely late in 1740, although it set the record only
265 in one of the series. In Figure 7 11 we present a very simple analysis of daily mean temperature in
266 Berlin from 1738 to 1743. For comparison we also show the monthly minimum and maximum
267 temperatures from Gdansk, which is over 400 km away. For Berlin, all measurements made on a
268 particular day were averaged without considering possible variations in the time of day of the
269 measurement. Note also that we do not know the scale of temperature, although it is reported that
270 Christfried and Christine Kirch used a Fahrenheit thermometer. The fact that slightly negative values
271 are reached in Jan. 1740 might indicate Fahrenheit temperature (where zero corresponds to -17.8 °C)
272 or it might indicate that the liquid dropped below a self-defined scale. The summer values are clearly
273 too low to be degrees Fahrenheit. Despite all these factors and despite the distance between the two
274 sites, we clearly see common variations. For instance, minimum temperatures were low in Nov. 1739,
275 but then high in in Dec. 1739 (with high maximum temperatures as well), then Jan. 1740 had very low
276 minimum temperatures but rather normal maximum temperature.

277

278 **Table 2.** Phenological spring data in Europe (from Reichen et al., 2022), rank of year 1740 and number of years
279 in the record.

Location	Proxy	rank	N
Mälaren	ice break-up	19	288
Tallinn	ice break-up	21	363
Tornio	ice break-up	4	311
St. Peterburg	ice break-up	6	173
Zurich	cherry blossom	1	283
Haarlem	days of freeze	5	147
Turku	ice break-up	2	79

280

281 For pressure, we can compare the Berlin series with that from Nuremberg (the observer was Johann
282 Gabriel Doppelmayr, who is also the author of the Atlas from which Fig. 2 3 is taken). The agreement
283 is very good, with a correlation coefficient of 0.83 despite some probable outliers. For the winter
284 1739/40, we find low pressure in Dec. 1739, but then high pressure in Jan. 1740. This is consistent
285 with temperatures.

286

287

Fig. ~~7~~11. Top: Daily averages of temperature from 1738-1742 by Christine Kirch (left axis; note that the temperature scale is not known). Also shown are the monthly minimum and maximum temperatures in Gdansk (right axis). Bottom: Daily mean pressure in Berlin (reduced by 30 hPa for comparison) and Nuremberg.

288
289
290
291

6. Discussion and conclusions

292

The Kirch family (Gottfried Kirch, his wife Maria Margaretha Kirch (née Winkelmann), their son Christfried Kirch and their daughter Christine Kirch) noted the weather for almost a century, from 1677 to 1774. From 1691 onward they were using a thermometer, later also a barometer, although regular observations only start in 1701. This body of measurements makes up the first part of the longest meteorological record in Germany. We imaged the diaries, totalling 10445 images, and make them available on a repository. Moreover, we digitised ca. 82000 instrumental observations after 1720 and publish them. Pressure data could be processed (although a reduction to 0 °C could not be performed). Temperature data could not be reduced because of an unknown (and changing) scale but were nevertheless digitised and are published as a supplement to this paper. Comparisons with other data suggest that the pressure series is trustworthy on the daily as well as monthly scale, although the scale remains uncertain. In fact, correlations with neighbouring series are very high. A brief analysis of the cold winter of 1739/40 suggests that also temperature measurements may contain useful information, even though the scale remains unknown. The newly digitised series will serve as the starting point for a new, daily Berlin series of temperature and pressure. The Kirch data set will be concatenated with other Berlin series from the 18th century currently under digitisation (Lambert, Jablonski, Gronau, Brand, and others) and homogenised to generate a more complete Berlin series.

293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308

The early Berlin data fall into a period in which not many other records are known and therefore they provide valuable information. However, there are some records with which the Berlin record can be compared. This includes long daily time series from Paris (Pliemon et al., 2022, Cornes et al., 2012a), London (Cornes et al., 2012b), Zwanenburg/DeBilt, shorter series from Nuremberg, Basel, Geneva, Zurich and Bern (e.g., Brugnara et al., 2022), St. Peterburg, Uppsala, and several Italian series from the IMPROVE project (Camuffo and Jones, 2002), among others (see inventory by Brönnimann et al., 2019). However, also other, non-instrumental weather diaries may be a good resource for comparisons, including those from Nuremberg (Brönnimann, 2023), Wroclaw (Przybylak, and Pospieszyńska, 2010), Gdansk (Filipiak et al., 2019) or Zurich (see weather diaries in EURO-CLIMHIST, Pfister et al., 2017). All sources taken together may provide a detailed view of weather in the 18th century.

Funding. The work was funded by the Swiss National Science Foundation project WeaR (188701) and the European Commission through H2020 (ERC Grant PALAEO-RA 787574).

Acknowledgements. We thank the FU Berlin for allowing us to photograph their material, the Paris Observatory, and the Crawford Collection of the Royal Observatory Edinburgh for allowing us to publish the data under a CC-BY-NC licence. We also thank Klaus-Dieter Herbst for his support and information on the Kirch family. In particular, we thank the students who performed the digitisation work.

Data availability

The images of the Kirch diaries can be found at: https://doi.org/10.48620/222

SEF data files for pressure in Berlin as well as Nuremberg have been submitted to the GLAMOD data base of Copernicus Climate Change service, they are also attached to this submission, together with the raw files for all variables for Berlin.

The monthly pressure data are part of the H-CLIM collection: https://doi.pangaea.de/10.1594/PANGAEA.940724

Author contributions

SB imaged the diaries, YB organised the digitisation, performed the quality control and all processing and formatting steps of the Berlin data. SB performed the analyses in the paper. Both authors wrote the paper.

Competing interests. The contact author has declared that none of the authors has any competing interests.

References

Bahr, R.-M.: Das Klima von Berlin (I). Geschichte der meteorologischen Beobachtungen der Stadt Berlin. Abh. Meteorol. Dienst DDR, 78, Bd. 10, 1966.

Brönnimann, S. Allan, R., Ashcroft, L., Baer, S., Barriendos, M., Brázdil, R., Brugnara, Y., Brunet, M., Brunetti, M., Chimani, B., Cornes, R., Domínguez-Castro, F., Filipiak, J., Founda, D., García Herrera, R., Gergis, J., Grab, S., Hannak, L., Huhtamaa, H., Jacobsen, K. S., Jones, P., Jourdain, S., Kiss, A., Lin, K. E., Lorrey, A., Lundstad, E., Luterbacher, J., Mauelshagen, F., Maugeri, M., Maughan, N., Moberg, A., Neukom, R., Nicholson, S., Noone, S., Nordli, Ø., Ólafsdóttir, K. B., Pearce, P. R, Pfister, L., Pribyl, K., Przybylak, R., Pudmenzky, C., Rasol, D., Reichenbach, D., Řezníčková, L., Rodrigo, F. S., Rohde, R., Rohr, C., Skrynyk, O., Slonosky, V., Thorne, P., Valente, M. A., Vaquero, J. M., Westcottt, N. E., Williamson, F., and Wyszyński, P.: Unlocking pre-1850 instrumental meteorological records: A global inventory, B. Am. Meteorol. Soc., 100, ES389–ES413, 2019.

347 Brönnimann, S.: From climate to weather reconstructions. PLOS Climate 1, e0000034, 2022.

348 Brönnimann, S.: The weather diary of Georg Christoph Eimmart for Nuremberg, 1695-1704. Clim. Past (submittedaccepted),
349   2023.

350 Brugnara, Y., Gilabert, A., Ventura, C., and Hunziker, S.: dataresqc: Quality control tools for climate data developed by the
351   C3S Data Rescue Service, available at: https://github.com/c3s-data-rescue-service/dataresqc, last access: 14 June 2023.

352 Brugnara, Y., Pfister, L., Villiger, L., Rohr, C., Isotta, F. A., and Brönnimann, S.: Early instrumental meteorological
353   observations in Switzerland: 1708–1873, Earth Syst. Sci. Data, 12, 1179–1190. 2020.

354 Brugnara, Y., Hari, C., Pfister, L., Valler, V., and Brönnimann, S.: Pre-industrial Temperature Variability on the Swiss
355   Plateau Derived from the Instrumental Daily Series of Bern and Zurich, Clim. Past 18, 2357–2379, 2022,
356   https://doi.org/10.5194/cp-18-2357-2022.

357 Brunet, M., Brugnara, Y., Noone, S., Stephens, A., Valente, A.A., Ventura, C., Jones, P., Gilabert, A., Brönnimann, S.,
358   Luterbacher, J., Allan, R., Brohan, P., and Compo, G.P.: Best Practice Guidelines for Climate Data and Metadata
359   Formatting, Quality Control and Submission. Copernicus Climate Change Services, 2020, https://doi.org/10.24381/kctk-
360   8j22

361 Camuffo, D., and Jones, P.: Improved understanding of past climatic variability from early daily European instrumental
362   sources. Climatic Change, 53, 1–4, 2002, https://doi.org/10.1023/A:1014902904197

363 Casty, C., Wanner, H., Luterbacher, J., Esper, J. and Böhm, R.: Temperature and precipitation variability in the European
364   Alps since 1500. Int. J. Climatol., 25: 1855-1880, 2005, https://doi.org/10.1002/joc.1216

365 Cornes R. C., Jones, P. D., Briffa, K. R., and Osborn, T. J.: A daily series of mean sea-level pressure for Paris, 1670–2007.
366   Int. J. Climatol. 32(8): 1135-1150. doi: 10.1002/joc.2349, 2012a.

367 Cornes R. C., Jones, P. D., Briffa, K. R., and Osborn, T. J.: A daily series of mean sea-level pressure for London, 1692-2007.
368   Int. J. Climatol., 32, 641–656, 2012b.

369 Cubasch, U., and Kadow, C.: Die Berliner Temperaturreihe. Acta Historica Astronomiae, 41, 112-132, 2010.

370 Filipiak, J., Przybylak, R., and Oliński, P.: The longest one-man weather chronicle (1721–1786) by Gottfried Reyger for
371   Gdańsk, Poland as a source for improved understanding of past climate variability. Int. J. Climatol., 39, 828-842, doi:
372   10.1002/joc.5845, 2019.

373 Gronau, K. L.: Hundertjährige meteorologische Tabellen der Witterung in Berlin von 1701 bis 1800. Ges. natf. Freunde z.
374   Berlin, 1, 1807.

375 Hellmann, G.: Das älteste Berliner Wetter-Buch 1700-1701 von Gottfried Kirch und seiner Frau Margaretha geb.
376   Winkelmann, Berlin, 1893.

377 Herbst, K.-D.: Gottfried Kirch (1639–1710). Astronom, Kalendermacher, Pietist, Frühaufklärer. Acta Calendariographica –
378   Forschungsberichte, 10, Jena, 2022.

379 Kadow, C., Wentzel, B., Jaekel, I., Bahlo, M., and Cubasch, U.: Berlin Climate Record - Daily Mean Temperature - Inner
380   City - 1766-1934. PANGAEA, https://doi.org/10.1594/PANGAEA.869767, 2016

381 Lenke, W.: Untersuchung der ältesten Temperaturmessungen mit Hilfe des strengen Winters 1708–1709. Berichte des
382   Deutschen Wetterdienstes, Nr. 92, Offenbach am Main, 1964.

383 Lundstad, E., Brugnara, Y. and Brönnimann, S.: Global Early Instrumental Monthly Meteorological Multivariable Database
384   (HCLIM). PANGAEA https://doi.pangaea.de/10.1594/PANGAEA.940724, 2022.

385 Lundstad, E., Brugnara, Y. Pappert, D., Kopp, J., Hürzeler, A., Andersson, A., Chimani, B., Cornes, R., Demarée, G.,
386   Filipiak, J., Gates, L., Ives, G. L., Jones, J. M., Jourdain, S., Kiss, A., Nicholson, S. E., Przybylak, R., Jones, P. D.,
387   Rousseau, D., Tinz, B., Rodrigo, F. S., Grab, S., Domínguez-Castro, F., Slonosky, V., Cooper, J., Brunet, N. and
388   Brönnimann, S.: Global historical climate database - HCLIM. Scientific Data, 10, 44, 2023.

389 Mädler, J.-H.: Die mittlere Temperatur Berlins für den Zeitraum 1701-1825 aus 126410 Beobachtungen berechnet. Zeitschr.
390 f. d. gesamt. Meteorol., 1, 1825.

391 Noone, S., Atkinson, C., Berry, D.I., Dunn, R.J.H., Freeman, E., Perez Gonzalez, I., Kennedy, J.J., Kent, E.C., Kettle, A.,
392 McNeill, S., Menne, M., Stephens, A., Thorne, P.W., Tucker, W., Voces, C., and Willett, K.M.: Progress towards a
393 holistic land and marine surface meteorological database and a call for additional contributions. Geosci Data J., 8, 103–
394 120, 2021. https://doi.org/10.1002/gdj3.109

395 Pappert, D., Barriendos, M., Brugnara1, Y., Imfeld, N., Jourdain, S., Przybylak, R., Rohr, C. and Brönnimann, S.: Statistical
396 reconstruction of daily temperature and sea-level pressure in Europe for the severe winter 1788/9. Clim. Past, 18, 2545–
397 2565, https://doi.org/10.5194/cp-18-2545-2022, 2022.

398 Pelz, J.: Die Wettertagebücher der Familie Kirch, 1700 bis 1774, Interne Information der WAG 6 2/78, 1978.

399 Pelz, J.: Die Berliner Jahresmitteltemperaturen von 1701 bis 1996. Beilage zur Berliner Wetterkarte, 6/97, 1997.

400 Pliemon, T., Foelsche, U., Rohr, C., and Pfister, C.: Subdaily meteorological measurements of temperature, direction of the
401 movement of the clouds, and cloud cover in the Late Maunder Minimum by Louis Morin in Paris, Clim. Past, 18, 1685–
402 1707, https://doi.org/10.5194/cp-18-1685-2022, 2022.

403 Przybylak, R., and Pospieszyńska, A.: Air temperature in Wrocław (Breslau) in the period 1710-1721 based on
404 measurements made by David von Grebner. Acta Agrophysica, 184, 35-43, 2010.

405 Reichen, L., Burgdorf, A.-M., Brönnimann, S., Rutishauser, M., Franke, J., Valler, V., Samakinwa, E., Hand, R., and
406 Brugnara, Y.: A Decade of Cold Eurasian Winters Reconstructed for the Early 19th Century. Nature Communications, 13,
407 2116, 2022, https://doi.org/10.1038/s41467-022-29677-8.

408 Schlaak, P.: 300 Jahre Wetterforschung in Berlin. Ihre Geschichte in Persönlichkeitsbildern. In: Fortak, H. (Ed.): 100 Jahre
409 Deutsche Meteorologische Gesellschaft in Berlin 1884–1984. Berlin, 1984. p. 86–124

410 Siebenhühner, H.: Das Wetterbuch der »Kirchin«. Fast 75jährige Beobachtungsreihe der Astronomenfamilie Kirch.
411 Berlinische Monatsschrift Heft 10/1996, 83-86, 1996.

412 Slivinski, L. C., Compo, G. P., Whitaker, J. S., Sardeshmukh, P. D., Giese, B. S., McColl, C., Allan, R., Yin, X., Vose, R.,
413 Titchner, H., Kennedy, J., Spencer, L. J., Ashcroft, L., Brönnimann, S., Brunet, M., Camuffo, D., Cornes, R., Cram, T. A.,
414 Crouthamel, R., Domínguez-Castro, F., Freeman, J. E., Gergis, J., Hawkins, E., Jones, P. D., Jourdain, S., Kaplan, A.,
415 Kubota, H., Le Blancq, F., Lee, T., Lorrey, A., Luterbacher, J., Maugeri, M., Mock, C. J., Moore, G. K., Przybylak, R.,
416 Pudmenzky, C., Reason, C., Slonosky, V. C., Smith, C., Tinz, B., Trewin, B., Valente, M. A., Wang, X. L., Wilkinson, C.,
417 Wood, K., and Wyszy´nski, P.: Towards a more reliable historical reanalysis: Improvements to the Twentieth Century
418 Reanalysis system, Q. J. Roy. Meteorol. Soc., 145, 2876–2908, 2019, https://doi.org/10.1002/qj.3598.

419 Valler, V., Franke, J., Brugnara, Y., and Brönnimann, S.: An updated global atmospheric paleo-reanalysis covering the last
420 400 years. Geosc. Data J., 9, 89–107, 2022.

421 Wanner, H., and Luterbacher, J.: The LOTRED approach – a first step towards a "paleoreanalysis" for Europe. Pages News
422 10: 9–11, 2002.

423 WMO: Chapter 6 - Quality Control Procedures. Guide on the Global Data-processing System, World Meteorological
424 Organization, Geneva, No. 305, VI.1-VI.27, ISBN 92-63-13305-0, 1993.