

# Evolution of winter precipitation in the Nile-River watershed since the last glacial

Vera D. Meyer<sup>1\*</sup>, Jürgen Pätzold<sup>1</sup>, Gesine Mollenhauer<sup>1,2</sup>, [Isla S. Castañeda<sup>3</sup>](#), [Stefan Schouten<sup>4</sup>](#), Enno Schefuß<sup>1\*</sup>

5 <sup>1</sup>MARUM – Center for Environmental Sciences, University of Bremen, Bremen, 28359, Germany

<sup>2</sup>Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, 27570, Germany

<sup>3</sup>[Department of Earth, Geographic and Climate Science, University of Massachusetts Amherst, Amherst, MA USA](#)

<sup>4</sup>[NIOZ Royal Netherlands Institute for Sea Research Department of Marine Microbiology and Biogeochemistry, 1790 AB Den Burg, Netherlands](#)

10 *Correspondence to:* Vera D. Meyer (vmeyer@marum.de), Enno Schefuß (eschefuss@marum.de)

**Abstract.** Between 14.5 and 5 ka BP, the Sahara was vegetated owing to a wet climate during the African Humid Period (AHP). However, the climatic factors sustaining the “Green Sahara” are still a matter of debate. Particularly the role of winter precipitation is poorly understood. Using the stable hydrogen isotopic composition ( $\delta D$ ) of high molecular weight (HMW) *n*-alkanoic acids in a marine sediment core from the Eastern Mediterranean (EM), we provide a continuous record for winter precipitation in the Nile-River delta spanning the past 18 ka. Pairing the data with regional  $\delta D$  records from HMW *n*-alkanes, we show that HMW *n*-alkanoic acids constantly derive from the delta while the HMW *n*-alkanes also receive significant contributions from the headwaters between ~15-1 ka BP due to enhanced fluvial runoff. This enables us to reconstruct the evolution of Mediterranean (winter) and monsoonal (summer) rainfall in the Nile River watershed in parallel. Heinrich Stadial 1 (HS1) evolved in two phases with a dry episode between ~17.5-16.0 ka BP followed by wet conditions between ~16-14.5 ka BP owing to movements of the Atlantic storm track. Winter rainfall enhanced substantially between 11-6 ka BP lagging behind the intensification of the summer monsoon by ca. 3 ka. Heavy winter rainfall resulted from a southern position of the Atlantic storm track combined with elevated sea-surface temperatures in the EM reinforcing local cyclogenesis. We show that during the “Green Sahara” monsoon precipitation and Mediterranean winter rainfall were simultaneously enhanced and infer that the winter-rainfall zone extended southwards delivering moisture to the Sahara. Our findings corroborate recent hypotheses according to which southward extended winter rains were a crucial addition to the northward displacement of the summer monsoon helping to sustain a “Green Sahara”.

## 1 Introduction

North Africa underwent dramatic oscillations between dry and wet climate states in the course of glacial-interglacial cycles ([e.g., deMenocal et al., 2000a; Menviel et al., 2021; Larrasoana et al., 2013; Ziegler et al., 2010](#)). The last wet phase, known as the “African Humid Period” (AHP), occurred between 14.5-5 ka BP reaching its climax between 11-6 ka BP (e.g. deMenocal et al., 2000a; Shannahan et al., 2015; Tierney et al., 2017; Dupont and Schefuß. 2018). Humid conditions in North Africa

transformed the formerly barren, hyperarid Sahara Desert into a fertile “Green Sahara” (~11.5-5 ka BP; Kuper and Kröpelin, 2006) where savannah, lakes, rivers and wetlands existed allowing human settlements in the Sahara (Kuper and Kröpelin, 2006; Quade et al., 2018; Larrasoña et al., 2013; Jolly et al., 1998). Although intensively studied, the drivers and spatio-temporal extent of the AHP are still a matter of debate (Lüning and Vahrenholt, 2019; Kutzbach et al., 2014; Otto-Bliesner et al., 2014; Menviel et al., 2021; Tierney et al., 2017; Sha et al., 2019; Cheddadi et al., 2021). There is consensus that the AHP initiated in response to insolation forcing which intensified the African summer monsoon and shifted the Intertropical Convergence Zone and the African rainbelt northward (Menviel et al., 2021; Lüning and Vahrenholt, 2019; Pausata et al., 2016; Braconnot et al., 2007; Claussen et al., 2017; deMenocal et al., 2006b). Next to orbital forcing positive, non-linear feedbacks from the land surface amplified the climatic changes (e.g. Chandan et al., 2020; Pausata et al., 2016; 2020). Controversy exists on the termination of the AHP (Kuper and Kröpelin, 2006; Shannahan et al., 2015; Schefuß et al., 2005; Costa et al., 2014; Blanchet et al., 2014; deMenocal et al., 2000a; Tierney and deMenocal, 2013; Ménot et al., 2020; Tierney et al., 2008; Berke et al., 2012; Weldeab et al., 2014; Junginger et al., 2014; Castañeda et al., 2016a; deMenocal, 2015; Collins et al., 2017) as well as on the climatic processes sustaining a vegetated Sahara (Cheddadi et al., 2021, Kutzbach et al., 2014; Alpert et al., 2006; Chandan et al., 2020; Braconnot et al., 2007; Hopcroft et al., 2017; Claussen et al., 2017; Sha et al., 2019; Tierney et al., 2017). Paleoclimate records suggest both a gradual as well as an abrupt ending of the AHP and indicate that the AHP terminated earlier in the North than in the South (Kuper and Kröpelin, 2006; Shannahan et al., 2015; deMenocal, 2015). For a long time, most studies focused on the northward extension of the summer monsoon seeking to explain the AHP and the Green Sahara (deMenocal et al., 2006b; Braconnot et al., 2007; Menviel et al., 2021; deMenocal et al., 2000b; Shannahan et al., 2015; Sha et al., 2019; Tierney et al., 2017). However, reproducing the AHP and the “Green Sahara” has challenged climate modellers for a long time because of inconsistencies between model results and proxy data, a problem that has not been overcome, yet (Chandan and Peltier., 2020; Braconnot et al., 2007; Hopcroft et al., 2017; Perez-Sanz et al., 2014). Climate models commonly suggest a northward extension of the summer monsoon (Chandan and Peltier., 2020; Braconnot et al., 2007; Hopcroft et al., 2017; Perez-Sanz et al., 2014) but in northern Africa these models underestimate precipitation amount inferred from the pollen-based reconstructions (e.g. Bartlein et al., 2011; Hély et al., 2014) and isotopic compositions of biomarkers and speleothems (Tierney et al., 2017; Sha et al., 2019). The representation of positive feedbacks from the land surface on the monsoon may partly account for the discrepancies as these feedbacks have been shown to substantially amplify the response of the monsoon to orbital forcing pushing the monsoon further north than orbital forcing alone (Chandan and Peltier, 2020; Pausata et al., 2016). Nevertheless, inconsistencies with proxy data remain (e.g. Chandan and Peltier, 2020; Pausata et al., 2016). The simulated precipitation in northern Africa is still too low compared to proxy data (Pausata et al., 2016; Chandan and Peltier, 2020; Bartlein et al., 2011) but are compatible with rainfall estimates based on leaf-wax isotopic compositions (Tierney et al., 2017). Given these inconsistencies contributions from additional moisture sources such as the Mediterranean winter rains increasingly receive attention seeking to explain the green Sahara (Kutzbach et al., 2014; Cheddadi et al., 2021; Tierney et al., 2017). Recently, Cheddadi et al. (2021) suggested that intensified Mediterranean winter rainfall and a southward

extension of the winter rainfall zone into the Sahara may have delivered the additional moisture needed for sustaining a green Sahara by decreasing rainfall seasonality. Beforehand, Tierney et al. (2017) had already invoked additional winter precipitation to fully explain their observed amplitudes in precipitation records from northwestern Africa. A model-based investigation of the response of Mediterranean winter rainfall confirms intensified precipitation in northern Africa and the Middle East during the AHP in response to orbital forcing during the AHP (Kutzbach et al., 2014).

70 However, the comparison of simulations and vegetation reconstructions shows that climate models probably underestimate precipitation in the northern Sahara and that the summer monsoon alone may have been insufficient to sustain a vegetated Sahara (Chandan et al., 2020; Braconnot et al., 2007; Hoperoft et al., 2017; Perez Sanz et al., 2014; Cheddadi et al., 2021; Hély et al., 2014). Recently, Cheddadi et al. (2021) suggested that intensified Mediterranean winter rainfall and a southward

75 extension of the winter rainfall zone into the Sahara may have delivered the additional moisture needed for sustaining a Green Sahara by decreasing rainfall seasonality. Also, Tierney et al. (2017) have already invoked additional winter precipitation to fully explain their observed amplitudes in precipitation change in Northwest Africa. Unfortunately, the glacial-to-Holocene development of winter precipitation in ~~northeast~~northeastern Africa remains elusive given the scarcity of proxy records. In order to improve the understanding of how winter and summer precipitation evolved around the AHP, continuous records for

80 precipitation are required to robustly investigate spatial variations in rainfall across North Africa. Constraints on the hydroclimatic development in the southeastern Mediterranean come from Earth System Models and various paleoenvironmental archives from the Asian borderlands of the Levantine Basin (Langgut et al., 2016; Tierney et al., 2022; Stein, 2001; Ludwig and Hochmann, 2022; Bar-Mathews et al., 1997; Goldsmith et al., 2017; Enzel et al., 2008; Cheng et al., 2015; Miebach et al., 2019). While there is consensus that during the Last Glacial Maximum (LGM) the eastern Mediterranean

85 realm was characterized by a more positive moisture balance (Stein, 2001; Goldsmith et al., 2017; Ludwig and Hochmann, 2022), the effects from changes in evaporation and precipitation on effective moisture are still debated. Some authors proposed increased precipitation (Enzel et al., 2008; Stein et al., 1997; Cheng et al., 2015) while others found indication for a reduction (Ludwig and Hochmann, 2022; Bar-Mathews et al., 1997; Langgut et al., 2021). While the sign of rainfall changes is ambiguous, there is a broad consensus among proxy data and model results that evaporation was lower under cold glacial

90 conditions (Stein, 2001; Tierney et al., 2022; Ludwig and Hochmann, 2022; Goldsmith et al., 2017). These uncertainties in terms of rainfall ask for more studies addressing the hydroclimate development in the Eastern Mediterranean since the last glacial.

A key region to study ~~Northeast~~northeastern-African climate change is the Nile-River basin which - extending over 3 million

95 km<sup>2</sup> (Figure 1) - currently is influenced by monsoonal summer rains south of the Sahara Desert and Mediterranean winter rainfall in the delta region. Furthermore, it is of societal relevance to address the climatic history of the Nile River catchment because the river is the lifeline of Egypt providing fertile ground and drinking water to millions of people. It also played an ~~important~~ vital role in the rise and demise of ancient Egyptian civilizations (e.g. Zaki et al., 2021). In northeastern Africa and the borderlands of the southern Levantine basin socioeconomic impacts due to increasing aridity are expected in the course of

100 [future warming](#). Continuous archives for precipitation are predominantly found in the headwaters of the Nile River and further south (Lakes Victoria, Tana, Tanganyika; [Figure 1](#); Berke et al., 2012; Costa et al., 2014; Tierney et al., 2008; 2010). In the northern part of the catchment where the hyperarid Sahara Desert extends, continuous records for precipitation are sparse as sedimentary sequences from the deglaciation and the Holocene - such as lacustrine deposits - were subject to strong wind erosion during arid periods (Hamdan et al., 2016; Hamdan and Lucarini, 2013). Sediment cores from the Levantine Basin  
105 (Figure 1) have often been used to reconstruct environmental changes in the Nile-River watershed. These records are commonly considered as integrators of the entire catchment and are mostly interpreted to reflect monsoonal rainfall variability (Castañeda et al., 2016a; Revel et al., 2015; Ménot et al., 2020, Blanchet et al., 2014). However, the Nile crosses several climate regimes which drastically differ in precipitation amount and seasonality (tropical conditions in the headwaters, hyperarid desert in its central part and Mediterranean winter rainfall in the delta region). For climate reconstructions it is crucial  
110 to address these climate zones separately in order to identify latitudinal differences and to understand how monsoonal (summer) and Mediterranean (winter) precipitation evolved around the AHP.



Here, we provide a new hydroclimate record based on the stable hydrogen isotopic composition ( $\delta^2\text{H}$  or also often termed:  $\delta\text{D}$  [where D stands for deuterium](#)) of HMW *n*-alkanoic acids. HMW *n*-alkanoic acids are major components of epicuticular leaf-waxes of higher plants (Eglinton and Hamilton, 1967). The record is obtained from marine sediment core GeoB7702-3 from the Eastern Mediterranean Sea (EM; Figure 1).  $\delta\text{D}$  of leaf-wax lipids ( $\delta\text{D}_{\text{wax}}$ ) is a powerful means to reconstruct past hydrological changes (e.g. Sachse et al., 2012) and has been successfully applied to infer hydroclimate variability across Africa (Schefuß et al., 2005; Tierney et al., 2008; 2017; Berke et al., 2012; Costa et al., 2014; Collins et al., 2013; Castañeda et al., 2016a; [Konecky et al., 2011](#)). We infer that  $\delta\text{D}_{\text{wax}}$  of our HMW *n*-alkanoic acids records winter precipitation in the Nile-delta  
115 region. By comparison to existing  $\delta\text{D}_{\text{wax}}$  records based on HMW *n*-alkanes from the Levantine Basin (Castañeda et al., 2016a) and HMW *n*-alkanoic-acid based  $\delta\text{D}_{\text{wax}}$  records from the headwaters (Berke et al., 2012; Costa et al., 2014; Tierney et al., 2008) we are able to address the individual hydroclimate developments of the southern and northern sections of the Nile-River catchment and shed light onto the interplay of Mediterranean (winter) and monsoonal (summer) rainfall changes in [Northeastnortheastern](#) Africa around the AHP.  
120

## 125 **2 Study Area**


Core GeoB7702-3 was recovered from the southeastern Levantine Basin off Israel (Figure 1) (Pätzold et al., 2003). The core receives terrigenous material from the Nile River as the Nile suspension load is transported eastward along the continental margin due to the anticlockwise direction of surface currents and eddies (e.g. Weldeab et al., 2002). This makes site GeoB7702-3 a suitable archive for environmental changes in the Nile-River basin (Castañeda et al., 2010a; 2016a).  
130 The Nile River is the longest river in the world extending over 34° of latitude (Figure 1). The catchment spans from Equatorial Africa to the Mediterranean coast draining Uganda, Ethiopia, South Sudan, Sudan and Egypt (Figure 1). The river consists of three major tributaries, i.e. the White Nile (sourced from Lake Victoria), the Blue Nile (sourced from Lake Tana) and the

Atbara Nile (source is situated north of Lake Tana). The confluence of the three tributaries forms the Main Nile (Figure 1). For the following discussion, we define three sub-catchments of the watershed as follows: upper catchment (headwaters in Ethiopia and Uganda; 4°S to 15°N), middle catchment (Sahara, Sudan and Egypt; 15-30°N) and lower catchment (Nile delta; 30-31°N). On its way to the North the Nile River crosses different climate zones and vegetation regimes. The climate zones encompass Mediterranean climate in the delta area, the hyperarid Sahara Desert, the semiarid Sahel zone and a wet, tropical


climate in Ethiopia and Uganda (e.g. Korecha and Barnston, 2007). In the lower and middle catchment, rainfall mainly occurs




- |               |               |
|---------------|---------------|
| 1: GeoB7702-3 | 2: MS27PT     |
| 3: Soreq Cave | 4: Jeita Cave |








(A)



(B)



- |                 |               |
|-----------------|---------------|
| 1: GeoB7702-3   | 2: MS27PT     |
| 3: Soreq Cave   | 4: Jeita Cave |
| 5: DSDDP 5017-1 |               |

140

145

Figure 1: General wind patterns over Africa during winter (A) and summer (B). The star marks site GeoB7702-3. The modern positions of the ITCZ during January (A) and July/August (B) are illustrated. CAB means Congo Air Boundary which separates Atlantic and Indian air masses over equatorial Africa. In (A) the positions of the North Atlantic storm track are sketched for positive and negative phases of the North Atlantic Oscillation (NAO). (C) Detailed map of the Nile-River watershed. Site GeoB7702-3 is marked by a red star. Other locations mentioned in the text are indicated by circles. The maps were created using Ocean Data View 5.6.3 (Schlitzer, 2006).

during winter. In the Nile delta annual rainfall spans from 1481 mm/yr at the coast (Alexandria; <https://en.climate-data.org>) to 18 mm/yr at its southern edge (Cairo; <https://en.climate-data.org>). Most rain falls between October and March. The Sahara usually does not receive any rainfall (e.g. Luxor; <https://en.climate-data.org>). The upper section of the watershed is characterized by heavy rainfall (e.g. about 1874 mm/yr annually in Addis Ababa/Ethiopia; about 1747 mm/yr in Kampala/Uganda; <https://en.climate-data.org>). In Ethiopia, most rain falls between June and September (<https://en.climate-data.org>). Uganda receives year-round precipitation with rainfall maxima during March till [Mai-May](#) and September till November (<https://en.climate-data.org>).

Precipitation in the upper Nile River watershed is mainly determined by the West African Monsoon, and thus related to the seasonal migration of the Intertropical Convergence Zone (ITCZ; Figure 1). During the summer months, the seasonal northward movement transports moisture-laden air to southern [Northeastnortheastern](#) Africa (up to ~15°N). Additionally, the Congo Air Boundary controls the relative contribution of moisture from the Gulf of Guinea, i.e. the Atlantic Ocean, and Indian Ocean to the headwaters (Figure 1) (Camberlin, 2009). North of 15°N the catchment is predominantly under the influence of the westerlies receiving moisture from the Atlantic Ocean and the Mediterranean Sea, the Arabian Peninsula and the Red Sea (Figure 1) (Viste and Sortberg, 2013). The Mediterranean climate is characterized by dry summers and wet winters. [Winter precipitation is governed by cyclogenesis over the eastern Mediterranean, so called “Cyprus Lows”. The cyclones form when relatively cold European air encounters the warm Mediterranean Sea during southward advection \(Alpert et al., 1990\). Next to the Cyprus Lows the Red Sea Trough is another local atmospheric circulation pattern controlling winter rainfall. It is commonly associated with warm and dry conditions but variations in its position can cause heavy precipitation events \(Tsvieli and Zangvil, 2005; Krichak et al., 2012\).](#) -The winter climate is largely dependent [to teleconnections with the on the](#) North Atlantic Oscillation (NAO) which forms a constituent of the Arctic Oscillation. Modulating the positions of the Atlantic and Mediterranean storm tracks (Figure 1) the NAO exerts strong control over moisture delivery to the Mediterranean borderlands. Warm and wet winters [in the Mediterranean](#) are generally associated with negative NAO-states while cold and dry winters occur during positive phases ([Eshel and Farrel, 2000](#)).

## 3 Material and Methods

### 3.1 Core Material and Chronology

Gravity core GeoB7702-3 was recovered from the continental slope off Israel at 562 m water depth during RV Meteor cruise M52/2 in 2002 (Pätzold et al., 2003). Prior to sampling the core was stored at 4°C. Age control for this core was previously established by Castañeda et al. (2010a) and is based on accelerator mass spectrometry (AMS) radiocarbon dates of planktic foraminifera. We refined the age model using up-to-date calibration curves and nine additional AMS dates (Table 1). To isolate foraminifera the samples were wet-sieved and specimens of the foraminifera *Globigerinoides ruber* alba were hand-picked from the (150-63 µm fraction). In cases where the abundance of *G. ruber* was insufficient, we mixed planktonic foraminifera species to obtain enough material (Table 1). 290-810 µg carbon was dated at the MICADAS AMS-dating facility at the Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research (Bremerhaven, Germany) according to [inhousein-house](#)



180 protocols (Mollenhauer et al., 2021). The dates were combined with the data set of Castañeda et al. (2010a) to create the age-  
depth model. All radiocarbon dates are listed in Table 1. The BACON 2.5.8 software (Blauw and Christen, 2011) was used  
for age-depth modelling. Radiocarbon ages were transferred into calendar ages based on the Marine20 calibration curve  
(Heaton et al., 2020). Today, the mean reservoir age offset ( $\Delta R$ ) in the Levantine Basin is  $-94 \pm 94$  yrs in relation to Marine20  
~~at present~~ (marine reservoir correction database; Reimer et al., 2001). BACON was run with a constant  $\Delta R = -100 \pm 100$  yrs,  
185 accordingly. Default settings were used for the prior, apart from the accumulation ~~rate prior~~, which was set to 50 yrs/cm.  
BACON operated with 118 core-slices. The age-depth model for GeoB7702-3 and the prior distributions are plotted in Figure  
2.

~~For biomarker analysis the core was sampled in 5-12 cm steps providing a mean temporal resolution of ~350 years between samples.~~

### 190 3.2 Lipid extraction, quantification and isotopic analysis

~~For biomarker analyses, a total of 51 samples were collected every 5-12 cm, providing a mean temporal resolution of ~350 years between samples. Target biomarkers of our study are HMW *n*-alkanes and HMW *n*-alkanoic acids and their stable hydrogen isotopic compositions.~~ The lipid extraction and isotopic analysis were performed at MARUM - Centre for Marine Environmental Sciences (University of Bremen, Germany). The sediment samples were freeze-dried and afterwards  
195 homogenized using a mortar and pestle. The soluble organic matter was extracted from ~5 g of sediment using an Accelerated Solvent Extractor (ASE200). Extraction was performed with three cycles lasting 5 minutes each, using Dichloromethane (DCM)-Methanol (9:1 v/v) at 100°C and 1000 psi. 19-Methyl-Arachidic Acid was added to the samples as internal standard prior to extraction. The total lipid extracts were saponified using potassium hydroxide (KOH). ~~Neutral lipids and fatty acids were recovered using *n*-hexane and DCM, respectively.~~ The fatty acids were methylated using methanol of known isotopic  
200 signature. The fatty-acid methyl esters (FAMES) were cleaned by means of column chromatography. Columns consisted of 4 cm deactivated silica (1% H<sub>2</sub>O in *n*-hexane) and 0.5 cm sodium sulfate (Na<sub>2</sub>SO<sub>4</sub>) in a Pasteur pipette with 5 mm diameter. FAMES were recovered using ~~Dichloromethane~~ DCM:Hexane (2:1). ~~FAMES were quantified by gas chromatography and flame ionization detection (GC-FID) using a Thermo FOCUS GC. The GC was equipped with a Restek Rxi-5ms column. FAMES were analysed using a similar method as in Gensel et al. (2022).~~ For the quantification of the FAMES of HMW *n*-alkanoic  
205 ~~acids the response factors of the target compounds were determined from an external standard mixture containing five HMW *n*-alkanoic acid homologues (C<sub>24:0</sub>, C<sub>26:0</sub>, C<sub>28:0</sub>, C<sub>30:0</sub> and C<sub>32:0</sub>) at a concentration of 50 ng/μl. The external standards were run every six sample. The relative standard deviation of the peak areas was <9%.~~

Isotope analysis of stable hydrogen ( $\delta D$ ) was performed using gas chromatography coupled to isotope ratio mass spectrometry (GC-IRMS). We used a Thermo Trace GC coupled to a MAT253 MS. Isotope values were measured against calibrated  
210 reference gas (H<sub>2</sub>). Values are reported in per mil relative to the VSMOW standards. A standard mixture consisting of 16 *n*-alkanes was run every sixth sample in order to monitor the performance of the system. For the  $\delta D$  analysis the accuracy and precision (mean deviation from offline values and the respective relative standard deviation, RSD) were 2.2‰ and 3.0‰,



215 respectively. The machine was operated only when the average absolute deviation from offline values was <5%. The  $H_3^+$  - factor was measured daily and was  $5.6 \pm 0.1$  throughout the measurement series. Replicate measurements of the samples yielded a standard deviation of 0.1-3.8‰ for  $\delta D$ . We report the  $\delta D$ -signatures of the  $n$ - $C_{26:0}$  and  $n$ - $C_{28:0}$  alkanolic acids as they turned out to be the most abundant homologues in our samples. The  $\delta D$  of the respective FAMES were corrected for the bias introduced during the methylation process using isotope mass balance (hereafter  $\delta D_{wax\ n\text{-alkanoic\ acid}}$ ).  $\delta D_{wax\ n\text{-alkanoic\ acid}}$  was further corrected for deglacial changes in global ice-volume applying stacked data of oxygen isotopic compositions ( $\delta^{18}O$ ) of benthic foraminifera (L04-stack; Lisiecki and Raymo, 2005) and the approach described in Ruan et al. (2019). We additionally analyzed the stable carbon isotopic composition of HMW  $n$ -alkanoic acids ( $\delta^{13}C_{wax\ n\text{-alkanoic\ acids}}$ ) as described in the supplementary material (S1).

220 Prior to this study, a separate set of samples from GeoB7702-3 were extracted at the Royal Netherlands Institute for Sea Research (NIOZ) (Castañeda et al, 2010). As described in Castañeda et al. (2016), neutral lipids were split in apolar, keto and polar fractions by alumina oxide column chromatography with the following solvent mixtures: hexane:dichloromethane (9:1 v/v), hexane:DCM (1:1 v/v) and DCM:methanol (1:1 v/v). The apolar hydrocarbon fraction was further separated into saturated and unsaturated compounds using silica gel coated with silver nitrate ( $AgNO_3$ ). Identification of  $n$ -alkanes was performed on a Thermo Finnigan Trace Gas Chromatograph (GC) Ultra coupled to Thermo Finnigan DSQ mass spectrometer (MS) using a CP Sil-5 fused silica capillary column (25 m $\times$ 0.32 mm; film thickness 0.12  $\mu$ m) with helium as the carrier gas. Mass scans were made from  $m/z$  50 to 800, with 3 scans per second and an ionization energy of 70 eV. The oven program started at 70°C, increased by 20°C min<sup>-1</sup> to 130°C, and subsequently by a rate of 4°C min<sup>-1</sup> until 320°C (held for 10 min). Quantification of  $n$ -alkanes was performed on an HP 6890 gas chromatograph (GC) using a 50 m CP Sil-5 column (0.32 mm diameter, film thickness 0.12  $\mu$ m) and helium as the carrier gas, and with the same oven temperature program as for GC-MS. Compound concentrations were determined by relating chromatogram peak areas to the concentration of an internal standard of known concentration. Details regarding hydrogen and carbon isotopic analyses of the  $n$ - $C_{29}$  and  $n$ - $C_{31}$  alkanes are reported by Castañeda et al. (2016a).

235 ~~Next to  $\delta D$ , we also analyzed the stable carbon isotopic composition of HMW  $n$  alkanolic acids ( $\delta^{13}C_{wax\ n\text{-alkanoic\ acids}}$ ) as described in the supplementary material (S1).~~

Note that the set of samples taken for HMW  $n$ -alkanoic acids (this study) are at different sampling depths from the HMW  $n$ -alkanes (Castañeda et al., 2016a).

240



**Figure 2** Output graph for the age-depth modelling approach for core GeoB7702-3 using the BACON software (Blaauw and Christen, 2011). The model is based upon the radiocarbon dates of foraminifera given in Table 1. The small panels on top represent the Markov Chain Monte Carlo iterations (left) and the distributions of the prior (green lines) and the posterior (grey histograms) for the accumulation rate (middle) and memory (right). The bottom panel illustrates the age-depth model. The blue dots depict the calibrated <sup>14</sup>C-ages. The grey stippled lines mark the 95% confidence intervals. The red line reflects the single best model based on the mean age for each depth. For more technical information the reader is referred to Blaauw and Christen (2011).

245

**Table 1: List of AMS-dates and corresponding calendar ages (cal. age) used to establish the revised age-depth model of GeoB7702-3. AMS dates were compiled from this study and Castañeda et al. (2010a; 2010b). Samples from this study were dated at the MICADAS-dating facility at Alfred Wegener Institute (Bremerhaven, Germany) while Castañeda et al. (2010a; 2010b) performed dating at the Leibnitz Laboratory for Radiometric Dating and Stable Isotope Research (University of Kiel, Germany). Our new age-depth model is based upon the median values of the calendar ages calculated.**

Depth [cm]	Sample label	Dated material	AMS date [ <sup>14</sup> C a BP]	2σ [±]	Cal. age min [a BP]	Cal. age max [a BP]	Cal. age median [cal a BP]	Cal. age mean [cal a BP]	Reference of AMS date
0	set*	-	100*	10	64	124	98	97	this study
10	KIA25649	G. ruber and O. universa	245	30	123	415	221	231	Castañeda et al. (2010b)
64.5	KIA25648	G. ruber and G. sacculifer	1725	25	1086	1682	1342	1356	Castañeda et al. (2010b)
81.5-84.5	6279.2.1	G. ruber	2340	26	1615	2285	1929	1933	this study
102	KIA24619	G. ruber	2965	55	2288	2849	2580	2576	Castañeda et al. (2010b)
130-133	6281.1.1	G. ruber	3586	24	3113	3685	3400	3400	this study
132	KIA24617	G. ruber	3500	35	3189	3643	3413	3416	Castañeda et al. (2010b)
198-201	6277.2.1	G. ruber	5399	25	5353	6014	5697	5697	this study
210	KIA24616	G. ruber	5600	40	5764	6383	6036	6043	Castañeda et al. (2010b)
231-234	7307.1.1	G. ruber	7393	45	7242	8138	7724	7718	this study
242.5	KIA25646	G. ruber and G. sacculifer	7845	40	8023	8866	8327	8358	Castañeda et al. (2010b)
251-254	6280.1.1	G. ruber	9309	28	9015	10110	9663	9652	this study
257	KIA24613	G. ruber	9070	60	9564	10290	9934	9928	Castañeda et al. (2010b)
278-281	6276.2.1	mixed planktonic foraminifera	10144	34	11051	12047	11502	11508	this study
279	KIA24612	mixed planktonic foraminifera	10470	70	11128	11839	11457	11463	Castañeda et al. (2010b)
297-300	6275.2.1	G. ruber	11827	32	12688	13729	13215	13213	this study
310	KIA24611	mixed planktonic foraminifera	12580	80	13593	14509	14035	14037	Castañeda et al. (2010b)
356	KIA24609	G. ruber	14130	100	16073	16781	16432	16431	Castañeda et al. (2010b)
359-362	6274.2.1	G. ruber	14420	37	16264	17074	16666	16664	this study
391	KIA24608	G. ruber	15830	120	17610	18515	18036	18048	Castañeda et al. (2010b)
393-396	6262.2.1	G. ruber	15275	39	17690	18728	18151	18176	this study
452	KIA25652	G. ruber	18810	150	21025	22260	21750	21717	Castañeda et al. (2010b)
497	KIA24605	G. ruber	20660	180	23106	24215	23747	23716	Castañeda et al. (2010b)
540.5	KIA24604	G. ruber	21840	220	24579	25711	25192	25179	Castañeda et al. (2010b)
581.5	KIA25653	mixed planktonic foraminifera	22230	190	25715	27340	26397	26436	Castañeda et al. (2010b)

\* the core-top age was defined as 100 calendar ages in order to guarantee BACON calculates positive dates for the depths above the first AMS-date (i.e. 10 cm).


#### 4. Results

The concentrations of HMW *n*-alkanoic acids and HMW *n*-alkanes are shown in Figure 3. Concentrations of HMW *n*-alkanoic acids are slightly lower during the early deglaciation (between 18-17 ka BP) than during the late Holocene (Figure 3c). At ~16.5 ka BP they decrease to a minimum. From 14.6 ka BP onwards the concentrations progressively increase in two steps culminating in a maximum at 10 ka BP (Figure 3c). Afterwards they decrease until the middle Holocene and increase again from 2 ka BP onwards. The concentrations of HMW *n*-alkanes show a similar development but differences are evident. The concentrations of HMW *n*-alkanes share the two-step increase starting at 14.6 ka BP and subsequent decline as well as the rise during the late Holocene (Figure 3b,c). However, while concentrations of HMW *n*-alkanoic acids are lowest during the early deglaciation (18-14.6 ka BP) the abundances of HMW *n*-alkanes are higher or similar to Holocene values (Figure 3b). Between ~17.5-16.5 ka BP concentrations of HMW *n*-alkanes are prominently elevated, reaching their maximum at that time. By contrast, HMW *n*-alkanoic acids reach maximum concentrations around 10 ka BP (Figure 3b,c).

$\delta D_{\text{wax } n\text{-alkanoic acid}}$  are given in Figure 4.  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  range between -104 and -153‰. The *n*-C<sub>26:0</sub> and *n*-C<sub>28:0</sub> alkanoic acids behave similarly regarding  $\delta D$ -values and trends (Figure 2d,d); green and orange lines). In general,  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  is higher during the deglacial than during the Holocene (Figure 2d,d). Two episodes of remarkable changes in  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  are evident in the record, i.e. between ~17.5-14.5 ka BP and between ~11-6 ka BP. At 17.5 ka BP  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  rapidly increase and reach maximal values at 17.0 ka BP. Afterwards, at ~16 ka BP,  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  decrease again reaching values similar to the late Holocene between 16.0 and 14.5 ka BP. At 14.5 ka BP  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  slightly increase again and then remain relatively constant until a striking minimum between 10.0-6.0 ka BP is registered between 10.0-6.0 ka BP. The minimum begins and terminates abruptly (where abruptly is defined as within a few hundred years). At this time HMW *n*-alkanoic acids became depleted by 34 ‰, relative to the values found at 12 ka BP (Figure 2d,d). After the minimum there is little variability throughout the late Holocene.

Castañeda et al. (2016a) investigated the hydroclimate development in the Nile River watershed using  $\delta D_{\text{wax}}$  of HMW *n*-alkanes (another leaf wax lipid;  $\delta D_{\text{wax } n\text{-alkanes}}$ ) in core GeoB7702-3 (Figure 2d, dark green line). Comparing our  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  with previously published  $\delta D_{\text{wax}}$  of HMW *n*-alkanes (another leaf-wax lipid;  $\delta D_{\text{wax } n\text{-alkanes}}$ ) from core GeoB7702-3 (Figure 4d, blue line Castañeda et al., 2016a) reveals distinct discrepancies. While  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  and  $\delta D_{\text{wax } n\text{-alkanes}}$  from GeoB7702-3 are similar before 15 ka BP and at 1 ka BP, they differ in the long-term trend of the glacial-to-Holocene evolution. The amplitude of the long-term development in the  $\delta D_{\text{wax } n\text{-alkanes}}$  is larger than in the  $\delta D_{\text{wax } n\text{-alkanoic acid}}$ , which is why the  $\delta D_{\text{wax } n\text{-alkanes}}$  are more negative than the  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  (up to 25 ‰; Figure 4d). This difference in amplitude mainly stems from a progressive decrease in  $\delta D_{\text{wax } n\text{-alkanes}}$  from 15 ka BP onwards and from a progressive increase after ~5 ka BP (Figure 4d) this is not recorded by  $\delta D_{\text{wax } n\text{-alkanoic acid}}$ . In contrast to the  $\delta D_{\text{wax } n\text{-alkanes}}$ , the  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  remained quite stable between 15-11 ka BP and after ~6 ka BP (Figure 4d). While  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  and  $\delta D_{\text{wax } n\text{-alkanes}}$  are similar during the periods between 18-14.5 ka BP and at 1 ka BP (Figure 2d)  $\delta D_{\text{wax } n\text{-alkanes}}$  are more negative than  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  for most of the time (up to ~25 ‰). The

285 offset results from different long term developments.  $\delta D_{wax\ n\text{-alkanoic\ acid}}$  was rather stable between 15-11 ka BP whereas  $\delta D_{wax\ n\text{-alkanes}}$  progressively become more negative from 15 ka BP onwards. Likewise, constancy is found in  $\delta D_{wax\ n\text{-alkanoic\ acid}}$  between 6 ka BP till present while  $\delta D_{wax\ n\text{-alkanes}}$  progressively increase throughout this period (Figure 2d). Both records agree with respect to the maximum at ~17 ka BP and the minimum between 10-7 ka BP (Figure 2d).



290 **Figure 3: Concentrations of HMW *n*-alkanoic acids (even numbered homologues  $\Sigma C_{26:0}$ - $C_{32:0}$ ; green (c) and HMW *n*-alkanes (odd**  
 295 **numbered homologues  $\Sigma C_{29}$ - $C_{33}$ ; blue (b) in core GeoB7702-3. Black lines denote running averages. (a) endmember-modelling**  
**results from the grain size analysis in core GeoTü SL112 adopted from Hamann et al. (2008). The contribution of endmember 1**  
**(EM1) to the silt fraction represents Saharan dust input to the Eastern Mediterranean (Hamann et al., 2008). (d) The oxygen isotopic**  
**composition of the planktic foraminifera species *Globigerinoides ruber alba* ( $\delta^{18}O_{G.ruber\ alba}$ ) from core MS27PT (Figure 1c), Nile deep**  
**sea fan (Revel et al., 2010; 2015), is given to reflect freshwater runoff in the Nile River. YD: Younger Dryas stadial, B/A:**  
**Bølling/Allerød interstadial, HS1: Heinrich Stadial 1, LGM: Last Glacial Maximum. The light grey and dark grey shadings mark**  
**the episodes of the AHP and the Green Sahara, and their optimum. The yellow bar denotes the interval when Lakes Tana and**  
**Victoria desiccated (Stager et al., 2011; Lamb et al., 2007; Marshall et al., 2011; Williams et al., 2006).**






Figure 4: (a)-(c)  $\delta D_{wax}$  records from Lakes Tana (Costa et al., 2014), Victoria (Berke et al., 2012) and Tanganyika (Tierney et al., 2008) reporting hydroclimate variability in tropical East Africa; (d)  $\delta D_{wax}$   $n$ -alkanoic acids (orange:  $n$ -C<sub>26:0</sub>; green:  $n$ -C<sub>28:0</sub>; this study) along with  $\delta D_{wax}$   $n$ -alkane (blue:  $n$ -C<sub>31</sub>; Castañeda et al., 2016a; 2016b) from core GeoB7702-3 and  $\delta D_{wax}$  based on the  $n$ -C<sub>30:0</sub> alkanolic acid from core DSDDP 5017-1 from the Dead Sea (black; Tierney et al., 2022); (e) Ratio of tetra and penta-methylated brGDGTs from the Nile-deep sea fan (core MS27PT; Figure 1) reporting on the input of delta-derived organic matter versus material from the soils in the upper Nile-River catchment to the Eastern Mediterranean (Ménot et al., 2020); (f) oxygen isotopic composition of the planktic foraminifera species *Globigerinoides ruber alba* from the Levantine Basin (core MS27PT; Figure 1) used to reconstruct changes in salinity related to Nile River runoff (Revel et al., 2010; 2015). The light grey and dark grey shadings mark the episodes

310 of the AHP and the Green Sahara, and their optimum. LGM: Last Glacial Maximum; HS1: Heinrich Stadial 1; B/A: Bølling-Allerød interstadial; YD: Younger Dryas stadial.

## 5. Discussion

$\delta D_{wax}$  is dependent to the  $\delta D$  of the source water taken up by the plant during biosynthesis and thus can be used as tracer for  $\delta D$  of precipitation (e.g. Sachse et al., 2012 and references within; Tierney et al., 2017). This, in turn, varies along with hydrological processes including rainfall amount, evapotranspiration and moisture source but also temperature changes affect the isotopic composition (Sachse et al., 2012). In low latitude regions  $\delta D_{wax}$  is predominantly influenced by rainfall amount (amount effect) as temperature effects are negligible (Sachse et al., 2012). This makes  $\delta D_{wax}$  a powerful tool to reconstruct glacial-interglacial fluctuations in rainfall amount and moisture source across Africa (Schefuß et al., 2005; Tierney et al., 2008; Berke et al., 2012; Costa et al., 2014; Collins et al., 2013; Castañeda et al., 2016a). As for the arid Sahara, Tierney et al. (2017) document that temperature effects play a subordinate role while precipitation exerts predominant control on the  $\delta D_{wax}$ .

320 However, changes in the relative abundance of  $C_3$  versus  $C_4$  plants may also overprint the hydrological signal in  $\delta D_{wax}$  owing to different fractionation factors in the Calvin and Hatch-Slack photosynthetic pathways (e.g. Sachse et al., 2012; Collins et al., 2013). The  $\delta^{13}C$  signature of leaf wax lipids such as HMW  $n$ -alkanoic acids ( $\delta^{13}C_{wax\ n\text{-alkanoic acids}}$ ) is a common means to assess past changes in the relative contributions of  $C_3$  versus  $C_4$  plants in the catchment (Sachse et al., 2012; Collins et al., 2013) and to evaluate potential impacts of vegetation changes on  $\delta D_{wax}$  (e.g. Castañeda et al., 2016a; Collins et al., 2013). In order to explore potential effects of vegetation changes on our  $\delta D_{wax\ n\text{-alkanoic acid}}$  we correlate  $\delta D_{wax\ n\text{-alkanoic acid}}$  and  $\delta^{13}C_{wax\ n\text{-alkanoic acids}}$  (Figure S1). Considering that the linear correlation yields low correlation coefficients ( $R^2 < 0.5$ ; supplementary Figures S1 and S2) the impact of vegetation changes on  $\delta D_{wax\ n\text{-alkanoic acid}}$  is considered minor when the entire 18 ka are considered. Nevertheless, considerable impacts from vegetation changes may have existed during shorter intervals despite the weak correlation. The amplitudes in the  $\delta^{13}C_{wax\ n\text{-alkanoic acids}}$  are relatively small (up to 3‰) rendering profound impacts of vegetation changes on  $\delta D_{wax\ n\text{-alkanoic acid}}$  unlikely. When correcting  $\delta D_{wax\ n\text{-alkanoic acid}}$  for vegetation changes (supplementary material) the patterns and amplitudes persist (Figures S1, S3). We conclude that hydrologic variations exerted dominant control on  $\delta D_{wax\ n\text{-alkanoic acid}}$ . ~~Hence, we conclude and~~ that  $\delta D_{wax\ n\text{-alkanoic acid}}$  is a robust proxy for changes in rainfall and moisture source in the Nile River watershed throughout the past 18 ka. In order to deduce deglacial changes in hydroclimate in the Nile-River catchment, we pair our  $\delta D_{wax\ n\text{-alkanoic acid}}$  with the  $\delta D_{wax\ n\text{-alkanes}}$  from Castañeda et al. (2016a).  $\delta D_{wax\ n\text{-alkanes}}$  has been

335 interpreted to reflect rainfall variability in the Nile-River watershed associated with changes in the African summer monsoon (Castañeda et al., 2016a). The ~~fact~~ The offset in  $\delta D_{wax\ n\text{-alkanoic acid}}$  and  $\delta D_{wax\ n\text{-alkanes}}$  between 15-1 ka BP (Figure 4d) that  $\delta D_{wax\ n\text{-alkanoic acid}}$  and  $\delta D_{wax\ n\text{-alkanes}}$  show dissimilar patterns in in the long term trends suggests that the compounds recorded different hydrological developments which in turn implies that HMW  $n$ -alkanoic acids and HMW  $n$ -alkanes derive from different source areas. This conclusion seems reasonable considering that the Nile-River watershed extends over several climate zones and covers strong hydrological contrasts (tropical to subtropical and hyperarid desert). These zones are influenced by different atmospheric circulation patterns with the westerlies influencing the middle and lower catchments and the African monsoon

controlling precipitation in the headwaters. Hence, the paired application of  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  and  $\delta D_{\text{wax } n\text{-alkanes}}$  has the potential to reconstruct monsoonal along with westerly precipitation in the Nile River watershed. In order to understand the hydrologic signals recorded in  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  and  $\delta D_{\text{wax } n\text{-alkanes}}$  it is crucial to investigate-evaluate from which section of the watershed HMW *n*-alkanoic acids and HMW *n*-alkanes derive.

### 5.1 Sources areas of leaf-wax lipids in the Nile-River watershed

Leaf wax lipids reach the ocean mainly via fluvial discharge but also by aeolian transport (Schlünz and Schneider, 2000; Kusch et al., 2010; Schreuder et al., 2018). The latter is most important in open ocean settings where river discharge has little influence (Kusch et al., 2010; Hedges et al., 1997). Off large river systems leaf-waxes are predominantly introduced by river discharge (e.g. Kusch et al., 2010) and accordingly leaf-waxes in the sediments of the Nile deep sea fan (e.g. Blanchet et al., 2014) and at site GeoB7702-2 (Castañeda et al., 2016a) have been interpreted as recorders of environmental change in the Nile River watershed. However, in sediments of the southeastern Levantine basin the proportion of Saharan dust rises with increasing distance to the Nile delta (e.g. Krom et al., 1999). In core GeoB7702-3 the abundance profile of HMW *n*-alkanoic acids resembles the  $\delta^{18}\text{O}$  record of the planktonic foraminifera species *Globigerinoides ruber alba* ( $\delta^{18}\text{O}_{G. ruber alba}$ ; Figure 3c,d) from a sediment core from the Nile deep sea fan (MS27PT, Figure 1) which tracks freshwater discharge from the Nile River (Revel et al., 2010). Concentrations and river runoff are relatively low during the LGM and the early deglaciation and increase between 14-10 ka BP. During the middle Holocene both decline (Figure 3c,d). Similarly, HMW *n*-alkane concentrations rise between 14-10 ka BP and drop afterwards. This confirms that fluvial discharge was mainly responsible for the delivery of HMW *n*-alkanoic acids and HMW *n*-alkanes to the eastern Mediterranean. However, the abundances of HMW *n*-alkanes also have similarities with the proportion of Saharan dust in sediments of the southeastern Levantine Basin, in particular during the early deglaciation, before ~14 ka BP (Figure 3a, b). Thus, dust-bound HMW *n*-alkanes were additionally deposited at site GeoB7702-3 at these times. Considerable Saharan dust contributions are documented until 10 ka BP (Hamann et al., 2008; Figure 3a). Accordingly, aeolian transport may have been relevant to HMW *n*-alkanes from the last glacial until the early Holocene, possible dust sources to the southeastern Levantine Basin are the eastern Lybian Desert, central Algeria, Egypt and Sudan (Hamann et al., 2008). These regions are scarcely vegetated due to the vast extension of hyperarid desert. Therefore, dust bound HMW *n*-alkanes from these distal regions probably constitute a minor part in the HMW *n*-alkane pool at site GeoB7702-3. Thus, environmental change in the Nile river watershed predominantly accounts for the signal recorded by the HMW *n*-alkanoic acids and HMW *n*-alkanes.


Leaf-wax based records from off river mouths often are interpreted as catchment integrating signals (e.g. Blanchet et al., 2014; Häggi et al., 2016; Hemmingway et al., 2016), as also done by Castañeda et al. (2016a) with respect to  $\delta D_{\text{wax } n\text{-alkanes}}$  in core GeoB7702-3. However, several studies indicate that this does not generally hold true for all types of lipids and not for all river systems (Hemmingway et al., 2016; Agrawal et al., 2014; Galy et al., 2011). It has been reported that leaf-wax lipids may be representative only for specific parts of a river catchment and source regions may vary between compound-classes and even

375 between homologues within a compound class (e.g. Hemmingway et al., 2016; Agrawal et al., 2014). Hemmingway et al. (2016) found that HMW *n*-alkanoic acids originate from a local source in the Congo River watershed while HMW *n*-alkanes serve as catchment-wide integrator. The differences between  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  and  $\delta D_{\text{wax } n\text{-alkanes}}$  found in GeoB7702-3 strongly imply that the two lipid types derive from different source areas ~~in the Nile River watershed~~, at least for times when records of  $\delta D_{\text{wax } n\text{-alkanoic acids}}$  and  $\delta D_{\text{wax } n\text{-alkanes}}$  diverge (i.e. between 15-1 ka BP; Figure 4d). Similar  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  and  $\delta D_{\text{wax } n\text{-alkanes}}$  are found before 15 ka BP and from 1 ka BP onwards (Figure 4d) suggesting that the compounds derive from the same area at these times. Since dust input from the Sahara was relatively stable between 14.6-10 ka BP (Figure 3a; Hamann et al., 2008), it is unlikely that aeolian contributions from distal source regions outside the Nile river watershed account for the divergence in the  $\delta D_{\text{wax}}$  records. Instead, the offset most likely stems from different source areas within the Nile River watershed. ~~Similar  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  and  $\delta D_{\text{wax } n\text{-alkanes}}$  are found before 15 ka BP and from 1 ka BP onwards (Figure 2d) suggesting that the~~ compounds derive from the same area at these times. In order to identify the source areas of HMW *n*-alkanes and HMW *n*-alkanoic acids in GeoB7702-3 at present, we convert the  $\delta D_{\text{wax}}$  values into  $\delta D$  values of precipitation corrected for [changes in](#)

380

385

vegetation and ice volume change ( $\delta D_{p-vc-ic}$ ) according to the methods described in the supplementary material.



390 **Figure 32:** (a)–(c)  $\delta D_{wax}$  records from Lakes Tana (Costa et al., 2014), Victoria (Berke et al., 2012) and Tanganyika (Tierney et al., 2008) reporting hydroclimate variability in tropical East Africa; (d)  $\delta D_{wax-n-alkanoic\ acids}$  (orange:  $n-C_{26:0}$ ; green:  $n-C_{28:0}$ ; this study) along with  $\delta D_{wax-n-alkane}$  (dark green:  $n-C_{31}$ ; Castañeda et al., 2016a; 2016b) from core GeoB7702-3; (e) Ratio of tetra and penta-

395 methylated brGDGTs from the Nile deep-sea fan (core MS27PT; Figure 1) reporting on the input of delta-derived organic matter versus material from the soils in the upper Nile River catchment to the Eastern Mediterranean (Ménot et al., 2020); (f) oxygen isotopic composition of the planktic foraminifera species *Globigerinoides ruber alba* from the Levantine Basin (core MS27PT; Figure 1) used to reconstruct changes in salinity related to Nile River runoff (Revel et al., 2010; 2015). The light grey and dark grey shadings mark the episodes of the AHP and the Green Sahara, respectively. Dashed lines indicate the optimum of the AHP.

We compare the near core-top values (here 15cm-~~bsf~~ translating into 0.31 ka BP) to mean weighted  $\delta D$  values of precipitation of the growing season ( $\delta D_{p-gs}$ ) in the Nile catchment (Bowen et al., 2005). Bowen et al. (2005) define the growing season as 400 months with mean temperatures  $>0^{\circ}\text{C}$ . According to this definition, the values reported for the Nile River watershed are monthly weighted annual means since mean temperatures never drop below  $0^{\circ}\text{C}$  (<https://en.climate-data.org>).  $\delta D_{p-gs}$  values from different parts of the catchment and the results from our  $\delta D_{p-vc-ic}$  are listed in Table 2. The results downcore are given in Figure S3. For the core top  $\delta D_{p-vc-ic}$  is around  $-14\text{‰}$  for the  $n\text{-C}_{31}$  alkane and is  $-11\text{‰}$  and  $-8\text{‰}$  for the  $n\text{-C}_{26,0}$  and  $n\text{-C}_{28,0}$  alkanolic acids, respectively. This matches the isotopic composition of the Nile Delta where the predicted  $\delta D_{p-gs}$  range between 405  $-15$  and  $-11\text{‰}$  (at Cairo,  $30^{\circ}3'N$  and Alexandria,  $31^{\circ}13'N$ ; Bowen et al., 2005). Today, the Nile-Delta region receives the most deuterium-depleted precipitation in the entire watershed (Table 2; Bowen et al., 2005). In the middle section of the catchment, along the main Nile ( $30^{\circ}N\text{-}15^{\circ}N$ ),  $\delta D_{p-gs}$  becomes progressively more enriched towards the South ( $-4\text{‰}$  at Assuan,  $24^{\circ}6'N$  and  $10\text{‰}$  in Khartoum,  $15^{\circ}35'N$ ). In the headwater region, precipitation is more depleted compared to the Main-Nile but still is enriched by  $10\text{-}17\text{‰}$  ( $3.5\text{‰}$  Lake Tana,  $12^{\circ}N$  and  $-5.6\text{‰}$  Lake Victoria,  $1^{\circ}S$ ; Table 2) relative to  $\delta D_{p-gs}$  in the delta 410 (Bowen et al., 2005). As such, we infer that HMW  $n$ -alkanoic acids and HMW  $n$ -alkanes predominantly derive from the delta region at present.

**Table 2: Results for  $\delta D_{p-vc-ic}$  calculated based on  $\delta D_{wax}$  for GeoB7702-3 at 15 cm depth ( $\sim 0.3$  ka BP) together with  $\delta D_{p-gs}$  values across the Nile River watershed. Locations listed are indicated in Figure 1.**

Location	$\delta D_{p-vc-ic}$ [‰]	$\delta D_{p-gs}^1$ [‰]
GeoB7702-3; 15cm; $C_{31}$ $n$ -alkane	$-14.87 \pm 4.95$	
GeoB7702-3; 15cm; $C_{26,0}$ $n$ -alkanoic acid	$-11.10 \pm 3.14$	
GeoB7702-3; 15cm; $C_{28,0}$ $n$ -alkanoic acid	$-8.78 \pm 2.51$	
Cairo/Nile Delta		$-14.6$ to $-11.4$
Alexandria/Nile Delta		$-13.1$
Assuan/Main Nile		$-4.4$
Dunqula/Main Nile		$4.5$ to $5.3$
Khartoum/Main Nile		$9.9$
Kassala/Atbara		$13.4$ to $9.7$
Bahir Dar/Lake Tana		$3.5$
Juba/White Nile		$5.9$
Kampala /Lake Victoria		$-5.6$

<sup>1</sup>: adopted from Bowen et al. (2005).



415 When comparing  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  and  $\delta D_{\text{wax } n\text{-alkanes}}$  from GeoB7702-3 with  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  from Lakes in the headwater region and equatorial Africa (Lakes Tana, Victoria and Tanganyika (Figure 1); Berke et al., 2012; Tierney et al., 2008; Costa et al., 2014) we find a strong similarity between long term developments of  $\delta D_{\text{wax } n\text{-alkanes}}$  and the lake records during the past 18 ka (Figure 2a, b, c, d). This suggests that  $\delta D_{\text{wax } n\text{-alkanes}}$  records deglacial climate change in the headwaters of the river (Figure 2a, b, c, d) and that the source of the HMW  $n$ -alkanes must have changed at some point in the past. Accordingly, we propose

420 that the HMW  $n$ -alkanes received substantial contributions from the headwater region between ~15-1 ka BP, the time when  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  and  $\delta D_{\text{wax } n\text{-alkanes}}$  have dissimilar values. While the source of the HMW  $n$ -alkanes was dynamic during the past 18 ka, the source of the HMW  $n$ -alkanoic acids most likely was rather stable considering the dissimilar patterns in  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  from core GeoB7702-3 and the records from the lakes. The offset between the  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  and  $\delta D_{\text{wax } n\text{-alkanes}}$  between 15-1 ka BP indicates that either the source of the HMW  $n$ -alkanoic acid or the HMW  $n$ -alkanes changed during this

425 timeframe. Considering that the northeastern African hydroclimate saw drastic changes in the course of the AHP (Shanahan et al., 2015; Berke et al., 2012; Tierney et al., 2008; Costa et al., 2014) it is very likely that the gradients in  $\delta D_{\text{p-gs}}$  changed, too. As such, the present-day distribution of  $\delta D_{\text{p-gs}}$  cannot be used for paleo source-apportionment. Unfortunately, the detailed spatial distribution of past  $\delta D_{\text{p-gs}}$  in the Nile River watershed remains unresolved preventing us from using  $\delta D_{\text{p-vc-ic}}$  to identify the source areas of the leaf-waxes during the last glacial and the deglaciation. Therefore, we compare our  $\delta D_{\text{wax}}$  records to

430 available  $\delta D_{\text{wax}}$  data from Lakes Tana, Victoria and Tanganyika (Figure 1) and to a  $\delta D_{\text{wax}}$  record from the Dead Sea representing hydroclimate variability in the headwaters, equatorial Africa and the Eastern Mediterranean realm (Figure 3a-d; Berke, et al., 2012; Costa et al., 2014; Tierney et al., 2008; 2022).

Our  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  has a remarkable resemblance with the  $\delta D_{\text{wax}}$  record from the Dead Sea (Tierney et al., 2022; Figures 1, 4d). Plotted on the same scale the two records track each other sharing trends and amplitudes (Figure 4d). Even the range of

435 values covered by the two records is almost the same varying between ca. -105 to -145 ‰ during the past 18 ka. These similarities suggest that the  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  in core GeoB7702-3 was determined by Mediterranean hydroclimate throughout the past 18 ka. **This also implies that the HMW  $n$ -alkanoic acids were constantly sourced from the lower Nile River catchment.** Due to the larger amplitude in the long-term development of the  $\delta D_{\text{wax } n\text{-alkanes}}$  the range of values covered (-105 to -170 ‰) is larger than in the  $\delta D_{\text{wax } n\text{-alkanoic acid}}$ . If the Dead-Sea  $\delta D_{\text{wax}}$  and our  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  reflects the composition of the Mediterranean

440 moisture (Tierney et al., 2022) the  $\delta D_{\text{wax } n\text{-alkanes}}$  must have been influenced by more depleted rainfall, and thus by another moisture source, given the offsets between the Dead Sea record and the  $\delta D_{\text{wax } n\text{-alkanes}}$  in core GeoB7702-3 (Figure 4d). At Lake Tana  $\delta D_{\text{wax}}$  values of up to -160 ‰ have been recorded during the AHP providing evidence that in the headwaters of the Blue Nile the rainfall was more depleted during the AHP than the Mediterranean precipitation (Figure 4a). It is most likely, that the HMW  $n$ -alkanes recorded climate variability in the southern reaches of the river where the summer monsoon determines the

445 hydroclimate. In equatorial Africa decreasing  $\delta D_{\text{wax}}$  in sediments from Lake Tanganyika (Tierney et al., 2008) point to increasing humidity from 15 to 11.5 ka BP (Figure 4c) onwards. Likewise, the levels from Lakes Tana and Victoria - which desiccated during the late glacial (Stager et al., 2011; Lamb et al., 2007) – rose at that time in response to a wetter climate (Williams et al., 2006; Marshall et al., 2011). The **progressive decrease in  $\delta D_{\text{wax } n\text{-alkanes}}$  starting at 15 ka BP thus is in harmony**

with the hydroclimate development in the headwaters during the early AHP which further corroborates our inference that the  $\delta D_{\text{wax } n\text{-alkanes}}$  was substantially influenced by rainfall in the southern Nile River catchment. Accordingly, we propose that the HMW *n*-alkanes in core GeoB7702-3 received considerable contributions from the headwater region between ~15-1 ka BP, the time when  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  and  $\delta D_{\text{wax } n\text{-alkanes}}$  deviate from each other. Before and afterwards both compounds derived predominantly from the delta. While the source of the HMW *n*-alkanes was dynamic during the past 18 ka, the source of the HMW *n*-alkanoic acids most likely was rather stable, ~~given the similarities in  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  from core GeoB7702-3 with the  $\delta D_{\text{wax}}$  from the Dead Sea (Figure 2a3a, b, c, d).~~ Considerable contributions of HMW *n*-alkanoic acids from the headwater region are unlikely. However, acknowledging that the Sahara was vegetated between ~11-5 ka BP (Kuper and Kröpelin, 2006; Watrin et al., 2009; Hély et al., 2014; Hamdan et al., 2016) the source region of the HMW *n*-alkanoic acids may have extended further south into the northern Sahara at this time. Likewise, the HMW *n*-alkanes probably received additional contributions material from the middle catchment of the river. During the AHP, site GeoB7702-3 may also have received additional material load from the Sinai Peninsula through Wadi El Arish. This drainage basin is rather dry at present but activated during the AHP (AbuBakr et al., 2013; Muhs et al., 2013).

Our scenario regarding dynamics in HMW *n*-alkane provenance fits to findings from Ménot et al. (2020) who analyzed branched Glycerol Dialkyl Glycerol Tetraethers (brGDGTs) in the Nile deep sea fan in order to reconstruct soil input from the Nile River into the EM over the deglaciation. BrGDGTs are synthesized by bacteria thriving in peat and soils (Weijers et al., 2006, Martin et al., 2019) but can also be produced in-situ in rivers and estuarine settings and in the marine sediments (deJonge et al., 2014). Using the ratio between tetra- and penta- methylated brGDGTs ( $\Sigma\text{IIIa}/\Sigma\text{IIa}$ -ratio) Ménot et al. (2020) inferred that the compounds derived from the Nile delta between 20-14.5 ka BP and after 4 ka BP (Figure 2e3e). The interval of soil-derived input (14.5-4 ka BP; Ménot et al., 2020) matches the interval of diverging  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  and  $\delta D_{\text{wax } n\text{-alkanes}}$  (Figure 42d) during which considerable amounts of HMW *n*-alkanes from the upper and middle catchment were deposited in the EM sediments.

Being hydrophobic the HMW *n*-alkanes and HMW *n*-alkanoic acids are among the recalcitrant fraction of organic matter that is preserved for long times in the geological record and consequently is able to survive riverine transport and intermediate storage in reservoirs prior to burial in the marine sediments (e.g. Eglinton and Eglinton, 2008). However, our north-south allocation regarding the sources of HMW *n*-alkanoic acids and HMW *n*-alkanoic acids in GeoB7702-3 suggests that in the Nile-River watershed, the most of the HMW *n*-alkanoic acids sourced from the upper catchment did not reach site GeoB7702-3, most likely due to degradation during riverine transport. This is in accordance with findings from Hemmingway et al. (2016) who inferred that in the Congo-River catchment HMW *n*-alkanoic acids in suspended sediments from the outflow derive from local sources while HMW *n*-alkanes have a local and distal origin providing a more catchment integrated signal. Similarly, Agrawal et al. (2014) suggested that in the Ganga-Bahamaputra River system HMW *n*-alkanoic acids from the Himalayan headwaters degrade during transport and get replaced by HMW *n*-alkanoic acids from the local floodplains. There is consensus that HMW *n*-alkanoic acids are more prone towards degradation than HMW *n*-alkanes (Meyers and Ishiwatari, 1993; Hoefs et al., 2002; Sinninghe-Damsté et al., 2002; Galy and Eglinton, 2011; Hemmingway et al., 2016) which probably accounts for

the discrepancies in provenance of HMW *n*-alkanoic acids and HMW *n*-alkanes observed in these large river systems (e.g. Hemmingway et al., 2016) and at our study site.

## 485 5.2 Environmental drivers of HMW *n*-alkane provenance

There is compelling evidence that during the LGM and the early deglaciation, ~~and~~ in particular Heinrich Stadial 1 (HS1), the Northeastnortheastern African climate was very arid (Stager et al., 2011; Castañeda et al., 2016; Tierney et al., 2008; Tierney and deMenocal, 2013; Revel et al., 2014; Ménot et al., 2020). Lakes Tana and Victoria, the sources of the Blue and White Nile tributaries, desiccated between-at 17-16 ka BP (Stager et al., 2011; Lamb et al., 2007) leading to a drastic reduction in runoff in the Nile River system relative to present (Williams et al., 2009; 2015) as recorded- ~~documented~~ by higher  $\delta^{18}\text{O}$  ~~values of the planktonic foraminifera species Globigerinoides ruber alba~~ ( $\delta^{18}\text{O}_{G. ruber alba}$ ; (Figure 42f) offshore the Nile River mouth (Revel et al., 2010; 2015). At the same time deposition of Saharan dust peaked in the EM (Hamann et al., 2008). The weak fluvial activity in the Nile River likely restricted the source of the leaf-wax lipids to the Nile delta and led to minimal export of HMW *n*-alkanoic acids in eastern Mediterranean sediments (Figure 3c). Relatively high concentrations of HMW *n*-alkanes in core 495 GeoB7702-3 despite the diminished fluvial activity (Figure 3b) can be best explained by aeolian transport of HMW *n*-alkanes sourced from the vegetated Nile delta region and floodplain deposits of the lower Nile River catchment considering that the input of Saharan dust to the eastern Mediterranean peaked at these times (Hamann et al., 2008; Figure 3a).  $-\delta\text{D}_{\text{wax}}$  records from Lake Tanganyika (Figure 2e3c) document that the East African climate became wetter around 14.5-15 ka BP (Tierney et al., 2008). In response to the wetter conditions the overflow of Lakes Tana and Victoria resumed around 15.5 and 14.5 ka BP 500 (Williams et al., 2006; Marshall et al., 2011) and freshwater input from the Nile River increased accordingly as documented by lower  $\delta^{18}\text{O}_{G. ruber alba}$  (Revel et al., 2010; 2014; 2015). The climate amelioration in Northeastnortheastern Africa coincided ~~ed~~ with the onset of divergence between  $\delta\text{D}_{\text{wax } n\text{-alkanes}}$  and  $\delta\text{D}_{\text{wax } n\text{-alkanoic acid}}$  as well as with the switch from deltaic to soil-derived brGDGTs (Figure 24d,e). Concentrations of HMW *n*-alkanes increase along with decreasing  $\delta^{18}\text{O}_{G. ruber alba}$  (Figure 3c,d) suggesting that ~~e~~Enhanced fluvial energy in the headwaters probably increased erosion and export of organic matter in the upper reaches of the river as well as in the middle catchment where vegetation expanded into the Sahara in the course of the AHP. As mentioned above, a considerable amount of HMW *n*-alkanoic acids mobilized in the headwaters probably got degraded during riverine transport and did not reach the core site. Rising concentrations of HMW *n*-alkanoic acids (Figure 3c) thus reflect intensified fluvial erosion in the delta and floodplains of the lower Nile catchment. During the late Holocene, at around 5-4 ka BP, drier conditions re-established in the headwaters and tropical East Africa (Tierney et al., 2008, Costa et al., 510 2014, Berke et al., 2012). ~~and T~~he Nile runoff reduced accordingly (Revel et al., ~~2015~~2010; 2015~~9~~). Again, the delta became the predominant source of HMW *n*-alkanes and brGDGTs.

## 5.3 Hydroclimate development during the past 20 ka

As for the reconstruction of the hydroclimate variability our findings imply that the HMW *n*-alkanoic acids and the HMW *n*-alkanes can be used to reconstruct the hydroclimatic developments in the lower and upper reaches of the Nile River in parallel

515 (between ~14.5-4 ka BP). As elaborated earlier, the delta is predominantly influenced by Mediterranean winter precipitation. According to our inference that the HMW *n*-alkanoic acids consistently derive from the delta  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  should record changes in winter precipitation throughout the past 18 ka. By contrast the  $\delta D_{\text{wax } n\text{-alkanes}}$  should predominantly be a summer signal, as we infer that the HMW *n*-alkanes receive major contributions from the headwaters south of 15°N where the African summer monsoon controls precipitation. ~~Given that the Sahara became vegetated during the Early Holocene,  $\delta D_{\text{wax } n\text{-alkanes}}$  potentially also recorded changes in the middle catchment.~~ The application of paired  $\delta D_{\text{wax } n\text{-alkanes}}$  and  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  allows  
520 to investigate how westerly (winter) and monsoonal (summer) precipitation evolved around the AHP.

#### 5.4 early deglaciation – Heinrich Stadial 1 (18-14.6 ka BP)


The last glacial maximum (~21 ka BP; Clark et al., 2009) was a relatively moist interval in the southeastern Mediterranean realm due to a more positive water balance (precipitation – evaporation) compared to today as indicated by higher levels of glacial Lake Lisan (paleo Dead Sea; Figure 1; Stein, 2001) and some pollen data from Dead Sea sediments (Langgut et al., 2021). In the northeastern Sahara, tufa deposits from the Eastern Desert provide evidence for sufficiently moist conditions to recharge aquifers (Hamdan and Brook, 2015). As for rainfall amount it is debated whether it increased, remained similar or reduced compared to today (Ludwig and Hochmann, 2022; Kageyama et al., 2021; Goldsmith et al., 2017; Enzel et al., 2008). In case of reduced or similar rainfall amount, reduced evaporation must have balanced reductions in annual rainfall (Ludwig and Hochmann, 2022). It is assumed that cold glacial conditions likely resulted in reduced evaporation and higher effective moisture during the LGM (Ludwig and Hochmann, 2021; Stockhecke et al., 2016). Moreover, the polar front and the westerlies were displaced southward relative to today (situated at ca. 40°N; Wang et al., 2018) owing to the presence of the Fennoscandian ice sheet (Wang et al., 2018; Kageyama et al., 2021). Accordingly, storms frequently steered moist air from the Atlantic Ocean and Mediterranean Sea to the southeastern Mediterranean borderlands accounting for higher effective moisture and potentially also enhanced rainfall (Hamdan and Brook, 2015; Wang et al., 2018; Goldsmith et al., 2017; Kageyama et al., 2021; Enzel et al., 2008).  
525  
530  
535

~~Several authors pointed out that the LGM (21-19 ka BP) was a relatively moist interval in the southeastsoutheastern Mediterranean realm where conditions may have been similar to today (Wang et al., 2018). In the northeastnortheastern Sahara precipitation must have been sufficient to recharge aquifers from which tufa deposits formed in the Eastern Desert (Hamdan and Brook, 2015). During the LGM, the polar front and the westerlies were displaced to the south relative to today (situated at ca. 40°N; Wang et al., 2018) owing to the presence of the Fennoscandian ice sheet (Wang et al., 2018). Accordingly, storms frequently reached the Eastern Mediterranean and Northern Sahara providing the region with moisture (Hamdan and Brook, 2015; Wang et al., 2018). The Northeastnortheastern African climate became harsher during the early deglaciation (from 19 ka onwards) and the most severe aridity occurred during HS1 (~18-14.5 kaBP; Tierney et al., 2008; Stager et al., 2008; Hamdan and Brook, 2015). Lakes Victoria and Tana desiccated (Stager et al., 2011; Lamb et al., 2007) and in the eastern Sahara, tufa formation ceased (Hamdan and Brook, 2015). Also, in the Nile-River delta conditions may have rapidly become more arid at the beginning of HS1 given the abrupt increase of  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  as well as the  $\delta D_{\text{wax } n\text{-alkanes}}$  (Castañeda et al.,~~  
540  
545

2016) at 17.8 ka BP (Figures 2d4d, 3d5d). Maximal values of  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  and  $\delta D_{\text{wax } n\text{-alkanes}}$  suggest that also in the Nile River delta experienced the most arid conditions since the LGM at that time. However, the drought was restricted to the first half of HS1. ~~And~~ at ~16 ka BP, the climate rapidly transitioned into a wet phase that lasted until the end of HS1 (14.6 ka BP) (Figures 2d4d, 3d5d). Also,  $\delta^{18}\text{O}$  of speleothems from the Soreq Cave/Israel suggest a progressive increase in precipitation at the same time (Figure 3b5b).  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  show that hydroclimatic conditions may have been similar to the late Holocene and today considering the similar values in  $\delta D$  (-130 to -125‰). The arid episode attests to a decrease in precipitation and/or a decrease in evaporation. In turn, the reversal to more humid conditions was either caused by increased rainfall and/or reduced evaporation. The dry episode coincided with a moderate temperature rise as documented by TEX<sub>86</sub>-based sea surface temperature (SST) reconstructions from core GeoB7702-3 (Castañeda et al., 2010a; Figure 5e). By contrast, the wet period occurred along with a cold phase in the EM eastern Mediterranean (Figure 5e). This suggest that early warming may have reduced moisture availability in the region by enhancing evaporation. Later, the reversal to colder conditions may have reduced evapotranspiration allowing for more effective moisture. Next to temperature-induced changes in evaporation, variations in rainfall amount due to altered atmospheric circulation patterns are potential drivers that need to be evaluated. As noted by Bartov et al. (2003) lower air temperatures that are possibly came along associated with low SSTs in the EM during HS1 (Bar-Mathews et al., 1999), would have minimized the cyclonic activity over the EM. A reduction of the intensity and frequency of Cyprus Lows would have reduced the advection of moist air and rainfall in the eastern Mediterranean realm fostering aridity. However, enhanced aridity is at odds with the isotopic depletion of the leaf waxes documented in core GeoB7702-3. If the local cyclonic activity was weakened, either effects of reduced evaporation must have balanced the reduced advection of moisture from the Cyprus Lows or another moisture source must have provided the region with rainfall. Interestingly, Italian speleothems imply that the delivery of Atlantic moisture to the central Mediterranean intensified from 16 ka BP onwards (Columbu et al., 2022). Analogous to the Italian site, a stronger influence of the Atlantic moisture may have caused more humid conditions in the southeastern Mediterranean. The succession of dry and wet episodes within HS1 recorded at our study site fits the growing view that HS1 evolved in two phases, the first one lasting from ~18.2- 16.2 ka and the second lasting from 16.2-14.5 ka BP (e.g. Naughton et al., 2023 and references within). Two-phase patterns with alternating dry and wet intervals have also been recorded by many marine and terrestrial archives in the entire Mediterranean realm (Valsecchi et al., 2012; Naughton et al., 2023; Naughton et al., 2009; 2016; Pérez-Mejías et al., 2021; Fletcher and Sanchez-Goñi et al., 2008) and even in western North America where they have been called “big dry” and “big wet” intervals (Broecker et al., 2009). However, the Mediterranean records are inconsistent regarding the order of wet and dry episodes indicating a complex hydroclimate development in the Mediterranean borderlands throughout HS1. For example, the Sea of Marmara/Northeast/northeastern/northern Mediterranean experienced wet conditions during the first phase of HS1 followed by drought during the second phase (Valsecchi et al., 2012). The opposite pattern is evident in our data at site GeoB7702-3. These regional differences are probably related to spatial variations of the Atlantic-Mediterranean storm track. Proxy-based and modeling studies suggest that the deglacial Mediterranean climate responded to abrupt climate fluctuations in the North Atlantic via atmospheric teleconnections involving the position of the westerly Jet-jet and the associated storm track (Columbu et al., 2022; Valsecchi et al., 2012; Li et

al., 2019). According to Li et al. (2019) a weakening of the Atlantic Meridional Overturning Circulation (AMOC) during HS1 would push the westerlies northward. ~~Considering that during the LGM, the polar front and the westerlies were situated at ca. 40°N; (Wang et al., 2018) a~~ northward movement of the westerlies and the ~~Atlantic~~ storm track during the first half of HS1 ~~would~~could explain why the climate became drier in the Nile River delta but concurrently wetter in the ~~NE~~northeastern Mediterranean. A subsequent return to the South at ~16 ka BP would have reversed the situation with wet conditions in the Nile-River delta and concurrent aridity in the NE Mediterranean realm. Our inference agrees with findings from the western Mediterranean basin since Naughton et al. (2023) conclude that opposing successions of dry and wet phases on the Iberian Peninsula during HS1 document a northward followed by a southward movement of the polar front and the associated westerly ~~Jet~~jet.








Figure 343: (a) Oxygen isotope records from speleothems ( $\delta^{18}\text{O}_{\text{carbonate}}$ ) in Jeita Cave, Lebanon (Cheng et al., 2015); (b) Oxygen isotope records from speleothems in Soreq Cave, Israel (Bar-Mathews et al., 2003); (c)  $\delta\text{D}_{\text{wax}}$  record based on the  $n\text{-C}_{30:0}$  alkanolic acid in core DSDDP 5017-1 from the Dead Sea (Figure 1; Tierney et al., 2022) (de) The red bar marks the interval during which spring-tufa deposits formed on the Sinai Peninsula (Hamdan and Brook, 2015); (e)  $\delta\text{D}_{\text{wax}}$   $n$ -alkanoic acids (orange:  $n\text{-C}_{26:0}$ ; green:  $n\text{-C}_{28:0}$ ; this study) along with  $\delta\text{D}_{\text{wax}}$   $n$ -alkane (dark green:  $n\text{-C}_{31}$ ; Castañeda et al., 2016a; 2016b) from core GeoB7702-3; (f) sea surface temperature (SST) in the Eastern Mediterranean (EM) based on TEX<sub>86</sub> from core GeoB7702-3 (Castañeda et al., 2010a; 2010c; 2016a); (g) Insolation during January and July at 15°N from Berger and Loutre (1991). The light grey and dark grey shadings indicate the episodes of the AHP and the Green Sahara and their optimum, respectively. The AHP optimum is marked by dashed lines. LGM: Last Glacial Maximum; HS1: Heinrich Stadial 1; B/A: Bølling-Allerød interstadial; YD: Younger Dryas stadial.

At the end of HS1, relatively dry conditions established in the Nile-Delta region ~~which suggests that the storm track moved north again. Relatively dry conditions and~~ prevailed ~~in the Nile River delta~~ until 11.5 ka BP as documented by increased  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  (Figure ~~3d5d~~). ~~indicating that~~ The storm frequency and westerly moisture supply must have remained weak in the region throughout this period. ~~Climate models propose that the polar front and the westerly jet moved northwards once the Fennoscandian Ice Sheet retreated (Wang et al., 2018; Kageyama et al., 2021). This northward movement may have diminished westerly-moisture supply to the Nile delta after 15 ka BP. Additionally, warming after HS1 (Figure 5e) may have intensified evapotranspiration fostering aridity in the region. This contradicts records from the western, central and the NE Mediterranean region (e.g. Jeita Cave/Lebanon, Figure 3a) where wet episodes already occurred earlier coincident with the Bølling/Allerød (B/A) interstadial in the North Atlantic (Fletcher et al., 2010; Columbu et al., 2022; Valsecchi et al., 2012; Cheng et al., 2015). These are attributed to enhanced moisture delivery by stronger westerlies in the region (e.g. Columbu et al., 2022; Li et al., 2019). Also, subsequent reversals to drier conditions during the Younger Dryas (YD) have been identified (Fletcher et al., 2010; Columbu et al., 2022; Valsecchi et al., 2012; Cheng et al., 2015). These fluctuations are interpreted to document that the teleconnection between north Atlantic climate dynamics and Mediterranean rainfall via the westerly Jet mediated rainfall in the western, central and northeastern Mediterranean throughout the deglaciation (Columbu et al., 2022). The absence of such variations in the Nile River delta suggests that Atlantic climate forcing was suppressed in the northeastern Sahara. Once the Fennoscandian Ice Sheet retreated the polar front and the westerly Jet moved northwards (Wang et al., 2018) and probably pushed the trajectory of storms northwards accordingly leading to dry conditions in Nile River delta throughout the B/A and YD. Consequently, dynamics of the westerlies associated with AMOC variations probably did not influence the Nile River delta anymore. Considering that Jeita Cave recorded a wet interval during the B/A (Cheng et al., 2015; Figure 3a), the southern boundary of the storm track was probably situated between  $\sim 31\text{--}33^\circ\text{N}$  at this time.~~

### 5.5 African Humid Period (14.5-5 ka BP)

Whereas  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  indicate relatively constant dryness after HS1,  $\delta D_{\text{wax } n\text{-alkanes}}$  document a progressive depletion of plant waxes. As previously elaborated by Castañeda et al. (2016a), this progressive depletion starting at  $\sim 15$  ka BP most likely stems from increasing rainfall amount in the headwaters of the Nile-River catchment. This development agrees with increasing humid conditions at Lake Tanganyika, Tana and Victoria and has been interpreted as intensification of the African summer monsoon (Castañeda et al., 2016, Berke et al., 2012; Tierney et al., 2008; Costa et al., 2014). This hydroclimate amelioration led into in the AHP which lasted from  $\sim 14.5\text{--}5$  ka BP and culminated between  $\sim 11\text{--}6$  ka BP (e.g. deMenocal et al., 2006a; Tierney et al., 2017; Dupont and Schefuß, 2018). More negative  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  indicate that also the Nile Delta received substantially more rainfall during the AHP optimum, i.e. between 11.0-6.0 ka BP (Figure ~~2d4d~~). A case study conducted by Breitenbach et al., (2010) in the Bay of Bengal/Indian Ocean shows that intensified river discharge due to heavy summer-monsoon rainfall leads to surface water freshening in the Ocean which in turn lowers the  $\delta D$ -signature of precipitation derived from these surface waters. Likewise, the discharge of isotopically light water from the Nile River watershed varied along with

635 [monsoon intensity during the Late Quaternary \(Almogi-Labin et al., 2009\). Isotopic variations in the Mediterranean surface waters have also been invoked to explain parts of the  \$\delta^{18}\text{O}\$ -signals recorded in speleothems from caves in Israel and the Lebanon \(Cheng et al., 2012\) and  \$\delta\text{D}\_{\text{wax}}\$  from the Dead Sea \(Tierney et al., 2022\).](#) Considering that the  $\delta^{18}\text{O}_{G. ruber\ alba}$  records from the Mediterranean Sea (Emeis et al., 2000; Revel et al., 2010; 2015) show massive surface freshening during the AHP [resulting from enhanced monsoon rains in the Nile-River watershed](#), this effect could have potentially accounted for depleted

640  $\delta\text{D}_{\text{wax } n\text{-alkanoic acid}}$  in addition to or instead of the amount effect. However, given that the  $\delta^{18}\text{O}_{G. ruber\ alba}$  (Revel et al., 2010; 2015) and  $\delta\text{D}_{\text{wax } n\text{-alkanoic acid}}$  show different temporal trends (Figure [2d4d,f](#)) freshening of the Mediterranean probably constitutes only a minor part of the signal in  $\delta\text{D}_{\text{wax } n\text{-alkanoic acid}}$ . Therefore, the minimum in  $\delta\text{D}_{\text{wax } n\text{-alkanoic acid}}$  most likely attests to drastically increased rainfall amount in the Nile River delta during the AHP. [Interestingly, Tierney et al. \(2022\) found that the glacial-interglacial hydroclimate is quite sensitive to temperature driven changes in the atmospheric evaporative demand. Warming enhances evapotranspiration and reduces effective moisture.  \$\text{TEX}\_{86}\$ -based SST reconstructions at site GeoB7702-3 show pronounced warming at 11-10 ka BP \(Figure 5e; Castañeda et al., 2010a\). Consequently, a deteriorated water balance due to elevated evapotranspiration would be expected for the Levant. However, an increase in evapotranspiration and reduction in effective moisture would cause an isotopic enrichment of the leaf waxes rather than a depletion \(Sachse et al., 2012\). As such, the decrease in  \$\delta\text{D}\_{\text{wax } n\text{-alkanoic acid}}\$  reported at site GeoB7702-3 clearly documents that increases in rainfall must have](#)

645 [been exceeded any potential increases in evaporation leading to a more positive water balance. It is well constrained that the ITCZ shifted northward during the AHP supplying the Sahara with regular rainfall \(e.g. deMenocal et al., 2006b; Menviel et al., 2021; Braconnot et al., 2007; Blanchet et al., 2021; Tierney et al., 2017; Kuper and Kröpelin, 2006; Hamdan and Brook, 2015\). Some studies suggest that the monsoon fringe even expanded up to 31°N \(Sha et al., 2019; Tierney et al., 2017\) which would mean that the Nile River delta \(situated at 30-31°N; Figure 1\) became influenced by the African summer monsoon. However,](#)

650 [in most climate simulations the northernmost position of the ITCZ is located at ~24°N \(e.g. Pausata et al., 2016\) which is also corroborated by proxy data \(Hamdan and Brook, 2015; Cheddadi et al., 2021\). Thus, we consider it unlikely that the northward migration of the monsoon rainbelt caused the increased rainfall in the delta region. Instead, increased winter precipitation associated with the westerlies most likely was responsible for the minimum in  \$\delta\text{D}\_{\text{wax } n\text{-alkanoic acid}}\$ .](#) [It is well constrained that the ITCZ and summer monsoon shifted northward during the AHP supplying the Sahara with seasonal rainfall \(e.g. deMenocal et al., 2006b; Menviel et al., 2021; Braconnot et al., 2007; Blanchet et al., 2021; Tierney et al., 2017; Kuper and Kröpelin, 2006; Hamdan and Brook, 2015\). However, the northernmost extent of the monsoon is debated. Today the northernmost limit is at 14°N. In many climate simulations for the AHP the monsoon reaches ~24°N \(e.g. Pausata et al., 2016; Chandan and Peltier, 2020; Thompson et al., 2021; Hopcroft et al., 2017; Dallmeyer et al., 2020\). This is a more modest extent compared to pollen-based reconstructions which indicate the limit at 23-28°N \(Bartlein et al., 2011; Hély et al., 2014\). However, some climate](#)

655 [models acknowledging several land-surface feedbacks set the northernmost boundary as far north as 30-31°N over northwestern Africa \(Pausata et al., 2016\). Likewise, analysing isotopic compositions of speleothems and leaf-wax lipids researchers concluded that the monsoon fringe expanded up to 27-31°N over northwestern Africa \(Sha et al., 2019; Tierney et al., 2017\), yet this interpretation recently was challenged by pollen data from northern Morocco \(Cheddadi et al., 2021\).](#) [A](#)

670 ~~similar northward extension over northeastern Africa would mean that the Nile River delta (situated at 30-31°N; Figure 1) became influenced by the African summer monsoon. However, while As for northeastern Africa climate models consistently compute very small positive anomalies or no change (e.g. Perez-Sanz et al., 2014; Dallmeyer et al., 2020; Pausata et al., 2016) even when simulate wet conditions reach as far north as 30-31°N over for the northwestern Sahara (Pausata et al., 2016)-. Records of Saharan Dust input to the North Atlantic and the Red Sea also point to a greater extent of the monsoonal rains in the northwestern than in the northeastern Sahara (Palchan and Torfstein, 2019; McGee et al., 2013). The wet dry west-east gradient in humidity has been attributed to the tilted axis of the monsoonal rainbelt (Pausata et al., 2016; Dallmeyer et al., 2020). From the climate model perspective monsoonal rains did not reach the Nile delta region during the AHP. This view is consistent with dust accumulation rates in the Red Sea which set the limit of monsoonal rains at 22°N over northeastern Africa (Palchan and Torfstein, 2019). Further support comes from Holocene spring tufa deposits in the Eastern Desert and the Sinai Peninsula (Hamdan and Brook, 2015). The tufas from the Eastern Desert (~25°N) were fed from monsoonal precipitation~~  
675 ~~Holocene Tufa deposits from the Eastern Desert (Egypt; Figure 1) provide evidence for monsoonal rains at ~25°N whereas spring tufa from the Sinai Peninsula (at ~27°N) were fed by precipitated from springs recharged by Mediterranean moisture rainfall (Hamdan and Brook, 2015) indicating that the monsoon limit over northeastern Africa was situated south of 27°N. Moreover, salinity reconstructions from the Red Sea (Arz et al., 2008) suggest that the Mediterranean was the predominant moisture source to north easternmost Africa during the Holocene. Collectively, models and proxies suggest that the monsoonal rainbelt never reached the Nile Delta region. Thus, we consider it unlikely that the northward migration of the monsoon rainbelt caused the increased rainfall in the delta region. Instead of monsoonal rain, increased winter precipitation associated with the westerlies most likely was responsible for the minimum in  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  between 11-6.5 ka BP.~~

### 5.5.1 Controls on enhanced winter precipitation

690 The rapid switch to wetter conditions at 11 ka BP as well as the abrupt return to dry conditions at 6.5 ka BP displayed by the  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  occurred congruently with a warm ~~spell-phase~~ in the EM between 10-7 ka BP documented by the TEX<sub>86</sub>-SST proxy from GeoB7702-3 (Castañeda et al., 2010a). The warm SST would have enhanced evaporation over the Mediterranean Sea and at the same time would have led to a higher water-holding capacity of the atmosphere in virtue of the Clausius-Clapeyron relation (e.g. Trenberth et al., 2003; Lenderink and Meijgaard, 2008). Additionally, warming would have  
695 ~~strengthened local cyclogenesis the formation of Cyprus Lows-~~ over the Levantine Basin. Both effects would have increased precipitation moisture delivery to over the Nile-delta region. Records from the Middle East (Figure ~~3a4a~~,b), southern Europe and northwest Africa provide evidence that mild and wet conditions were widespread across the Mediterranean borderlands during the early Holocene (Cheddadi et al., 2021, Wagner et al., 2019, Bar-Mathews et al., 2003; Cheng et al., 2015). Today, relatively mild and wet winters in the Mediterranean realm are associated with negative phases of the NAO-Index during which  
700 the westerlies weaken and the Atlantic storm track is situated south steering Atlantic moisture and storms frequently into the Mediterranean (Figure 1). As invoked by several data- as well as model-based studies, negative NAO-like circulation patterns

probably prevailed during the early Holocene promoting intense winter precipitation in the Mediterranean borderlands (Kutzbach et al., 2014, Arz et al., 2003, Dixit et al., 2020, Wassenburg et al., 2016). Previous work suggests that the southward displacement of the westerlies and the associated storm trajectory was a response to decreasing Northern Hemisphere winter insolation and thus a result of precessional forcing (Kutzbach et al., 2014; Wagner et al., 2019; Li et al., 2019). Analyzing lacustrine sediments from Lake Ohrid (Balkan Peninsula) Wagner et al. (2019) proposed that the Mediterranean winter precipitation varied in-phase with the African summer monsoon strength over glacial-interglacial cycles as both were driven by precessional and thus insolation forcing. The monsoon responded to rising summer insolation changes and Mediterranean precipitation to weakened winter insolation (Wagner et al., 2019; Cheng et al., 2016). Indeed,  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  and  $\delta D_{\text{wax } n\text{-alkanes}}$  from core GeoB7702-3 confirm that Mediterranean winter rains and the African summer monsoon (Castañeda et al., 2016a; Costa et al., 2014, Berke et al., 2012, Tierney et al., 2008) were concurrently strong in the Nile River watershed during the AHP optimum but wet conditions began later (~11 ka BP) and also ended earlier (~6 ka BP) in the delta than in the upper river catchment (~14.5-5 ka BP; Figures 2a4a-d, 53d). The relatively short wet phase recorded in our  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  agrees with pronounced freshening of the northern Red Sea which has been interpreted to result from enhanced winter rainfall over Egypt and the Sinai Peninsula (Arz et al., 2003). Moreover, spring tufa deposits formed on the Sinai Peninsula only between 9.0-6.7 ka BP (Figure 3e5c) and were fed by Mediterranean moisture (Hamdan and Brook, 2015). Together these data show that in the northeastern Sahara wet conditions were restricted to the AHP optimum (11.0-6.5 ka BP) ~~although winter insolation had already begun to decrease 7 ka earlier (Figure 3d,f)~~. Likely, the storm track reached the northeastern Sahara only when the difference between summer insolation and winter insolation was maximal (Figures 3a5a,b,c,d). Interestingly, speleothem  $\delta^{18}\text{O}$ -records from caves in Lebanon (Jeita Cave; Figure 1) suggest that the hydroclimate may have ameliorated slightly earlier ~~than in the Nile Delta~~, i.e. around 12 ka BP (Figure 3a5a,d) while Soreq Cave ~~and  $\delta D_{\text{wax}}$  from the Dead Sea~~ indicates an increase of humid conditions at the same time as  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  in core GeoB7702-3 (Figure 3b5b,d). ~~Likewise, wet conditions lasted longer at Jeita Cave than in the Nile-River delta region (Figure 5a, d)~~. The staggered onset ~~and termination~~ of wet conditions in the south eastern Mediterranean realm may attest to the progressive southward ~~and subsequent northward~~ migration of the storm track ~~if the temporal offset does not stem from uncertainties in the chronologies~~.

### 5.5.2 Implications for the “Green Sahara” (11-5 ka BP)

~~Providing insights into the development of summer and winter precipitation in the Nile River watershed our findings are important for the understanding of the moisture sources which sustained the “Green Sahara”. As elaborated by Castaneda et al. (2016), the  $\delta D_{\text{wax } n\text{-alkane}}$  attest to enhanced monsoon rainfall in the Nile River watershed. It is widely accepted that the northward migration and intensification of the summer monsoon was critical to trigger plant growth in the Sahara (Braconnot et al., 2007; Menviel et al., 2021; deMenocal et al., 2000b; Shannahan et al., 2015; Sha et al., 2019; Tierney et al., 2017). However, discrepancies between proxy data and climate simulations regarding precipitation and vegetation anomalies question that the monsoon was the only moisture source to the Sahara (Chandan et al., 2020; Braconnot et al., 2007; Hoperoff et al., 2017; Perez Sanz et al., 2014; Cheddadi et al., 2021; Hély et al., 2014). Monsoonal precipitation may not have reached~~



735 beyond  $\sim 24^{\circ}\text{N}$  (Chandan and Peltier, 2020; Braconnot et al., 2007; Hoperoff et al., 2017; Perez Sanz et al., 2014) but vegetation  
extended up to  $28\text{--}31^{\circ}\text{N}$  (e.g. Hély et al., 2014; Tierney et al., 2017; Giraudi et al., 2013; Hamdan et al., 2016; Larrasoña et  
al., 2013). In the northern Sahara, south of the Nile Delta, the landscape turned into a semi-arid desert zone covered by  
740 vegetation patches comprising open Savannah with mainly grasslands. Wetlands and trees likely flourished near permanent  
water bodies (Hamdan et al., 2016; Larrasoña et al., 2013). Some tropical plants migrated north of  $25^{\circ}\text{N}$  but were restricted  
along rivers and lakes (Hamdan et al., 2016; Larrasoña et al., 2013). Thus, with the monsoon extending up to  $24^{\circ}\text{N}$ , it likely  
was insufficient to support such vegetation in the northern Sahara. Our  $\delta\text{D}_{\text{wax-}n\text{-alkanoic-acid}}$  show that Mediterranean winter rainfall  
enhanced when the Sahara became vegetated (Figure 3d). This suggests that winter rainfall provided additional moisture to  
the Northern Sahara. Our finding matches the results of Cheddadi et al. (2021) who proposed that Mediterranean winter rainfall  
745 played a crucial role in the genesis and sustaining of a green Sahara. Combining proxy data from Morocco with a vegetation  
model they show that intensified winter precipitation could have provided the additional moisture needed if the winter rainfall  
zone—which is restricted to a small area along the Mediterranean coast today—extended further south into the Sahara and  
perhaps even overlapped with the northward extended monsoon zone. The formation of the paleo Faiyum Lake (Egypt, Figure  
1) is dated to 10 ka BP (Hamdan et al., 2016; Hassan et al., 2012) and may give a first hint that also in the eastern Sahara  
750 winter rainfall increased south of the present day limit of the Mediterranean rainfall zone (i.e. approximately the southern tip  
of the Nile delta, Cairo). However, the inundation of the Faiyum depression to a large extent is a result from high Nile floods  
which are linked to monsoonal rainfall and local precipitation is considered of minor importance (Hamdan et al., 2016; Hassan  
et al., 2012). Therefore, more supporting evidence for the southward extension of frequent winter rains during the AHP is  
needed. Seeking for indication that also in Northeast Africa the winter rainfall zone extended deeper into the Sahara the  
information should be potentially recorded in the  $\delta\text{D}_{\text{wax-}n\text{-alkane}}$  considering that HMW  $n$ -alkanes received substantial  
755 contributions from the upper and middle catchment while the HMW  $n$ -alkanoic acids derived mainly from the delta. Between  
11–10 ka BP—at the onset of Saharan greening—the  $\delta\text{D}_{\text{wax-}n\text{-alkane}}$  shows a reversal to enriched  $\delta\text{D}$  values (Figure 3). The  
reversal to enriched  $\delta\text{D}_{\text{wax-}n\text{-alkane}}$  may either be interpreted as a reduction in rainfall amount or as a change in the moisture  
source. A similar reversal is not recorded in the  $\delta\text{D}_{\text{wax}}$  records from Lakes Tana, Victoria and Tanganyika (Berke et al., 2012,  
Tierney et al., 2008, Costa et al., 2014). These records even suggest that the monsoon strengthened at that time. Therefore, it  
760 is unlikely that the reversal in  $\delta\text{D}_{\text{wax-}n\text{-alkane}}$  between 11.5–10.5 ka BP stems from a weakened monsoon. Lake Tanganyika as  
well as other East African archives show that the monsoon weakened during the Younger Dryas (13.8–11.5 kaBP; Figure 2b,e;  
Tierney et al., 2008; Berke et al., 2014; Talbot et al., 2007). Castañeda et al. (2016) suggested the reversal would attest to drier  
conditions during Younger Dryas stadial. However, we rule out that the enrichment in  $\delta\text{D}_{\text{wax-}n\text{-alkane}}$  is a delayed climate  
deterioration associated with the Younger Dryas considering our revised age model for core GeoB7702-3 (uncertainties:  $\pm$   
765  $\pm 360$  yrs at 379 cm, Table 1). Given the absence of a comparable signal in the headwaters, the signal most likely was generated  
in the middle part of the Nile catchment, i.e. in the Sahara, south of the Nile delta. The reversal coincides with the beginning  
decrease of  $\delta\text{D}_{\text{wax-}n\text{-alkanoic-acid}}$  and consequently with the starting intensification of winter precipitation at 11 ka BP (Figure 3).  
As such, the reversal to more enriched  $\delta\text{D}_{\text{wax-}n\text{-alkane}}$  between 11.5–10.5 ka BP probably attests to increased influence of

770 Mediterranean precipitation in the Sahara. The general enrichment of  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  relative to  $\delta D_{\text{wax } n\text{-alkane}}$  during the early  
AHP (~14.5–11 ka BP) provides evidence that the Mediterranean moisture was isotopically heavier than moisture from the  
Indian and Atlantic Oceans, the latter feeding the African summer monsoon (Figure 1). This further argues for Mediterranean  
influence on the  $\delta D_{\text{wax } n\text{-alkane}}$  during this period. The minimum in  $\delta D_{\text{wax } n\text{-alkane}}$  succeeding the reversal then probably also stems  
from the amplification of winter precipitation, given the strong similarity to the minimum in  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  which occurs  
concurrently (Figures 2, 3). This inference is supported by the absence of similar minima in  $\delta D_{\text{wax}}$  from Lakes Tana, Victoria  
775 and Tanganyika at that time (Berke et al., 2012; Costa et al., 2014; Tierney et al., 2008). Seeking to explain the minimum in  
 $\delta D_{\text{wax } n\text{-alkane}}$  between 10 and 7 ka BP Castañeda et al. (2016a) invoked an eastward shift of the Congo Air Boundary during the  
AHP optimum which would enhance the influence of Atlantic moisture over the headwaters. However, the striking similarity  
between the  $\delta D_{\text{wax } n\text{-alkane}}$  and  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  is a profound argument for the influence of winter precipitation. Acknowledging  
that  $\delta D_{\text{wax } n\text{-alkane}}$  remain depleted relative to the  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  despite the influence of Mediterranean moisture (Figure 3),  
780 monsoon precipitation probably still substantially influenced the  $\delta D_{\text{wax } n\text{-alkanes signal}}$ . We conclude that during the peak phase of  
the AHP (11–6 ka BP), the  $\delta D_{\text{wax } n\text{-alkanes}}$  record monsoonal precipitation superimposed by Mediterranean winter precipitation.  
As such, we deduce that the winter rainfall zone shifted southward during the AHP optimum. Our data confirm that the  
interplay of monsoon (summer) and Mediterranean (winter) may have provided the Sahara with sufficient rainfall throughout  
the year allowing for plants to occupy the nowadays barren desert (Cheddadi et al., 2021).

785 Providing insights into the development of summer and winter precipitation in the Nile-River watershed our findings are  
important for the understanding of the moisture sources which sustained the “Green Sahara”. As elaborated above and by  
Castañeda et al. (2016a), the  $\delta D_{\text{wax } n\text{-alkane}}$  attest to enhanced monsoon rainfall in the upper Nile-River watershed. By contrast,  
our  $\delta D_{\text{wax } n\text{-alkane}}$  is a signal of winter rainfall from the lower catchment. It is widely accepted that the northward migration and  
intensification of the summer monsoon was critical to trigger plant growth in the Sahara during the AHP (Braconnot et al.,  
790 2007; Menviel et al., 2021; deMenocal et al., 2000b; Shanahan et al., 2015; Sha et al., 2019; Tierney et al., 2017). However,  
discrepancies between proxy data and climate simulations regarding precipitation and vegetation anomalies (Chandan et al.,  
2020; Braconnot et al., 2007; Hopcroft et al., 2017; Perez-Sanz et al., 2014; Cheddadi et al., 2021; Hély et al., 2014) question  
that the monsoon was the only moisture source to sustain a green Sahara (Cheddadi et al., 2021). According to climate model  
simulations monsoonal precipitation may not have provided enough moisture to sustain vegetation beyond ~24°N given  
795 mismatches with proxy data (Chandan and Peltier, 2020; Braconnot et al., 2007; Perez-Sanz et al., 2014; Cheddadi et al.,  
2021). Savannah centred between 20–25°N partly reaching to ~28–31°N (Larrasoña et al., 2013; Hély et al., 2014; Giraudi et  
al., 2013; Hamdan et al., 2016). The northeastern Sahara, south of the Nile Delta, turned into a semi-arid landscape covered  
by patchy vegetation comprising desert and open grass savannah as well as Mediterranean vegetation near the coast (Hamdan  
et al., 2016; Larrasoña et al., 2013). Wetlands and trees likely flourished near permanent water bodies such as the Faiyum  
800 Lake (Hamdan et al., 2016; Watrin et al., 2009). Thus, since the monsoon extended no further than ~22°N over northeastern  
Africa (e.g. Palchan and Torfstein, 2019; Arz et al., 2003), it likely was insufficient to support such vegetation in the lower  
Nile-River Basin (between 22–31°N). Our  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  show that Mediterranean winter rainfall enhanced during the AHP

suggesting that enhanced winter rainfall created favourable conditions for the establishment of the desert-steppe vegetation around the Nile Delta. The modern winter rainfall zone is restricted to a small area along the Mediterranean coast (between ~30-31°N over northeastern Africa). If winter rains not only intensified but also protruded deeper into the Sahara they may have provided the moisture needed to sustain the grasslands north of 22°N in the northeastern Sahara (Cheddadi et al., 2021). Seeking for indication that the winter rainfall zone extended deeper into the Sahara the information should be potentially discovered in the  $\delta D_{\text{wax } n\text{-alkane}}$  considering that HMW *n*-alkanes receive substantial contributions from the upper and middle catchment while the HMW *n*-alkanoic acids derive mainly from the delta region. Between 11-10 ka BP – at the onset of the AHP and green Sahara optimum – the  $\delta D_{\text{wax } n\text{-alkane}}$  shows a reversal to enriched  $\delta D$  values (Figure 5d). The reversal to enriched  $\delta D_{\text{wax } n\text{-alkane}}$  may either be interpreted as a reduction in rainfall amount or as a change in the moisture source. A similar reversal is not recorded in the  $\delta D_{\text{wax}}$  records from Lakes Tana, Victoria and Tanganyika (Berke et al., 2012, Tierney et al., 2008, Costa et al., 2014). These records even suggest that the monsoonal rainfall reached its maximum at this time. Therefore, it is unlikely that the reversal in  $\delta D_{\text{wax } n\text{-alkane}}$  between 11.5-10.5 ka BP stemmed from a weakened monsoon. Lake Tanganyika as well as other East African archives show that the monsoon weakened during the Younger Dryas (13.8-11.5 kaBP; Figure 4b,c; Tierney et al., 2008; Berke et al., 2014; Talbot et al., 2007). Analogously, Castañeda et al. (2016a) suggested the reversal would attest to drier conditions during Younger-Dryas stadial. However, we rule out that the enrichment in  $\delta D_{\text{wax } n\text{-alkane}}$  is a delayed climate deterioration associated with the Younger Dryas considering our revised age model for core GeoB7702-3 (uncertainties:  $\sim \pm 370$  yrs at 279 cm, median age: 11457 yrs, Table 1). Given the absence of a comparable signal in the  $\delta D_{\text{wax}}$  records from the headwaters, the Dead Sea and the  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  (Figure 5a-d), the signal most likely was generated in the middle part of the Nile catchment, i.e. in the Sahara, south of the Nile delta. The reversal coincides with the beginning decrease of  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  and consequently with the starting intensification of winter precipitation at 11 ka BP (Figure 5d). As such, the reversal to more enriched  $\delta D_{\text{wax } n\text{-alkane}}$  between 11.5-10.5 ka BP probably attests to increased influence of Mediterranean precipitation in the Sahara. The general enrichment of  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  relative to  $\delta D_{\text{wax } n\text{-alkane}}$  during the early AHP (~14.5-11 ka BP) provides evidence that the Mediterranean moisture was isotopically heavier than moisture from the Indian and Atlantic Oceans, the latter feeding the African summer monsoonal rainfall (Figure 1). This further argues for Mediterranean influence on the  $\delta D_{\text{wax } n\text{-alkane}}$  during this period. The minimum in  $\delta D_{\text{wax } n\text{-alkane}}$  succeeding the reversal (10.5-7 ka BP; Figure 5d) then probably also stemmed from the amplification of winter precipitation, given the strong similarity to the minimum in  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  which occurred concurrently (Figures 4d, 5d). This inference is supported by the absence of similar minima in  $\delta D_{\text{wax}}$  from Lakes Tana, Victoria and Tanganyika at these times (Figure 4a-c; Berke et al., 2012, Costa et al., 2014, Tierney et al., 2008). Seeking to explain the minimum in  $\delta D_{\text{wax } n\text{-alkane}}$  between 10 and 7 ka BP Castañeda et al. (2016a) invoked an eastward shift of the Congo Air Boundary during the AHP optimum which would enhance the influence of Atlantic moisture over the headwaters. However, the striking similarity between the  $\delta D_{\text{wax } n\text{-alkane}}$  and  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  is a profound argument for the influence of winter precipitation. Acknowledging that  $\delta D_{\text{wax } n\text{-alkane}}$  remain depleted relative to the  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  despite the influence of Mediterranean moisture (Figure 4d, 5d), monsoon precipitation probably still substantially influenced the  $\delta D_{\text{wax } n\text{-alkanes-signal}}$ . We conclude that the  $\delta D_{\text{wax } n\text{-alkanes}}$  recorded monsoonal precipitation superimposed by Mediterranean winter precipitation

840 between ~12-6 ka BP, broadly consistent with the AHP optimum. As such, we deduce that this episode corresponds to the  
timeframe of the southward extension of Mediterranean winter rainfall zone. Our finding supports earlier studies proposing  
southward extended winter rains during the early-to-mid Holocene in northeast Africa. South of the Nile Delta, the formation  
of the paleo Faiyum-Lake (Egypt, Figure 1) which is dated to ~10 ka BP (Hamdan et al., 2016; Hassan et al., 2012) hints to  
increased winter rains south of the present-day limit of the Mediterranean rainfall zone (i.e. approximately the southern tip of  
the Nile delta, Cairo). However, the inundation of the Faiyum depression to a large extent is a result from high Nile floods  
which are linked to monsoonal rainfall in the headwaters and local precipitation is considered of minor importance for the lake  
formation (Hamdan et al., 2016; Wendorf and Schild, 1976; Hassan, 1986). Lacustrine sediments at the northern shore of the  
845 modern Lake Quarun, the relict of the paleo Faiyum Lake, attest to Wadi activity in the region which is independent from Nile  
flow and thus indicative of stronger Mediterranean winter rainfall (Hamdan et al., 2020; Koopmann et al., 2016). Moreover,  
several authors argue that the arrival of East Asian domesticated cereals in the Faiyum region during the early Holocene was  
enabled by a southward extension of the winter rainfall zone as most of these crops rely on precipitation in winter to grow and  
on long daylight hours in summer (Phillipps et al., 2016; Shirai et al., 2016). Salinity reconstructions from the Red Sea (Art et  
850 al., 2003) and tufa formation on the southern Sinai Peninsula (Hamdan and Brook, 2015) provide evidence that Mediterranean  
winter rains reached further south than nowadays. Cheddadi et al. (2021) proposed that the Mediterranean winter rainfall may  
have reached as far south as 18-24°N based on a combination of proxy data from Morocco and vegetation modelling.  
Accordingly, winter rainfall may have played a crucial role in sustaining the green Sahara. Our data confirm that during the  
AHP optimum an interplay of monsoon (summer) and Mediterranean (winter) rainfall existed in the Nile River watershed  
855 which may have provided the northeastern Sahara with sufficient rainfall throughout the year allowing for plants to occupy  
the nowadays barren desert.

## 6. Conclusions

860 Analyzing  $\delta D_{\text{wax } n\text{-alkanoic acid}}$  in core GeoB7702-3 from the Eastern Mediterranean we provide a continuous record for winter  
precipitation in the Nile-River delta for the past 18 ka BP, in a region where continuous records are sparse. By comparison to  
previously published records of  $\delta D_{\text{wax } n\text{-alkanes}}$  from the same sediment core we gain new information about the provenance of  
leaf-wax lipids in the Nile River watershed and about the interplay of Mediterranean (winter) and monsoonal (summer)  
precipitation around the AHP and their significance for the genesis and sustaining of the Green Sahara. Our key findings can  
be summarized as follows:

865 1) HMW *n*-alkanoic acids predominantly derived from the delta during the past 18 ka while the source of the HMW *n*-alkanes  
varied through time as a function of river runoff and vegetation coverage in the Sahara. Between 15-4 ka BP the HMW *n*-  
alkanes received major contributions from the headwaters and the Sahara once the river runoff was relatively high due to  
intensified summer rains. Before and after this interval HMW *n*-alkanes derived from the delta region, like the HMW *n*-

alkanoic acids. Around the AHP (14.5-5 ka BP) the paired application of  $\delta D_{\text{wax } n\text{-alkanoic acids}}$  and  $\delta D_{\text{wax } n\text{-alkane}}$  allows to reconstruct winter rainfall ( $\delta D_{\text{wax } n\text{-alkanoic acid}}$ ) along with summer monsoonal precipitation ( $\delta D_{\text{wax } n\text{-alkane}}$ ) in the Nile-River watershed

2) HS1 occurred in two phases in the Nile River delta due to rapid variations in the position of the westerly  $\bar{J}$ Jet and the associated storm track over the Mediterranean. First the climate became arid due to a northward shift of the storm track. At 16 ka BP wet conditions established as the storm track was pushed south. Afterwards the climate remained relatively dry until 11 ka BP as the westerly  $\bar{J}$ Jet moved north in response to ice sheet retreat.

3) In the Nile-delta we find evidence for increased Mediterranean winter rainfall between 11-6 ka BP corresponding the optimum of the AHP. Amplified winter precipitation resulted from a combination of regional/local and large-scale factors. Local cyclogenesis (Cyprus Lows) promoted by elevated SSTs and southward displaced westerlies (negative NAO like pattern) steering Atlantic storms into the Mediterranean intensified the moisture delivery and precipitation in the Nile-River delta region. The episode of intensified rainfall was much shorter in the delta region than in the southern Nile catchment, where the summer monsoon determined precipitation. In the headwaters the hydroclimate had already begun to ameliorate as early as 14.5 ka BP. Likely, winter precipitation only increased when the storm track reached its southernmost position during maximal difference between NH summer and NH winter insolation.

4) At the time of Sahara greening and enhanced precipitation in the delta region (11-6 ka BP) the monsoonal signal in the  $\delta D_{\text{wax } n\text{-alkane}}$  was superimposed by intensified Mediterranean precipitation in the Sahara. Given that we find proof for concurrently intensified winter and summer precipitation in the Nile-River watershed and infer that westerly precipitation was enhanced not only in the delta region but also penetrated farther south into the Sahara, our data support the recently raised hypothesis that monsoonal precipitation only together with winter precipitation could allow for vegetation to occupy the Sahara Desert.

### Data availability

The data generated in this study will be accessible on the PANGAEA database:

<https://doi.pangaea.de/10.1594/PANGAEA.961489>

<https://www.pangaea.de/>

### Competing interests

The authors declare that they have no conflict of interest.

### Author contributions

ES designed the study together with VM, GM and JP. JP provided sediment material for core GeoB7702-3. VM carried out the sample preparation and analysis in the laboratories and conducted the data processing. IC and SS provided the contents of HMW *n*-alkanes. VM drafted the manuscript with contributions from all co-authors.

ES designed the study together with VM, GM and JP. JP provided sediment material for core GeoB7702-3. VM carried out the sample and data processing. VM drafted the manuscript with contributions from all co authors.

## Acknowledgements

We thank the captain and the crew of RV Meteor for their effort during cruise M52/2. Pushpak Nadar is thanked for his valuable help during processing the biomarker and foraminifera samples in the laboratories at MARUM. Ralph Kreutz is acknowledged for his support during the isotope analysis. We thank Liz Bonk, Hendrik Grotheer and Torben Gentz for their efforts during the AMS measurements. This work was funded by the DFG Research Center/Excellence Cluster “The Ocean Floor – Earth’s Uncharted Interface”. We are grateful to two anonymous reviewers and the editor Christo Buizert for constructive comments during the review process which helped to improve the quality of the manuscript.

## References

- 910 AbuBakr, M., Ghoneim, E., El-Baz, F., Zeneldin, M. and Zeid, S.: Use of radar data to unveil the paleolakes and the ancestral course of Wadi El-Arish, Sinai Peninsula, Egypt, *Geomorphology*, 194, 34–45, doi:10.1016/j.geomorph.2013.04.005, 2013.
- Agrawal, S., Galy, V. V., Sanyal, P. and Eglinton, T. I.: C4 plant expansion in the Ganga Plain during the last glacial cycle: insights from isotopic composition of vascular plant biomarkers. *Org. Geochem.*, 67, 58–71, doi:10.1016/j.orggeochem.2013.12.007, 2014.
- 915 [Almogi-Labin, A., Bar-Matthews, M., Shriki, D., Kolosovsky, E., Paterne, M., Schilman, B., Ayalon, A., Aizenshtat, Z. and Matthews, A.: Climatic variability during the last ~90 ka of the southern and northern Levantine Basin as evident from marine records and speleothems. \*Quat. Sci. Rev.\*, 28, 2882–2896, doi:10.1016/j.quascirev.2009.07.017, 2009.](#)
- Alpert, P., Baldi, M., Ilani, R., Krichak, S., Price, C., Rodó, X., Saaroni, H., Ziv, B., Kishcha, P., Barkan, J., Mariotti, A. and Xoplaki, E.: Relations between climate variability in the Mediterranean region and the tropics: ENSO, South Asian and African monsoons, hurricanes and Saharan dust, *Dev. Earth Environ. Sci.*, 4, 149–177, doi:10.1016/S1571-9197(06)80005-4, 2006.
- Alpert, P. Neeman, B. and Shay-El, Y.: Climatological analysis of Mediterranean cyclones using ECMWF data, *Tellus*, 42A, 65–77, DOI:10.3402/tellusa.v42i1.11860, 1990.
- Arz, H., Lamy, F., Pätzold, J., Müller, P. and Prins, M. A.: Mediterranean moisture source for an Early-Holocene Humid Period in the Northern Red Sea, *Science*, 300, 118–121, doi:10.1126/science.1080325, 2003.
- 925 Bar-Matthews, M., Ayalon, A., Gilmour, M., Matthews, A. and Hawkesworth, C. J.: Sea - land oxygen isotopic relationships from planktonic foraminifera and speleothems in the Eastern Mediterranean region and their implication for paleorainfall during interglacial intervals, *Geochim. Cosmochim. Acta*, 67, 3181–3199, doi:10.1016/S0016-7037(02)01031-1, 2003.



- Bar-Matthews, M., Ayalon, A. and Kaufman, A.: Late Quaternary climate in the eastern Mediterranean region and inferences from the stable isotope systematics of speleothems of the Soreq cave (Israel), *Quat. Res.* 47, 155-168, doi:10.1006/qres.1997.1883, 1997.
- 930 Bartlein, P. J., Harrison, S. P., Brewer, S., Connor, S., Davis, B. A. S., Gajewski, K., Guiot, J., Harrison-Prentice, T. I., Henderson, A., Peyron, O., Prentice, I. C., Scholze, M., Seppä, H., Shuman, B., Sugita, S., Thompson, R. S., Viau, A. E., Williams, J. and Wu, H.: Pollen-based continental climate reconstructions at 6 and 21 ka: A global synthesis, *Clim. Dyn.*, 37, 775–802, doi:10.1007/s00382-010-0904-1, 2011.
- 935 Berger, A. and Loutre, M. F.: Insolation values for the climate of the last 10 million years, *Quat. Sci. Rev.*, 10, 297–317, doi:10.1016/0277-3791(91)90033-Q, 1991.
- Berke, M. A., Johnson, T. C., Werne, J. P., Grice, K., Schouten, S. and Sinninghe Damsté, J. S.: Molecular records of climate variability and vegetation response since the Late Pleistocene in the Lake Victoria basin, East Africa, *Quat. Sci. Rev.*, 55, 59–74, doi:10.1016/j.quascirev.2012.08.014, 2012.
- 940 Berke, M. A., Johnson, T. C., Werne, J. P., Livingstone, D. A., Grice, K., Schouten, S. and Sinninghe Damsté, J. S.: Characterization of the last deglacial transition in tropical East Africa: Insights from Lake Albert, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 409, 1–8, doi:10.1016/j.palaeo.2014.04.014, 2014.
- Blanchet, C. L., Contoux, C. and Leduc, G.: Runoff and precipitation dynamics in the Blue and White Nile catchments during the mid-Holocene: A data-model comparison, *Quat. Sci. Rev.*, 130, 222–230, doi:10.1016/j.quascirev.2015.07.014, 2015.
- 945 Blanchet, C., Frank, M. and Schouten, S.: Asynchronous Changes in Vegetation, Runoff and Erosion in the Nile River Watershed during the Holocene, *PLoS One*, 9, 1–18, doi:10.1371/journal.pone.0115958, 2014.
- Blanchet, C. L., Osborne, A. H., Tjallingii, R., Ehrmann, W., Friedrich, T., Timmermann, A., Brückmann, W. and Frank, M.: Drivers of river reactivation in North Africa during the last glacial cycle, *Nat. Geosci.*, 14, 97-103, doi:10.1038/s41561-020-00671-3, 2021.
- 950 Blaauw, M. and Christen, A.: "Flexible paleoclimate age-depth models using an autoregressive gamma process." *Bayesian Anal.* 6, 457-474, doi:10.1214/11-BA618, 2011.
- Bowen, G. J., Wassenaar, L. I. and Hobson, K. A.: Global application of stable hydrogen and oxygen isotopes to wildlife forensics, *Oecologia*, 143, 337–348, doi:10.1007/s00442-004-1813-y, 2005.
- Braconnot, P., Otto-Bliesner, B., Harrison, S., Joussaume, S., Peterchmitt, J.-Y., Abe-Ouchi, A., Crucifix, M., Driesschaert, E., Fichefet, Th., Hewitt, C. D., Kageyama, M., Kitoh, A., Lâiné, A., Loutre, M.-F., Marti, O., Merkel, U., Ramstein, G., 955 Valdes, P., Weber, S. L., Yu, Y., and Zhao, Y.: Results of PMIP2 coupled simulations of the Mid-Holocene and Last Glacial Maximum – Part 1: experiments and large-scale features, *Clim. Past*, 3, 261–277, <https://doi.org/10.5194/cp-3-261-2007>, 2007
- Breitenbach, S. F. M., Adkins, J. F., Meyer, H., Marwan, N., Kumar, K. K. and Haug, G. H.: Strong influence of water vapor source dynamics on stable isotopes in precipitation observed in Southern Meghalaya, NE India, *Earth Planet. Sci. Lett.*, 292, 960 212–220, doi:10.1016/j.epsl.2010.01.038, 2010.



[Broecker, W. S., McGee, D., Adams, K. D., Cheng, H., Edwards, R. L., Oviatt, C. G. and Quade, J.: A Great Basin-wide dry episode during the first half of the Mystery Interval?, \*Quat. Sci. Rev.\*, 28, 2557–2563, doi:10.1016/j.quascirev.2009.07.007, 2009.](#)

Camberlin, P.: Nile Basin Climates. In: *The Nile*. Springer, pp. 307–333, 2009.

965 Castañeda, I. S., Schefuß, E., Pätzold, J., Sinninghe Damsté, J. S., Weldeab, S. and Schouten, S.: Millennial-scale sea surface temperature changes in the eastern Mediterranean (Nile River Delta region) over the last 27,000 years, *Paleoceanography*, 25, 1–13, doi:10.1029/2009PA001740, 2010a.

Castañeda, I. S., Schefuß, E., Pätzold, J., Sinninghe Damsté, Weldeab, S. and Schouten, S.: (Table 1) Radiocarbon dating of sediment core GeoB7702-3. PANGAEA, doi:10.1594/PANGAEA.736913, 2010b.

970 Castañeda, I. S., Schefuß, E., Pätzold, J., Sinninghe Damsté, Weldeab, S. and Schouten, S.: Isoprenoidal GDGT and alkenone based proxies of sediment core GeoB7702-3. PANGAEA, doi:10.1594/PANGAEA.736909, 2010c.

Castañeda, I. S., Schouten, S., Pätzold, J., Lucassen, F., Kasemann, S., Kuhlmann, H. and Schefuß, E.: Hydroclimate variability in the Nile River Basin during the past 28,000 years, *Earth Planet. Sci. Lett.*, 438, 47–56, doi:10.1016/j.epsl.2015.12.014, 2016a.

975 Castañeda, I. S., Schouten, S., Pätzold, J., Lucassen, F., Kasemann, S. A., Kuhlmann, H., and Schefuß.: Organic geochemical analyses of sediment core GeoB7702-3, PANGAEA, doi.org:10.1594/PANGAEA.858559, 2016b

Chandan, D. and Peltier, W. R.: African humid period precipitation sustained by robust vegetation, soil, and Lake feedbacks. *Geophys. Res. Lett.*, 47, e2020GL088728, doi:10.1029/2020GL088728, 2020.

980 Cheddadi, R., Carré, M., Nourelbait, M., François, L., Rhoujjati, A., Manay, R., Ochoa, D. and Schefuß, E.: Early Holocene greening of the Sahara requires Mediterranean winter rainfall, *Proc. Natl. Acad. Sci.*, 118, e2024898118, doi:10.1073/pnas.2024898118, 2021.

[Cheng, H., Sinha, A., Wang, X., Cruz, F.W. and Edwards, R.L.: The Global Paleomonsoon as seen through speleothem records from Asia and the Americas, \*Clim. Dym.\*, 39, 1045-1062, doi:10.1007/s00382-012-1363-7, 2012.](#)

985 Cheng, H., Sinha A., Verheyden, S., Nader, F. H., Li, X. L., Zhang, P. Z., Yin, J. J., Yi, L., Peng, Y. B., Rao, Z. G., Ning, Y. F. and Edwards, R. L.: The climate variability in northern Levant over the past 20,000 years, *Geophys. Res. Lett.*, 42, 8641–8650, doi:10.1002/2015GL065397, 2015.

Cheng, H., Lawrence Edwards, R., Sinha, A., Spötl, C., Yi, L., Chen, S., Kelly, M., Kathayat, G., Wang, X., Li, X., Kong, X., Wang, Y., Ning, Y. and Zhang, H.: The Asian monsoon over the past 640,000 years and ice age terminations, *Nature*, 534, 640–646, <https://doi.org/10.1038/nature18591>, 2016.

990 Claussen, M., Dallmeyer, A. and Bader, J.: Theory and modeling of the African humid period and the green Sahara, in: *Oxford Research Encyclopedia: Climate Science Vol. 1*, Oxford Research Encyclopedia, Ed., Oxford University Press USA, 2017.

[Clark, P. U., Dyke, A. S., Shakun, J. D., Carlson, A. E., Clark, J., Wohlfarth, B., Mitrovica, J. X., Hostetler, S. W. and McCabe, A. M.: The Last Glacial Maximum, \*Science\*, 325\(5941\), 710–714, doi:10.1126/science.1172873, 2009.](#)

- Collins, J. A., Schefuß, E., Mulitza, S., Prange, M., Werner, M., Tharammal, T., Paul, A. and Wefer, G.: Estimating the hydrogen isotopic composition of past precipitation using leaf-waxes from western Africa, *Quat. Sci. Rev.*, 65, 88–101, doi:10.1016/j.quascirev.2013.01.007, 2013.
- Collins, J. A., Prange, M., Caley, T., Gimeno, L., Beckmann, B., Mulitza, S., Skonieczny, C., Roche, D. and Schefuß, E.: Rapid termination of the African Humid Period triggered by northern high-latitude cooling, *Nat. Commun.*, 8, doi:10.1038/s41467-017-01454-y, 2017.
- Columbu, A., Spötl, C., Fohlmeister, J., Hu, H. M., Chiarini, V., Hellstrom, J., Cheng, H., Shen, C. C. and De Waele, J.: Central Mediterranean rainfall varied with high northern latitude temperatures during the last deglaciation, *Commun. Earth Environ.*, 3, 1–9, doi:10.1038/s43247-022-00509-3, 2022.
- Costa, K., Russell, J., Konecky, B. and Lamb, H.: Isotopic reconstruction of the African Humid Period and Congo Air Boundary migration at Lake Tana, Ethiopia, *Quat. Sci. Rev.*, 83, 58–67, doi:10.1016/j.quascirev.2013.10.031, 2014.
- De Jonge, C., Stadnitskaia, A., Hopmans, E. C., Cherkashov, G., Fedotov, A. and Sinninghe Damsté, J. S.: In situ produced branched glycerol dialkyl glycerol tetraethers in suspended particulate matter from the Yenisei River, Eastern Siberia, *Geochim. Cosmochim. Acta*, 125, 476–491, doi:10.1016/j.gca.2013.10.031, 2014.
- deMenocal, P. B.: Palaeoclimate: End of the African humid period, *Nat. Geosci.*, 8, 86–87, doi:10.1038/ngeo2355, 2015.
- deMenocal, P., Ortiz, J., Guilderson, T., Adkins, J., Sarnthein, M., Baker, L. and Yarusinsky, M.: Abrupt onset and termination of the African Humid Period: rapid climate responses to gradual insolation forcing, *Quat. Sci. Rev.*, 19, 347–361, doi:10.1016/S0277-3791(99)00081-5, 2000a.
- deMenocal P., Ortiz J., Guilderson T. and Sarnthein M.: Coherent high- and low-latitude climate variability during the Holocene warm period, *Science*, 288, 2198–2202, doi: 10.1126/science.288.5474.2198, 2000b.
- Dixit, Y., Toucanne, S., Fontanier, C., Pasquier, V., Lora, J. M., Jouet, G. and Tripathi, A.: Enhanced western Mediterranean rainfall during past interglacials driven by North Atlantic pressure changes, *Quat. Int.*, 553, 1–13, doi:10.1016/j.quaint.2020.08.017, 2020.
- Dupont, L. M. and Schefuß, E.: The roles of fire in Holocene ecosystem changes of West Africa, *Earth Planet. Sci. Lett.*, 481, 255–263, 2018.
- Eglinton, T. I. and Eglinton, G.: Molecular proxies for paleoclimatology, *Earth Planet. Sci. Lett.*, 275, 1–16, doi:10.1016/j.epsl.2008.07.012, 2008.
- Eglinton, G. and Hamilton, R. J.: Leaf epicuticular waxes, *Science*, 156, 1322–1335, doi: 10.1126/science.156.3780.1322, 1967.
- Emeis, K. C., Struck, U., Schulz, H. M., Rosenberg, R., Bernasconi, S., Erlenkeuser, H., Sakamoto, T. and Martinez-Ruiz, F.: Temperature and salinity variations of Mediterranean Sea surface waters over the last 16,000 years from records of planktonic stable oxygen isotopes and alkenone unsaturation ratios, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 158, 259–280, doi:10.1016/S0031-0182(00)00053-5, 2000.

- Enzel, Y., Amit, R., Dayan, U., Crouvi, O., Kahana, R., Ziv, B. and Sharon, D.: The climatic and physiographic controls of the eastern Mediterranean over the late Pleistocene climates in the southern Levant and its neighboring deserts, *Global Planet. Change* 60, 165-192, doi:10.1016/j.gloplacha.2007.02.003, 2008.
- 1030 Eshel, G. and Farrel, B. F.: Mechanisms of eastern Mediterranean rainfall variability, *J. Atmos. Sci.*, 57, 3219–32, doi:10.1175/1520-0469(2000)057<3219:MOEMRV>2.0.CO;2, 2000.
- Fletcher, W. J., Goni, M. F. S., Peyron, O. and Dormoy, I.: Abrupt climate changes of the last deglaciation detected in a Western Mediterranean forest record, *Clim. Past*, 6, 245–264, doi:10.5194/cp-6-245-2010, 2010.
- Fletcher, W. J. and Sanchez Goñi, M. F.: Orbital- and sub-orbital-scale climate impacts on vegetation of the western  
1035 Mediterranean basin over the last 48,000 yr. *Quat. Res.*, 70, 451-464, doi:10.1016/j.yqres.2008.07.002, 2008.
- Galy, V. and Eglinton, T.: Protracted storage of biospheric carbon in the Ganges-Brahmaputra basin, *Nat. Geosci.*, 4, 843–847, doi:10.1038/ngeo1293, 2011.
- Garcin, Y., Schefuß, E., Schwab, V. F., Garreta, V., Gleixner, G., Vincens, A., Todou, G., Séné, O., Onana, J. M., Achoundong, G. and Sachse, D.: Reconstructing C3 and C4 vegetation cover using n-alkane carbon isotope ratios in recent lake sediments  
1040 from Cameroon, Western Central Africa, *Geochim. Cosmochim. Acta*, 142, 482–500, doi:10.1016/j.gca.2014.07.004, 2014.
- Gensel, J., Humphries, M. S., Zabel, M., Sebag, D., Hahn, A. and Schefuß, E.: Origin, transport, and retention of fluvial sedimentary organic matter in South Africa's largest freshwater wetland, Mkhuzi Wetland System, *Biogeosciences*, 19, 2881–2902, doi:10.5194/bg-19-2881-2022, 2022.
- Giraudi, C., Mercuri, A. M. and Esu, D.: Holocene palaeoclimate in the northern Sahara margin (Jefara Plain, northwestern  
1045 Libya), *The Holocene*, 23, 339–352, doi:10.1177/0959683612460787, 2013.
- Goldsmith, Y., Polissar, P. J., Ayalon, A., Bar-Matthews, M., deMenocal, P. B. and Broecker, W. S.: The modern and Last Glacial Maximum hydrological cycles of the Eastern Mediterranean and the Levant from a water isotope perspective, *Earth Planet. Sci. Lett.*, 457, 302–312, doi:10.1016/j.epsl.2016.10.017, 2017.
- Häggi, C., Sawakuchi, A. O., Chiessi, C. M., Mülitz, S., Mollenhauer, G., Sawakuchi, H. O., Baker, P. A., Zabel, M. and  
1050 Schefuß, E.: Origin, transport and deposition of leaf-wax biomarkers in the Amazon Basin and the adjacent Atlantic, *Geochim. Cosmochim. Acta*, 192, 149–165, doi:10.1016/j.gca.2016.07.002, 2016.
- Hamann, Y., Ehrmann, W., Schmiidl, G., Krüger, S., Stuetz, J. B. and Kuhnt, T.: Sedimentation processes in the Eastern Mediterranean Sea during the Late Glacial and Holocene revealed by end-member modelling of the terrigenous fraction in marine sediments, *Mar. Geol.*, 248, 97–114, doi:10.1016/j.margeo.2007.10.009, 2008.
- 1055 Hamdan, M. A. and Brook, G. A.: Timing and characteristics of Late Pleistocene and Holocene wetter periods in the Eastern Desert and Sinai of Egypt, based on 14C dating and stable isotope analysis of spring tufa deposits, *Quat. Sci. Rev.*, 130, 168–188, doi:10.1016/j.quascirev.2015.09.011, 2015.
- Hamdan, M. A., Flower, R. J., Hassan, F. A. and Leroy, S. A. G.: Geochemical and palynological analysis of Faiyum Lake sediments, Egypt: Implications for Holocene paleoclimate, *J. African Earth Sci.*, 167, 103864, doi:10.1016/j.jafrearsci.2020.103864, 2020.
- 1060

- Hamdan, M. A., Ibrahim, M. I. A., Shiha, M. A., Flower, R. J., Hassan, F. A. and Eltelet, S. A. M.: An exploratory Early and Middle Holocene sedimentary record with palynofossils and diatoms from Faiyum lake, Egypt, *Quat. Int.*, 410, 30–42, doi:10.1016/j.quaint.2015.12.049, 2016.
- Hamdan, M.A. and Lucarini, G.: Holocene paleoenvironmental, paleoclimatic and geoarchaeological significance of the Sheikh El-Obeiyid area (Farafra Oasis, Egypt), *Quat. Int.*, 302, 154–168, doi:10.1016/j.quaint.2013.01.009, 2013.
- 1065 [Hassan, F. A.: Holocene lakes and prehistoric settlements of the Western Faiyum, Egypt, \*J. Archaeol. Sci.\*, 13, 483–501, doi:10.1016/0305-4403\(86\)90018-X, 1986.](https://doi.org/10.1016/0305-4403(86)90018-X)
- Hassan, F. A., Hamdan, M. A., Flower, R. J. and Keatings, K.: Oxygen and carbon isotopic records in Holocene freshwater mollusc shells from the Faiyum palaeolakes, Egypt: Palaeoenvironmental and palaeoclimatic implications, *Quat. Int.*, 266, 175–187, doi:10.1016/j.quaint.2011.11.024, 2012.
- 1070 Heaton, T. J., Köhler, P., Butzin, M., Bard, E., Reimer, R. W., Austin, W. E. N., Bronk Ramsey, C., Grootes, P. M., Hughen, K. A., Kromer, B., Reimer, P. J., Adkins, J., Burke, A., Cook, M. S., Olsen, J. and Skinner, L. C.: Marine20—The Marine Radiocarbon Age Calibration Curve (0–55,000 cal. BP), *Radiocarbon*, 62: IntCal20: Calibration Issue, 779–820, doi:10.1017/RDC.2020.68, 2020.
- 1075 [Hedges, J. I., Keil, R. G. and Benner, R.: What happens to terrestrial organic matter in the ocean?, \*Org. Geochem.\*, 27, 195–212, doi:10.1016/S0146-6380\(97\)00066-1, 1997.](https://doi.org/10.1016/S0146-6380(97)00066-1)
- Hély, C., Lézine, A. M., Ballouche, A., Cour, P., Duzer, D., Guinet, P., Jahns, S., Maley, J., Mercuri, A. M., Pons, A., Ritchie, J. C., Salzmann, U., Schulz, E., Van Campo, M. and Waller, M. P.: Holocene changes in African vegetation: Tradeoff between climate and water availability, *Clim. Past*, 10, 681–686, doi:10.5194/cp-10-681-2014, 2014.
- 1080 Hemingway, J. D., Schefuß, E., Dinga, B. J., Pryer, H. and Galy, V. V.: Multiple plant-wax compounds record differential sources and ecosystem structure in large river catchments, *Geochim. Cosmochim. Acta*, 184, 20–40, doi:10.1016/j.gca.2016.04.003, 2016.
- Hoefs, M. J. L., Rijpstra, W. I. C. and Sinninghe Damste, J.S.: The influence of oxic degradation on the sedimentary biomarker record I: evidence from Madeira Abyssal Plain turbidites. *Geochim. Cosmochim. Acta*, 66, 2719–2735, doi:10.1016/S0016-1085 7037(02)00864-5, 2002.
- Hopcroft, P. O., Valdes, P. J., Harper, A., B. and Beerling, D. J.: Multi vegetation model evaluation of the green Sahara climate regime, *Geophys. Res. Lett.*, 44, 6804–6813, doi:10.1002/2017GL073740, 2017.
- <https://en.climate-data.org/africa/egypt/cairo-governorate/cairo-3392/>; (checked December, 14th 2022)
- <https://en.climate-data.org/africa/egypt/alexandria-governorate/alexandria-515/> (checked December, 14th 2022)
- 1090 <https://en.climate-data.org/africa/egypt/luxor-governate/luxor-6345/> (checked December, 14th 2022)
- <https://en.climate-data.org/africa/ethiopia/addis-ababa/addis-abeba-532/> (checked December, 14th 2022)
- <https://en.climate-data.org/africa/uganda/central-region/kampala-5578/> (checked December, 14th 2022)
- Jolly, D., Prentice, I. C., Bonnefille, R., Ballouche, A., Bengo, M., Brenac, P., Buchet, G., Burney, D., Cazet, J. P., Cheddadi, R., Ector, T., Elenga, H., Elmoutaki, S., Guiot, J., Laarif, F., Lamb, H., Lézine, A. M., Maley, J., Mbenza, M., Peyron, O.,

- 1095 Reille, M., Reynaud-Farrera, I., Riollet, G., Ritchie, J. C., Roche, E., Scott, L., Ssemmanda, I., Straka, H., Umer, M., Van Campo, E., Vilimumbalo, S., Vincens, A. and Waller, M.: Biome reconstruction from pollen and plant macrofossil data for Africa and the Arabian peninsula at 0 and 6000 years, *J. Biogeogr.*, 25, 1007–1027, doi:10.1046/j.1365-2699.1998.00238.x, 1998.
- Junginger, A., Roller, S., Olaka, L. A. and Trauth, M. H.: The effects of solar irradiation changes on the migration of the Congo Air Boundary and water levels of paleo-Lake Suguta, Northern Kenya Rift, during the African Humid Period (15–5 ka BP), *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 396, 1–16., doi:10.1016/j.palaeo.2013.12.007, 2014.
- 1100 [Kageyama, M., Harrison, S. P., Kapsch, M. L., Lofverstrom, M., Lora, J. M., Mikolajewicz, U., Sherriff-Tadano, S., Vadsaria, T., Abe-Ouchi, A., Bouttes, N., Chandan, D., Gregoire, L. J., Ivanovic, R. F., Izumi, K., Legrande, A. N., Lhardy, F., Lohmann, G., Morozova, P. A., Ohgaito, R., Paul, A., Richard Peltier, W., Poulsen, C. J., Quiquet, A., Roche, D. M., Shi, X., Tierney, J. E., Valdes, P. J., Volodin, E. and Zhu, J.: The PMIP4 Last Glacial Maximum experiments: Preliminary results and comparison with the PMIP3 simulations, \*Clim. Past\*, 17\(3\), 1065–1089, doi:10.5194/cp-17-1065-2021, 2021.](#)
- 1105 [Konecky, B. L., Russell, J. M., Johnson, T. C., Brown, E. T., Berke, M. A., Werne, J. P. and Huang, Y.: Atmospheric circulation patterns during late Pleistocene climate changes at Lake Malawi, Africa, \*Earth Planet. Sci. Lett.\*, 312\(3–4\), 318–326, doi:10.1016/j.epsl.2011.10.020, 2011.](#)
- 1110 [Koopman, A., Kluiving, S., Holdaway, S. and Wendrich, W.: The Effects of Holocene Landscape Changes on the Formation of the Archaeological Record in the Fayum Basin, Egypt, \*Geoarchaeology\*, 31, 17–33, doi:10.1002/gea.21538, 2016.](#)
- Korecha, D. and Barnston, A.G.: Predictability of June–September Rainfall in Ethiopia. *Mon. Wea. Rev.*, 135, 628–650, doi:10.1175/MWR3304.1, 2007
- 1115 [Krichak, S. O., Breitgand, J. S. and Feldstein, S. B.: A conceptual model for the identification of active Red Sea Trough synoptic events over the south-eastern Mediterranean, \*J. Appl. Meteorol. Climatol.\*, 51, 962–71, doi:org/10.1175/JAMC-D-11-0223.1, 2012.](#)
- [Krom, M. D., Cliff, R. A., Eijsink, L. M., Herut, B. and Chester, R.: The characterisation of Saharan dusts and Nile particulate matter in surface sediments from the Levantine basin using Sr isotopes, \*Mar. Geol.\*, 155, 319-330, doi:10.1016/S0025-3227\(98\)00130-3, 1999.](#)
- 1120 Kuper, R. and Kröpelin, S.: Climate-controlled Holocene occupation in the Sahara: Motor of Africa’s evolution, *Science*, 313, 803–807, doi:10.1126/science.1130989, 2006.
- [Kusch, S., Rethemeyer, J., Schefuß, E. and Mollenhauer, G.: Controls on the age of vascular plant biomarkers in Black Sea sediments, \*Geochim. Cosmochim. Acta\*, 74, 7031–7047, doi:10.1016/j.gca.2010.09.005, 2010.](#)
- 1125 Kutzbach, J. E., Chen, G., Cheng, H., Edwards, R. L. and Liu, Z.: Potential role of winter rainfall in explaining increased moisture in the Mediterranean and Middle East during periods of maximum orbitally-forced insolation seasonality, *Clim. Dyn.*, 42, 1079–1095, doi:10.1007/s00382-013-1692-1, 2014.
- Lamb, H. F., Bates, C. R., Coombes, P. V., Marshall, M. H., Umer, M., Davies, S. J. and Dejen, E.: Late Pleistocene desiccation of Lake Tana, source of the Blue Nile, *Quat. Sci. Rev.*, 26, 287–299, doi:10.1016/j.quascirev.2006.11.020, 2007.

- Langgut, D., Cheddadi, R. and Sharon, G.: Climate and environmental reconstruction of the Epipaleolithic Mediterranean Levant (22.0-11.9 ka cal. BP), *Quat. Sci. Rev.* **270**, 107170, doi:10.1016/j.quascirev.2021.107170, 2021.
- Larrasoña, J. C., Roberts, A. P. and Rohling, E. J.: Dynamics of Green Sahara Periods and Their Role in Hominin Evolution, *PLoS One*, **8**, e76514, doi:10.1371/journal.pone.0076514, 2013.
- Li, Y., Song, Y., Yin, Q., Han, L. and Wang, Y.: Orbital and millennial northern mid-latitude westerlies over the last glacial period, *Clim. Dyn.*, **53**, 3315–3324, doi:10.1007/s00382-019-04704-5, 2019.
- Lenderink, G. and Meijgaard, E. V.: Increase in hourly precipitation extremes beyond expectations from temperature changes, *Nat. Geosci.* **1**, 511-514. doi: 10.1038/ngeo262, 2008.
- Lisiecki, L. E., and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic  $\delta^{18}O$  records, *Paleoceanography*, **20**, PA1003, doi:10.1029/2004PA001071, 2005.
- Ludwig, P. and Hochman, A.: Last glacial maximum hydro-climate and cyclone characteristics in the Levant: A regional modelling perspective, *Environ. Res. Lett.*, **17**, doi:10.1088/1748-9326/ac46ea, 2022.
- Lüning, S. and Vahrenholt, F.: Holocene Climate Development of North Africa and the Arabian Peninsula, in: Bendaoud, A., Hamimi, Z., Hamoudi, M., Djemai, S. and Zoheir, B.: The Geology of the Arab World - An Overview, pp. 507-546, Springer Geology, 2019, https://doi.org/10.1007/978-3-319-96794-3
- Martin, C., Ménot, G., Thouveny, N., Davtian, N., Andrieu-Ponel, V., Reille, M. and Bard, E.: Impact of human activities and vegetation changes on the tetraether sources in Lake St Front (Massif Central, France), *Org. Geochem.*, **135**, 38–52, doi:10.1016/j.orggeochem.2019.06.005, 2019
- Marshall, M.H., Lamb, H.F., Huws, D., Davies, S.J., Bates, R., Bloemendal, J., Boyle, J., Leng, M.J., Umer, M. and Bryant, C.: Late Pleistocene and Holocene drought events at Lake Tana, the source of the Blue Nile, *Glob. Planet. Change*, **78**, 147–161, doi:10.1016/j.gloplacha.2011.06.004, 2011.
- McGee, D., deMenocal, P. B., Winckler, G., Stuut, J. B. W. and Bradtmiller, L. I.: The magnitude, timing and abruptness of changes in North African dust deposition over the last 20,000 yr, *Earth Planet. Sci. Lett.*, **371–372**, 163–176, 2013.
- ~~Meyers, P. A. and Ishiwatari, R.: Lacustrine organic geochemistry – an overview of indicators of organic matter sources and diagenesis in lake sediments, *Org. Geochem.*, **20**, 867–900, doi:10.1016/0146-6380(93)90100-P, 1993.~~
- Ménot, G., Pivot, S., Bouloubassi, I., Davtian, N., Hennekam, R., Bosch, D., Ducassou, E., Bard, E., Migeon, S. and Revel, M.: Timing and stepwise transitions of the African Humid Period from geochemical proxies in the Nile deep-sea fan sediments, *Quat. Sci. Rev.*, **228**, doi:10.1016/j.quascirev.2019.106071, 2020.
- Menviel, L., Govin, A. and Grant, K. M.: Drivers of the evolution and amplitude of African Humid Periods, *Commun. Earth Environ.*, **2**, 237, doi:10.1038/s43247-021-00309-1, 2021.
- ~~Meyers, P. A. and Ishiwatari, R.: Lacustrine organic geochemistry – an overview of indicators of organic matter sources and diagenesis in lake sediments, *Org. Geochem.*, **20**, 867–900, doi:10.1016/0146-6380(93)90100-P, 1993.~~
- Miebach, A., Stolzenberger, S., Wacker, L., Hense, A. and Litt, T.: A new Dead Sea pollen record reveals the last glacial paleoenvironment of the southern Levant, *Quat. Sci. Rev.*, **214**, 98-116, doi:10.1016/j.quascirev.2019.04.033, 2019.

- Mollenhauer, G., Grotheer, H., Gentz, T., Bonk, E. and Hefter, J.: Standard operation procedures and performance of the MICADAS radiocarbon laboratory at Alfred Wegener Institute (AWI), Germany, Nucl. Instruments Methods Phys. Res. Sect. B Beam Interact. with Mater. Atoms, 496, 45–51, doi:10.1016/j.nimb.2021.03.016, 2021.
- Muhs, D. R., Roskin, J., Tsoar, H., Skipp, G., Budahn, J. R., Sneh, A., Porat, N., Stanley, J. D., Katra, I. and Blumberg, D. G.: Origin of the Sinai-Negev erg, Egypt and Israel: Mineralogical and geochemical evidence for the importance of the Nile and sea level history, Quat. Sci. Rev., 69, 28–48, doi:10.1016/j.quascirev.2013.02.022, 2013.
- Naughton, F., Sánchez Goñi, M.F., Kageyama, M., Bard, E., Cortijo, E., Desprat, S., Malaizé, B., Joly, C., Rostek, F. and Turon, J.-L.: Wet to dry climatic trend in north western Iberia within Heinrich events, Earth Planet. Sci. Lett., 284, 329-342, doi:10.1016/j.epsl.2009.05.001, 2009.
- Naughton, F., Sanchez Goñi, M.F., Rodrigues, T., Salgueiro, E., Costas, S., Desprat, S., Duprat, J., Michel, E., Rossignol, L., Zaragosi, S., Voelker, A. H. L. and Abrantes, F.: Climate variability across the last deglaciation in NW Iberia and its margin, Quat. Int., 414, 9-22, doi:10.1016/j.quaint.2015.08.073, 2016.
- Naughton, F., Toucanne, S., Landais, A., Rodrigues, T., Vazquez Riveiros, N. and Sánchez-Goñi, M. F.: Chapter 5 – Heinrich Stadial 1. In: Palacios, D., Hughes, P. D., García-Ruiz, J. M. and Andrés N.: European Glacial Landscapes – The Last Deglaciation, Elsevier, 37-44, ISBN 9780323918992, doi.org/10.1016/B978-0-323-91899-2.00049-8, 2023.
- Otto-Bliesner, B. L., Russell, J. M., Clark, P. U., Liu, Z., Overpeck, J. T., Konecky, B., Nicholson, S. E., He, F. and Lu, Z.: Coherent changes of southeastern equatorial and northern African rainfall during the last deglaciation, Science, 346, 1223–1227, doi:10.1126/science.1259531, 2014.
- [Palchan, D. and Torfstein, A.: A drop in Sahara dust fluxes records the northern limits of the African Humid Period, Nat. Commun., 10, doi:10.1038/s41467-019-11701-z, 2019.](https://doi.org/10.1038/s41467-019-11701-z)
- Pausata, F. S. R., Messori, G. and Zhang, Q.: Impacts of dust reduction on the northward expansion of the African monsoon during the green Sahara period, Earth Planet. Sci. Lett., 434, 298–307, doi:10.1016/j.epsl.2015.11.049, 2016.
- Pausata, F. S. et al.: The greening of the Sahara: Past changes and future implications. One Earth 2, 235–250, doi:10.1016/j.oneear.2020.03.002, 2020.
- Pätzold, J., Bohrmann, G. and Hübscher, C.: Black Sea – Mediterranean – Red Sea. Cruise No. 52, January 2 – March 27, 2002, Istanbul – Limassol, Universität Hamburg, METEOR-Berichte 3-2, 178 pp, 2003.
- Pérez-Mejías, C., Moreno, A., Bernal-Wormull, J., Cacho, I., Osácar, M.C., Lawrence, E. and Chang, E.: Oldest Dryas hydroclimate reorganization in the eastern Iberian Peninsula after the iceberg discharges of Heinrich Event 1, Quat. Res., 101, 67-83, doi:10.1017/qua.2020.112, 2021.
- Perez-Sanz, A., Li, G., González-Sampériz, P., and Harrison, S. P.: Evaluation of modern and mid-Holocene seasonal precipitation of the Mediterranean and northern Africa in the CMIP5 simulations, Clim. Past, 10, 551–568, <https://doi.org/10.5194/cp-10-551-2014>, 2014.



- 1195 [Phillipps, R., Holdaway, S., Ramsay, R., Emmitt, J., Wendrich, W. and Linseele, V.: Lake level changes, lake edge basins and the paleoenvironment of the fayum North shore, Egypt, during the early to mid-holocene, \*Open Quat.\*, 2\(1929\), 1–12, doi:10.5334/oq.19, 2016.](#)
- [Phillipps, R., Holdaway, S., Wendrich, W. and Cappers, R.: Mid-Holocene occupation of Egypt and global climatic change, \*Quat. Int.\*, 251, 64–76, doi:10.1016/j.quaint.2011.04.004, 2012.](#)
- 1200 Quade, J., Dente, E., Armon, M., Ben Dor, Y., Morin, E., Adam, O. and Enzel, Y.: Megalakes in the Sahara? A Review, *Quat. Res.*, 90, 253–275, doi:10.1017/qua.2018.46, 2018.
- Reimer, P. J. and Reimer, R. W.: A marine reservoir correction database and on-line interface. *Radiocarbon*, 43,461-3, doi:10.1017/S0033822200038339, 2001.
- Revel, M., Ducassou, E., Grousset, F. E., Bernasconi, S. M., Migeon, S., Revillon, S., Mascle, J., Murat, A., Zaragosi, S. and
- 1205 Bosch, D.: 100,000 Years of African monsoon variability recorded in sediments of the Nile margin, *Quat. Sci. Rev.*, 29, 1342–1362, doi:10.1016/j.quascirev.2010.02.006, 2010.
- Revel, M., Colin, C., Bernasconi, S., Combourieu-Nebout, N., Ducassou, E., Grousset, F. E., Rolland, Y., Migeon, S., Bosch, D., Brunet, P., Zhao, Y. and Mascle, J.: 21,000 years of Ethiopian African monsoon variability recorded in sediments of the western Nile deep-sea fan, *Reg. Environ. Chang.*, 14, 1685–1696, doi:10.1007/s10113-014-0588-x, 2014.
- 1210 Revel, M., Ducassou, E., Skonieczny, C., Colin, C., Bastian, L., Bosch, D., Migeon, S. and Mascle, J.: 20,000 years of Nile River dynamics and environmental changes in the Nile catchment area as inferred from Nile upper continental slope sediments, *Quat. Sci. Rev.*, 130, 200–221, doi:10.1016/j.quascirev.2015.10.030, 2015.
- [Ruan, Y., Mohtadi, M., van der Kaas, S., Dupont, L. M., Hebbeln, D. and Schefuß, E.: Differential hydro-climatic evolution of East Javanese ecosystems over the past 22,000 years, \*Quat. Sci. Rev.\*, 218, 49-60, doi: 10.1016/j.quascirev.2019.06.015, 2019.](#)
- 1215 Sachse, D., Billault, I., Bowen, G. J., Chikaraishi, Y., Dawson, T. E., Feakins, S. J., Freeman, K. H., Magill, C. R., McInerney, F. A., van der Meer, M. T. J., Polissar, P., Robins, R. J., Sachs, J. P., Schmidt, H.-L., Sessions, A. L., White, J. W. C., West, J. B. and Kahmen, A.: Molecular paleohydrology: interpreting the hydrogen-isotopic composition of lipid biomarkers from photosynthesizing organisms, *Annu. Rev. Earth Planet. Sci.*, 40, 221–249, doi:10.1146/annurev-earth-042711-105535, 2012.
- 1220 Schefuß, E., Kuhlmann, H., Mollenhauer, G., Prange, M. and Pätzold, J.: Forcing of wet phases in southeast Africa over the past 17,000 years, *Nature*, 480, 509–512, doi:10.1038/nature10685, 2011.
- Schefuß, E., Schouten, S. and Schneider, R. R.: Climatic controls on central African hydrology during the past 20,000 years. *Nature*, 437, 1003–1006, doi:10.1038/nature03945, 2005.
- Schlitzer, R. Ocean Data View v.5.6.3 (Alfred Wegener Institute, 2006); <https://odv.awi.de/>
- 1225 Sha, L., Ait Brahim, Y., Wassenburg, J. A., Yin, J., Peros, M., Cruz, F. W., Cai, Y., Li, H., Du, W., Zhang, H., Edwards, R. L. and Cheng, H.: How Far North Did the African Monsoon Fringe Expand During the African Humid Period? Insights From Southwest Moroccan Speleothems, *Geophys. Res. Lett.*, 46, 14093–14102, doi:10.1029/2019GL084879, 2019.

- Schlünz, B. and Schneider, R. R.: Transport of terrestrial organic carbon to the oceans by rivers: re-estimating flux- and burial rates, *Int. J. Earth Sci. (Geol. Rundsch.)*, 88, 599–606, 2000.
- 1230 Schreuder, L. T., Stuut, J.-B. W., Korte, L. F., Sinninghe Damsté, J. S. and Schouten, S.: Aeolian transport and deposition of plant wax *n*-alkanes across the tropical North Atlantic Ocean. *Organic Geochemistry*, 115, 113–123. doi:10.1016/j.orggeochem.2017.10.010, 2018.
- Shanahan, T. M., McKay, N. P., Hughen, K. A., Overpeck, J. T., Otto-Bliesner, B., Heil, C. W., King, J., Scholz, C. A. and Peck, J.: The time-transgressive termination of the African humid period, *Nat. Geosci.*, 8, 140–144, doi:10.1038/ngeo2329, 1235 2015.
- Sinninghe Damsté, J. S., Rijpstra, W. I. C. and Reichart, G.-J.: The influence of oxic degradation on the sedimentary biomarker record II. Evidence from Arabian Sea sediments, *Geochim. Cosmochim. Acta*, 66, 2737–2754, doi:10.1016/S0016-7037(02)00865-7, 2002.
- Stager, J. C., Ryves, D. B., Chase, B. M. and Pausata, F. S.: Catastrophic drought in the Afro-Asian monsoon region during Heinrich event 1. *Science*, 331, 1299–1302, doi:10.1126/science.1198322, 2011.
- 1240 [Stein, M.: The sedimentary and geochemical record of Neogene-Quaternary water bodies in the Dead Sea basin - inferences for the regional paleoclimate history, \*J. Paleolimnol.\*, 26, 271–282, doi: 10.1023/A:1017529228186, 2001.](#)
- [Stein, M., Starinsky, A., Katz, A., Goldstein, S., Machlus, M. and Schramm, A.: Strontium isotopic, chemical, and sedimentological evidence for the evolution of Lake Lisan and the Dead Sea, \*Geochim. Cosmochim. Acta\* 61, 3975-3992, 1245 doi:10.1016/S0016-7037\(97\)00191-9, 1997.](#)
- [Stockhecke, M., Timmermann, A., Kipfer, R., Haug, G.H., Kwiecien, O., Friedrich, T., Menviel, L., Litt, T., Pickarski, N. and Anselmetti, F.S.: Millennial to orbital-scale variations of drought intensity in the Eastern Mediterranean, \*Quat. Sci. Rev.\* 133, 77-95, doi:org/10.1016/j.quascirev.2015.12.016, 2016.](#)
- Talbot, M. R., Filippi, M. L., Jensen, N. B. and Tiercelin, J. J.: An abrupt change in the African monsoon at the end of the Younger Dryas, *Geochemistry, Geophys. Geosystems*, 8, Q03006, doi:10.1029/2006GC001465, 2007.
- 1250 Tierney, J. E., Russell, J. M., Huang, Y., Sinninghe Damsté, J. S., Hopmans, E. C. and Cohen, A. S.: Northern hemisphere controls on tropical ~~southeast~~southeastern African climate during the past 60,000 years, *Science*, 322, 252–255, doi:10.1126/science.1160485, 2008.
- [Thompson, A. J., Tabor, C. R., Poulsen, C. J. and Skinner, C. B.: Water isotopic constraints of the mid-Holocene West African Monsoon, \*EPSL\*, 554, 11667, doi: 10.1016/j.epsl.2020.116677, 2021.](#)
- 1255 Tierney, J. E., Russell, J. M. and Huang, Y.: A molecular perspective on Late Quaternary climate and vegetation change in the Lake Tanganyika basin, East Africa, *Quat. Sci. Rev.*, 29, 787–800, doi:10.1016/j.quascirev.2009.11.030, 2010.
- Tierney, J. E., deMenocal, P. B.: Abrupt shifts in Horn of Africa hydroclimate since the Last Glacial Maximum. *Science*, 342, 843–846, doi:10.1126/science.1240411, 2013.
- 1260 Tierney, J. E., Pausata, F. S. R. and deMenocal, P. B.: Rainfall regimes of the Green Sahara, *Sci. Adv.*, 3, 1–10, doi:10.1126/sciadv.1601503, 2017.

- 1265 [Tierney, J. E., Torfstein, A. and Bhattacharya, T.: Late Quaternary hydroclimate of the Levant: The leaf wax record from the Dead Sea, \*Quat. Sci. Rev.\*, 289, 107613, doi:10.1016/j.quascirev.2022.107613, 2022.](#)
- [Tsvieli, Y. and Zangvil, A.: Synoptic climatological analysis of the ‘wet’ and ‘dry’ Red Sea troughs over Israel, \*Int. J. Climatol.\*, 25, 1997–2015, doi:10.1002/joc.1232, 2005.](#)
- [Trenberth, K. E., Dai, A., Rasmussen, R. M. and Parsons, D. B.: The changing character of precipitation. \*Bull. Am. Meteorol. Soc.\*, 84, 1205-1217. Doi: 10.1175/BAMS-84-9-1205, 2003.](#)
- 1270 Valsecchi, V., Sanchez Goñi, M. F. and Londeix, L.: Vegetation dynamics in the Northeastern Mediterranean region during the past 23 000 yr: Insights from a new pollen record from the Sea of Marmara, *Clim. Past*, 8, 1941–1956, doi:10.5194/cp-8-1941-2012, 2012.
- Viste, E. and Sorteberg, A.: The effect of moisture transport variability on Ethiopian summer precipitation, *Int. J. Climatol.*, 33, 3106–3123, doi:10.1002/joc.3566, 2013.
- 1275 Wagner, B., Vogel, H., Francke, A., Friedrich, T., Donders, T., Lacey, J. H., Leng, M. J., Regattieri, E., Sadori, L., Wilke, T., Zanchetta, G., Albrecht, C., Bertini, A., Combourieu-Nebout, N., Cvetkoska, A., Giaccio, B., Grazhdani, A., Hauffe, T., Holtvoeth, J., Joannin, S., Jovanovska, E., Just, J., Kouli, K., Kousis, I., Koutsodendris, A., Krastel, S., Lagos, M., Leicher, N., Levkov, Z., Lindhorst, K., Masi, A., Melles, M., Mercuri, A. M., Nomade, S., Nowaczyk, N., Panagiotopoulos, K., Peyron, O., Reed, J. M., Sagnotti, L., Sinopoli, G., Stelbrink, B., Sulpizio, R., Timmermann, A., Tofilovska, S., Torri, P., Wagner-Cremer, F., Wonik, T. and Zhang, X.: Mediterranean winter rainfall in phase with African monsoons during the past 1.36 million years, *Nature*, 573, 256–260, doi:10.1038/s41586-019-1529-0, 2019.
- 1280 Wang, N., Jiang, D. and Lang, X.: Northern westerlies during the last glacial maximum: Results from CMIP5 simulations, *J. Clim.*, 31, 1135–1153, doi:10.1175/JCLI-D-17-0314.1, 2018.
- Wassenburg, J. A., Dietrich, S., Fietzke, J., Fohlmeister, J., Jochum, K. P., Scholz, D., Richter, D. K., Sabaoui, A., Spötl, C., Lohmann, G., Andreae, M. O. and Immenhauser, A.: Reorganization of the North Atlantic Oscillation during early Holocene deglaciation, *Nat. Geosci.*, 9, 602–605, doi:10.1038/ngeo2767, 2016.
- 1285 [Watrin, J., Lézine, A. M., Hély, C., Cour, P., Ballouche, A., Duzer, D., Elenga, H., Frédoux, A., Guinet, P., Jahns, S., Kadamura, H., Maley, J., Mercuri, A. M., Pons, I. A., Reynaud-Farrera, I., Ritchie, J. C., Salzmann, U., Schulz, E., Tossou, M. G., Vincens, A. and Waller, M. P.: Plant migration and plant communities at the time of the “green Sahara,” \*Comptes Rendus - Geosci.\*, 341\(8–9\), 656–670, doi:10.1016/j.crte.2009.06.007, 2009.](#)
- [Weijers, J. W. H., Schouten, S., Hopmans, E. C., Geenevasen, J. A. J., David, O. R. P., Coleman, J. M., Pancost, R. D., Sinninghe Damsté, J. S.: Membrane lipids of mesophilic anaerobic bacteria thriving in peats have typical archaeal traits, \*Environ. Microbiol.\*, 8, 648–657, doi:10.1111/j.1462-2920.2005.00941.x, 2006.](#)
- 1290 Weldeab, S., Emeis, K.-C., Hemleben, C. and Siebel, W.: Provenance of lithogenic surface sediments and pathways of riverine suspended matter in the Eastern Mediterranean Sea: evidence from  $^{143}\text{Nd}/^{144}\text{Nd}$  and  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios. *Chem. Geol.* 186, 139–149, doi:10.1016/S0009-2541(01)00415-6, 2002.

- 1295 Weldeab, S., Menke, V. and Schmiedl, G: The pace of East African monsoon evolution during the Holocene, *Geophys. Res. Lett.*, 41, 1724–1732, doi:10.1002/2014GL059361, 2014.
- ~~Weijers, J. W. H., Schouten, S., Hopmans, E. C., Geenevasen, J. A. J., David, O. R. P., Coleman, J. M., Pancost, R. D., Sinninghe Damsté, J. S.: Membrane lipids of mesophilic anaerobic bacteria thriving in peats have typical archaeal traits, *Environ. Microbiol.*, 8, 648–657, doi:10.1111/j.1462-2920.2005.00941.x, 2006.~~
- 1300 [Wendorf, F. and Schild, R.: Prehistory of the Nile Valley, Academic Press, London, 1976.](#)
- Williams, M., Talbot, M., Aharon, P., Abdl Salaam, Y., Williams, F. and Inge Brendeland, K.: Abrupt return of the summer monsoon 15,000 years ago: new supporting evidence from the lower White Nile valley and Lake Albert. *Quat. Sci. Rev.* 25, 2651–2665, doi:10.1016/j.quascirev.2005.07.019, 2006.
- [Williams, M. A. J.: Late Pleistocene and Holocene environments in the Nile basin, \*Glob. Planet. Change\*, 69, 1–15, doi:10.1016/j.gloplacha.2009.07.005, 2009.](#)
- [Williams, M. A. J., Usai, D., Salvatori, S., Williams, F. M., Zerboni, A., Maritan, L. and Linseele, V.: Late Quaternary environments and prehistoric occupation in the lower White Nile valley, central Sudan, \*Quat. Sci. Rev.\*, 130, 72–88, doi:10.1016/j.quascirev.2015.03.007, 2015.](#)
- Zaki, A. S., King, G. E., Hahipour, N., Giegengack, R., Watkins, S. E., Gupta, S., Schuster, M., Khairy, H., Ahmed, S., El-
- 1310 Wakil, M., Eltayeb, S. A., Herman, F. and Castellort, S.: Did increased flooding during the African Humid Period force migration of modern humans from the Nile Valley?, *Quat. Sci. Rev.*, 272, doi:10.1016/j.quascirev.2021.107200, 2021.
- [Ziegler, M., Tuenter, E. and Lourens, L.: The precession phase of the boreal summer monsoon as viewed from the eastern Mediterranean \(ODP Site 968\), \*Quat. Sci. Rev.\*, 29, 1481–1490, doi:10.1016/j.quascirev.2010.03.011, 2010.](#)