

# Accumulation rates over the past 260 years archived in Elbrus ice core, Caucasus

Vladimir Mikhalenko<sup>1</sup>, Stanislav Kutuzov<sup>1,2</sup>, Pavel Toropov<sup>1,3</sup>, Michel Legrand<sup>4,5</sup>, Sergey Sokratov<sup>3</sup>, Gleb Chernyakov<sup>1</sup>, Ivan Lavrentiev<sup>1</sup>, Susanne Preunkert<sup>5</sup>, Anna Kozachek<sup>6</sup>, Mstislav Vorobiev<sup>1</sup>,  
5 Aleksandra Khairedinova<sup>1</sup>, Vladimir Lipenkov<sup>1,6</sup>

<sup>1</sup>Institute of Geography, Russian Academy of Sciences, Moscow, 119017, Russia

<sup>2</sup>Byrd Polar and Climate Research Center, Columbus, OH 43210, USA

<sup>3</sup>Lomonosov Moscow State University, Moscow, 119991, Russia

<sup>4</sup>Laboratoire Interuniversitaire des Systèmes Atmosphériques, Université de Paris and Univ Paris Est Creteil, CNRS, LISA, F-  
10 75013, France

<sup>5</sup>Institut des Géosciences de l'Environnement, Université Grenoble Alpes, 38402 Grenoble, France

<sup>6</sup>Arctic and Antarctic Research Institute, St. Petersburg, 199397, Russia

*Correspondence to:* Stanislav Kutuzov (kutuzov.1@osu.edu)

**Abstract.** In this study, we present a seasonally ~~resolution-resolved~~ accumulation record spanning the period from 1750 to  
15 2009 Common Era (CE), based on a 181.8-m ice core obtained from the Elbrus Western Plateau in the Caucasus. We  
implemented various methods to account for uncertainties associated with glacier flow, layer thinning, and dating.  
Additionally, we ~~developed-applied~~ a novel approach to calculate a seasonal calendar for meteorological data, enabling  
comparison with ice core records. The reconstructed accumulation data were compared with available meteorological data,  
gridded precipitation records, and paleo reanalysis data. Reconstructed accumulation is representative for a large region south  
20 of the Eastern European plain and Black sea region with summer precipitation being the primary driver of precipitation  
variability. ~~Reconstructed accumulation is representative for a large region south of Eastern European plain and Black sea  
region. Summer precipitation was found to be the primary driver of precipitation variability. We identified a statistically  
significant relationship between changes in regional precipitation and fluctuations in the North Atlantic Oscillation (NAO)  
index, which is however not stable over the entire period covered by the ice core.~~ ~~We identified a statistically significant but  
25 unstable in time relationship between changes in precipitation in the region and fluctuations of the North Atlantic Oscillation  
(NAO) index.~~

## 1 Introduction

Reconstructing past precipitation is a crucial aspect of understanding the Earth's climate system, particularly with respect to  
its regional variability. Precipitation plays a critical role in shaping the environment including, water resources, vegetation and  
30 ecosystems. Unlike for most key climate indicators, which show similar trends in most parts of the world, a strong regional  
variation in the sign of trend is observed for changes in precipitation amounts (IPCC, 2014). ~~Unlike most key climate indicators~~

which are changing in the same direction in most parts of the world, there are strong regional variations in the sign of observed changes in precipitation (IPCC, 2014). The spatial distribution of precipitation is much more variable compared to air temperature and a denser network of observations is required to obtain homogeneous data. A particularly large, small-scale variations are large mosaic of precipitation records is observed in mountainous areas due to the complex interaction of circulation factors with the underlying surface. In situ Precipitation-precipitation in situ-measurements can be highly uncertain, particularly especially of for snowfall events. This is often due to gauge under-catch, which can lead to significant errors in the measurement of precipitation amounts (Rasmussen et al., 2012). Currently there are 30 gridded global precipitation data sets available, which include data based on gauge measurements, derived from satellites and reanalysis products Currently there are 30 gridded global precipitation data sets are available including gauge based, satellite related, and reanalysis data sets (Sun et al., 2018). The discrepancies between these data sets highlights their limitation, and the general difficulty, to investigate long-term precipitation changes. In any case, a major drawback is their generally coarse spatial resolution, which is especially problematic in mountain environments where orographic effects play an important role. Despite its discrepancies they are often used for investigating long term climatic changes. However, their major limitation is generally coarse spatial resolution which is especially crucial in mountain environments where orographic effects play important role.

Precipitation variability in the pre-instrumental period is mainly obtained from proxy data and climate models (Bunde et al., 2013; Pauling et al., 2006; Valler et al., 2022). Spore-pollen data characterizing the change of plant communities can be used as indirect evidence of the oscillations of wet and dry periods (Barber et al., 2004; Borisova, 2019). They reproduce the qualitative pattern of climatic variability, but generally suffer from a low temporal resolution generally they have a low temporal resolution-(decades, centuries) and do not always allow quantitative reconstructions.

In contrast, tree rings have an annual resolution and can easily be calibrated using instrumental records. These data are widely used for reconstructions of aridity (Büntgen et al., 2021; Cook et al., 2015, 2020; Solomina et al., 2017) and precipitation (e.g. Touchan et al., 2007; Zhang et al., 2017). The relationship between tree ring parameters of and precipitation strongly depends on the region. Tree-ring width is often a good indirect source of precipitation data in arid conditions, while wood isotope composition can also be closely related to precipitation in some non-arid regions (Loader et al., 2020).

Unlike other proxy, glaciers contain a more direct precipitation signal. Annual layer thickness in ice cores depend on the total annual precipitation amount, although the amount of precipitation may not always be equal to net accumulation. Thus, the most accurate data can be obtained in areas where the loss of deposited snow mass due to melt, sublimation and/or erosion and redistribution by wind and avalanches is minimal. Many polar ice cores were recovered in the central parts of the ice sheets near the ice divides where wind erosion and accumulation of snow is close to equilibriumthere is a balance between snow wind erosion and accumulation. On mountain glaciers, the underestimation of wind-driven snow erosion can lead to significant errors (Bohleber et al., 2013).

To obtain past accumulation rates, the annual-layer thickness has to be corrected for the cumulative effect of layer thinning with depth, which is caused by ice flow.To obtain past accumulation rates, the annual layer thickness has to be corrected for the cumulative effect of ice flow. With the processes of ice flow being well understood, a number of rather simple models and

approaches for calculating the initial thickness of deposited annual layers have been developed over the past decades (e.g. Dansgaard and Johnsen, 1969; Nye, 1963; Paterson and Waddington, 1984; Schwerzmann et al., 2006). The algorithm for calculating the initial thickness of deposited annual layers at the surface is quite well developed (Dansgaard and Johnsen, 1969; Nye, 1963; Paterson and Waddington, 1984; Schwerzmann et al., 2006).

70 In order to reconstruct accumulation from the determined thickness of annual layers, an ice flow model is required to correct for the amount of thinning with depth due to the flow of ice (e.g. Winski et al., 2017). This is particularly challenging for the deepest parts where bedrock topography can become an important factor (e.g. Licciulli et al., 2019). ~~The accuracy of accumulation reconstruction depends on the use of ice flow models to estimate the displacement of the drilling site due to the movement of the glacier and the thinning of the annual horizons, especially for the deep parts of the glacier (Licciulli et al.,~~  
75 ~~2019).~~ For these reasons detailed ice-core reconstructions of accumulation and precipitation are relatively rare both for Polar (e.g. Dahl-Jensen et al., 1993; Goodwin et al., 2016; Henderson et al., 2006; Pohjola et al., 2002; Winstrup et al., 2019; Yao et al., 2008) and High-elevation ice cores (Henderson et al., 2006; Herren et al., 2013; Mariani et al., 2014; Winski et al., 2017; Yao et al., 2008; Zhang et al., 2022) compared to other climate and environmental parameters.

In this paper, we present a high-resolution reconstruction of snow accumulation based on the ice-core records from Mt. Elbrus,  
80 Caucasus and interpretation of the obtained results.

## 2 Data and Methods

### 2.1 Site description

The Caucasus mountain system is situated between the Black and the Caspian seas, and generally trend east-southeast, with the Greater Caucasus range often considered as the divide between Europe and Asia. The 2223 glaciers in the Caucasus cover  
85 an area of  $1060.9 \pm 33.6 \text{ km}^2$  (Tielidze et al., 2022). The Elbrus mountain glaciers contain about 10% of the Caucasus ice volume and cover an area of  $112.6 \text{ km}^2$  (Mikhaleenko, 2020) (Fig. 1a). Glaciers cover an altitudinal range from 2683 to 5642 m asl with the coldest conditions present above 5200 m asl where mean summer air temperature stays below  $0^\circ\text{C}$ . The ice cores used for climate and environment reconstructions were recovered at two sites on Elbrus Mnt.: Western Plateau (WP) at 5115 m asl and the glacier in the crater of the eastern summit of Elbrus at 5600 m asl (Mikhaleenko et al., 2021). A 181.8 m ice  
90 core was recovered at the WP in August-September 2009 and 96 m ice core was drilled the crater in August 2020 at eastern summit (Mikhaleenko et al., 2020). Here we report the results of the WP ice core of 2009.

The WP area is  $\sim 0.5 \text{ km}^2$  and is surrounded by lava ridges to the south and southeast, and by a vertical wall of Mt. Elbrus to the east (Fig. 1b). Ice thickness according to ground-based survey in 2004-2007 varies from 100 to 200 m with a maximum of  $255 \pm 8 \text{ m}$  in the north-eastern part of the plateau (Lavrentiev et al., 2010). 10-m depth temperature is  $-17.3^\circ\text{C}$  (Mikhaleenko et al., 2015). ~~A 181.8 m ice core was recovered at the WP in August-September 2009 and the crater in August 2020 (Mikhaleenko et al., 2020).~~




Figure 1: Location of the Mt Elbrus drilling site: (a) the Caucasus; (b) the Mt Elbrus and meteorological stations; (c) Western Elbrus plateau drill site (photos by I. Lavrentiev, September 2009). ArcGIS World Imagery Basemap used as the background. Source: DigitalGlobe.

The amount of precipitation can be determined as the difference between the measured accumulation layer and the loss caused by melting, sublimation, and wind-driven snow redistribution. Ice formation at the drilling site occurs predominantly in cold, dry conditions and the thickness of the infiltration ice layers, which do not form every year, does not exceed 10 mm (Mikhalenko et al., 2015). The contribution of sublimation rate to glacier mass balance and snow cover balance is estimated to be between 5-10% (Bintanja, 1998; Palm et al., 2017) but can reach up to 30% in certain climatic conditions (Pomeroy and Gray, 1995). On the WP in 2018, sublimation during the short season of possible surface melting was estimated to be 3% of the accumulation layer or  $45 \text{ mm w.e. a}^{-1}$ . Sublimation of ice crystals during snowstorms was measured as 100 mm w.e. or 7% of the accumulation (Mikhalenko, 2020). The contribution of sublimation rate to glacier mass balance and snow cover balance is estimated to be between 5-10% (Bintanja, 1998; Palm et al., 2017) but can reach up to 30% in certain climatic conditions (Pomeroy and Gray, 1995).

Wind-driven snow redistribution was measured by stakes on the plateau during three field seasons, showing a zero balance between scouring-erosion and accumulation of snow near the drilling site (Mikhalenko, 2020). An analysis of the fields of summer and winter accumulation on the plateau from 2015-2017 shows that snow accumulation in the summer period is higher than in winter and is occurs more evenly distributed over the plateau, with its distribution being more stable from year to year. The relocation of winter snow was observed leading to net loss in the southern and western parts of the plateau and net gain on the northern and eastern parts of the plateau limited by the northern ridge and the steep wall of the Western summit of Elbrus In winter, the maximum snow accumulation shows a clear shift to the northern and eastern parts of the plateau, where

120 it is limited by the northern ridge and the steep wall of the Western summit of Elbrus. In the southern and western parts of the plateau, absolute minima are observed in the winter accumulation fields, which are likely caused by strong winds during winter (Lavrentiev et al., 2022). ~~The drill site is roughly in-between these two extremes and can thus be assumed to be in equilibrium also in winter. The area near the drilling site is characterized by mean values of snow accumulation. The total value of snow loss on the WP is estimated to be about 10% (Mikhaleiko, 2020). While we cannot dismiss the possibility of snow accumulation losses driven by wind for winter layers in specific years at the drilling site. Although we cannot rule out the higher snow accumulation losses for winter layers at the drilling site.~~

## 125 2.2 Ice core analysis

The 2009 WP ice core was processed and analysed at the Institute of Environmental Geosciences in Grenoble, (France) for major ions ( $K^+$ ,  $Na^+$ ,  $Ca^{2+}$ ,  $Mg^{2+}$ ,  $NH_4^+$ ,  $SO_4^{2-}$ ,  $NO_3^-$ ,  $Cl^-$ ,  $F^-$ ), succinic acid ( $HOOCCH_2COOH$ ), dust concentration, and black carbon (Kutuzov et al., 2019; Lim et al., 2017; Preunkert et al., 2019). Stable isotopes ( $\delta^{18}O$  and  $\delta D$ ) were analysed using Piccarro at the Arctic and Antarctic Research Institute in St. Petersburg, Russia (Kozachek et al., 2017). Determination of the tritium content was carried out at the University of Bern, Switzerland (Mikhaleiko et al., 2015). 3724 samples were analysed in total down to ~~the~~ 168.6 m depth. Sampling resolution was 10 cm ~~for~~ at the upper part of the ice core and ~~decreased~~ increased to 5 cm at 70 m depth and to 2 cm at 157 depth and below. Discrete sampling in the cold room was performed according to clean sampling protocol (Preunkert and Legrand, 2013). Detailed results of ice-core chemical records are given in (Preunkert et al., 2019).

### 135 2.2.1 Ice core dating

The upper 168.55–6 m (131.566 mwe) depth of the ice core were first dated by annual layer counting primarily using pronounced seasonal variations in ammonium and succinate concentrations, both exhibiting well-marked winter minima (Mikhaleiko et al., 2015; Preunkert et al., 2019). ~~Ammonium can reach the WP (5115 m) as a result of deep convection or with streams of well developed mountain valley circulation. Both phenomena are observed in most cases in the warm half of the year and their intra-annual frequency and intensity are closely related to the seasonality of free convection in the atmosphere. The very low winter  $NH_4^+$  levels are related to precipitation of the cold half year. Specific thresholds were used to differentiate between winter and summer conditions in the upper ice layers, reaching down to a depth of 75.6 meters, corresponding to the year 1963 (Mikhaleiko et al., 2015). These thresholds were set at 100 ppb for ammonium and 5 ppb for succinate concentrations. To account for the~~  
145 ~~observed increase in ammonium concentrations during the industrial era, these criteria were adjusted. Specifically, between depths of 75.6 to 86.8 meters, covering the period from 1950 to 1963, the ammonium winter criteria were modified to 50 ppb, and further reduced to 30 ppb below that depth. In contrast, no substantial depth-related~~

variation was observed for succinate, and the concentration limit of 5 ppb remained consistent in deeper layers. Below a depth of approximately 150 meters, winter layers often consist of only one or two samples, while summer layers consist of more than six samples, making the process of annual dating challenging (Preunkert et al., 2019).


The annual counting was found to be accurate (a 1-year uncertainty) over the last hundred years when anchored with the stratigraphy of the 1912 CE Katmai horizon located at 116.72 m (87.75 mwe) depth (Mikhalenko et al., 2015). Although several other volcanic horizons were suspected to be recorded, particularly below ~~137-139 m (405~~ 106 mwe) depth, none of them was unambiguously attributed to a particular event. A series of sulfate spikes present between 152.58-6 and 154.92 m (118-120 mwe) depth dated at ~ 1840-1833 CE (Table S1), of which two were characterized by an increase in acidity (up to 7.8  $\mu\text{Eq L}^{-1}$ ), were, however, suspected to be possibly related to the 1835 CE Cosigüina~~Coseguina~~ eruption (Nicaragua) (volcanic explosivity index (VEI) of 5, Simkin & Siebert, 1994). It has to be emphasized that given the time period covered by the ice core (1774-2009 CE): ~~S~~surprisingly the layer corresponding to the large eruption of Tambora (1815 CE, VEI of 7, Simkin and Siebert, 1994) was not identified in previous dating of the Elbrus ice core. Uncertainty was not estimated for the dating below the 1912 CE Katmai horizon.

With the aim to estimate uncertainties in ~~dating~~ dating established from annual layers ~~counting~~ deposited prior to 1912 CE, we revisited the chemical profiles including sulfate and acidity. ~~By its nature, d~~Dating using annual layer counting becomes more uncertain with depth because ~~identification~~ layers become thinner resulting from glacier ice flow and sufficient sampling resolution becomes critical to resolve the seasonal variations~~of winter layers is less straightforward due to the decrease of annual layer thicknesses resulting from glacier ice flow~~ (e.g. Paterson and Waddington, 1984). To evaluate the subjective character of the annual layer counting method, four co-authors (denoted A1, A2, A3, and A4) performed the annual layer counting using seasonal variation of ammonium and succinate concentrations without considering a strict criterion. The results were compared to those (denoted A5) previously proposed by Preunkert et al. (2019). As seen in Figure 2, the five chronologies are quite consistent until 131.5 m (100 mwe) depth dated at 1890 CE  $\pm$  3 years and 143.4 m (110 mwe) depth dated at 1866 CE  $\pm$  6 years. Below 149 m ~~The dating uncertainty increases with depth, the layer at 143.48 m (110-115 mwe) depth, A3 and A4 independently counted more years than A1, A2, and A5 by attributing much more ammonium and succinate minima to winter layers (S1). As a result, the layer at~~ and 154.93 m (120 mwe) is dated between 1828 CE and 1842 CE by A1, A2, and A5 whereas A3 and A4 dated it at 1813 CE or 1814 CE. depth being dated at 1867 CE  $\pm$  6 years and 1828 CE  $\pm$  15 years, respectively. A3 and A4 independently counted more years than A1, A2, and A5 by attributing much more ammonium and succinate minima to winter layers (S1). At 168.565 m (131.566 mwe), A1, A2, and A5

proposed an age ranging between 1770 CE and 1783 CE (~~of 1774~~  $CE \pm 6$  years), whereas A3 and A4 proposed an age of 1750-1752 CE  $\pm 4$  years (i.e., a departure of ~25 years).

180 As detailed in the SI, examination of the records of acidity and sulfate (Figure S2) permitted to evaluate the extent to which the two chronologies (A1-A2-A5 versus A3-A4) are consistent with volcanic horizons and from that dating uncertainties. The A3-A4 ~~Using this new dating we~~ attributed layer of increased acidity and sulfate located at 112.57 mwe (peak c dated at 1851 CE) to the 1854 CE Shiveluch eruption (Kamchatka) (~~of which the volcanic~~  
185 mwe (peaks d and f) ~~was were~~ previously identified as possibly related to the 1835 CE Cosigüina eruption (A5) are now attributed by A3-A4 to the 1815 Tambora eruption (VEI of 7, Simkin and Siebert, 1994). The layer dated at 1789 CE (peak e) may be related to the 1783 CE Laki eruption (see further discussions in supplementary).

~~In conclusion, The the WP two chronologies are consistent within one or two years back to the horizon of Shiveluch in 1854.~~ In conclusion, the A1-A2-A3 and A3-A4 chronologies are consistent until 131.5 m (100 mwe) depth (i.e.,  
190 1890 CE  $\pm 3$  years) but differ ~~Further down, dating differ~~ by some 25 years at 130 ~~131.9-6~~ mwe (1750-1752 CE  $\pm 4$  years instead of 1774-1776 CE  $\pm 6$  years), ~~depending on the results of the annual layer counting.~~ The new A3-A4  
195 datings (1750-1752 CE  $\pm 4$  years) is-are consistent with a Tambora layer located at 118.96 or 119.84 mwe and possibly a Laki layer at 124.71 mwe. Given the expected larger impact of the Tambora eruption compared to that of Cosigüina, we give more credit to the A3-A4 chronology. Note also that the u  
Use of the two possible datings (A1-A2-A5 and A3-A4) only slightly affects the general trends and tendencies of the accumulation reconstruction (S4). In this study, the new dating will be used as the basis for the accumulation reconstruction.-



200 **Figure 2. Age-depth (in mwe) relation of the Elbrus ice core derived by annual layer counting achieved by 4 co-authors of the present paper (denoted A1, A2, A3, and A4) and by Preunkert et al. (2019) (denoted A5).**

## 2.3 Snow accumulation reconstruction

### 2.3.1 Thinning of annual layers due to ice flow

205 The accumulation rate history at Mt. Elbrus can be inferred from depth profiles of annual-layer thicknesses in the WP ice core when corrected for ~~firn densification and~~ thinning of ice layers due to ice flow. Density measurements, as reported by Mikhalenko et al. (2015), reveal kinks in the curves of the studied ice cores, corresponding to critical density values of 550 and 840 kg m<sup>-3</sup> (Maeno and Ebinuma, 1983). The third critical density value of 730 kg m<sup>-3</sup>, which marks the transition of firn into ice by complete closure of air inclusions, is not clearly visible on the density curve. This indicates ice formation without meltwater involvement (Hörhold et al., 2011; Ligtenberg et al., 2011). The depth at which air bubbles separate from the surrounding ice matrix and pore closure occurs is approximately 55 m, with a measured density of about 840 kg m<sup>-3</sup> (Mikhalenko et al., 2015).

210 ~~Although the annual layer thickness exhibits high variability, the data suggest that layer thinning occurs with increasing depth due to ice flow. To reconstruct initial accumulation values from the annual layers' water equivalents as observed in the ice~~



215 ~~core we used To determine snow accumulation values, we used~~ a simple J. Nye flow model (Nye, 1963) (~~Dansgaard and Johnsen, 1969~~) incorporating mean accumulation rate and constant glacier thickness over the time span represented by an ice core (Dansgaard and Johnsen, 1969):

$$h_r = h_o \times e^{\frac{h_o \times A}{H}}, \quad (1)$$

$$h_o = \frac{H \times W\left(\frac{A \times h_r}{H}\right)}{A}, \quad (2)$$

220 where  $h_r$  and  $h_o$  – recent and original depths of annual layer respectively (m),  $A$  – age (years),  $H$  – depth of glacier (157 m ice eq.),  $W$  – Lambert W function.

Scaling of the best fit curve and the residuals of the raw data from the best fit curve (S3a) to the linear mean accumulation value (1.5 m ice eq.) was considered as the representation of the variability of the “original” annual layers’ depths ~~as if no variability in accumulation over the glacier surface was expected~~ (S3b).

### 225 2.3.2 Correction for the upstream effect ~~Calculation of backward trajectories~~


Since the ice core includes layers that were deposited upstream of the drilling site, where annual snow accumulation conditions may differ from those at the drilling site, it is crucial to identify the ice core catchment area in order to investigate the connection between the ice core data and surface accumulation. To ~~verify the representativeness of the ice core for the accumulation conditions at the drilling site and~~ account for upstream effect, we reconstructed the backward trajectories of the ice flow. To  
230 this end, we calculated the backward trajectories of the ice/firn particles positioned along a vertical line segment connecting the drill site and the bedrock (Fig. 3). In the post-processing step, the backward trajectories were calculated by utilizing the Runge-Kutta integrator implemented in the stream tracer of the ParaView visualization application, based on the modeled glacier velocity field. (Ahrens et al., 2005). The velocity field is simulated based on a 3-D full Stokes ice-flow model with the firn rheological law (Gagliardini and Meyssonier, 1997). The model is implemented using the finite element software  
235 Elmer/Ice (Gagliardini et al., 2013). We performed a steady-state simulation with fixed glacier geometry.

The mathematical formulation of the ice-flow problem follows (Zwinger et al., 2007) with some simplifications and includes the Stokes and the volume balance evolution equations, the stress-free surface boundary condition and the no-slip bedrock boundary condition. Unlike (Zwinger et al., 2007), we used only dynamical equations and did not consider thermo-mechanical coupling. Therefore, the flow rate factor in the firn rheological law considered constant. The model ~~used~~ utilizes an  
240 approximation smoothed vertical profile of the density profile measured in ~~inferred from the 2009 WP ice core the borehole drilled in 2009~~ (Mikhalenko et al., 2015). Model simulations show that ~~constant the~~ value of the flow rate ~~factor in the firn rheological law~~ has very little effect on the geometry of the backward trajectories and the position of their sources on the glacier surface. Therefore, the rate factor may be chosen arbitrary from a wide range of values (from 4.82 MPa<sup>-3</sup>a<sup>-1</sup> to 14.46 MPa<sup>-3</sup>a<sup>-1</sup> in our computational experiments).

245 Since the digital elevation model data (Lavrentiev et al., 2022) covered only part of the glacier bedrock, the computational domain was restricted to this area, and the lateral boundary connecting the surface and the bedrock boundaries was a fragment of a cylindrical surface (vertical wall) (Fig. 3c). As most of the lateral boundary was not a physical boundary in the glacier, the flow of ice/firn through it was possible. We considered two alternative boundary conditions for the lateral area: (i) no outflow (zero horizontal velocity) at the whole lateral boundary, and (ii) constant nonzero outflow velocity at the western and southern parts of the lateral boundary and no outflow at the eastern side where a vertical wall of Mt. Elbrus is situated (Fig. 3a). In case (ii), the outflow velocity vectors were directed outward from the computational domain along the normal to its lateral surface. The outflow velocity value was considered constant in space and time.

250 Our simulation showed that the geometry and the length of the backward trajectories were influenced by the presence or absence of the outflow rather than the particular value of nonzero outflow velocity. Therefore, this velocity value was not an essential parameter in our modeling. Additionally, no field data on velocities were available.

255 In our model experiments, we obtained the backward trajectories of maximal length under conditions (ii). Since the maximum extent of the ice core catchment area was of interest in our study, we present the results of this model case. We used the flow rate factor  $B = 4.82 \text{ MPa}^{-3} \text{ a}^{-1}$  and the outflow velocity  $v_{\text{out}} = 1 \text{ m a}^{-1}$  in the case illustrated in Figure 3.



260 **Figure 3.** Accumulation map for the Western plateau of Elbrus in 2016–2017 CE (Lavrentiev et al., 2022) (a); the modeled catchment area for the 2009 CE ice core (b); and the computational domain (c) with the borehole position (white line) and the calculated backward trajectories (red lines). An orange line shows the outflow zone. The source points of the backward trajectories are marked with red dots; the corresponding borehole depths (rounded, in m) according to DEM (2017) are given as dots labels. The green triangle indicates the position of the ice core (2009).

265

Based on our simulation, we have determined that the sources of the backward trajectories are located on the glacier surface, no more than 140 meters northeast from the ~~drill site used in 2009 CE~~ drill site. To estimate the correction factor for the difference in snow accumulation at the initial source location (S5), we utilized average accumulation distribution maps, which were generated using a combination of ice core data and a high-frequency radar survey (Lavrentiev et al. in 2022). Snow accumulation increases linearly with distance from the drill site along the backward trajectory. Therefore, we applied the same linear relationship to calculate the accumulation correction (%) based on the ice age. The most significant correction of 30% was applied to the oldest layers dating back to 1750 CE (S6).

## 2.4 Climate data

Data from 16 weather stations located in the foothill or mountainous regions of the Caucasus (Russia and Georgia) were used for the statistical estimates of the precipitation field in the North Caucasus, as well as for comparing the time series of precipitation with snow accumulation on the WP drilling site (Table 1). Most of the stations are located in lowland and foothill areas (less than 1000 m asl), 5 - in the middle mountain zone, and 4 - in high mountains (more than 2000 m asl). The CAPE (convective available potential energy) was calculated for separating summer and winter seasons using the ERA5 reanalysis data. We also used [the](#) Global Precipitation Climatology Centre (GPCC) v2020, 0.25° monthly precipitation dataset from 1891-present calculated from global station data for field correlation analysis.

**Table 1. Meteorological data used for analysis**

Weather station or reanalysis	Distance from drilling site, km	<u>Altitude</u> H, m asl	Coordinates	Observation period	<u>ValuesParameters</u>
Terskol (RUS)	12	2350	42.73N; 42.33E	1951–2010	P
Klukhorskiy pereval (RUS)	50	1815	43.15N; 41.52E	1965–2010	P
Sulak (RUS)	250	2927	42.22N; 46.25E	1936–2010	P
Teberda (RUS)	60	1325	43.27N; 41.25E	1926–2010	P
Shadjatmaz (RUS)	45	2056	43.7N; 42.40E	1961–2010	P
Kislovodsk (RUS)	75	860	43.90N; 42.50E	1936–2010	P
Mineralnie Vody (RUS)	100	420	44.10N; 43.10E	1955–2010	P
Krasnaya polyana (RUS)	170	960	43.60N; 40.20E	1936–2010	P
Sochi (RUS)	190	50	43.50N; 39.60E	1875–2010	P
Zelenchukskaya (RUS)	70	925	43.90N; 41.50E	1960–2010	P
Vladikavkaz (RUS)	170	750	43.00N; 44.60E	1951–2010	P
Mestia (GEO)	40	1445	43.00N; 42.60E	1961–2010	P
Kutaisi (GEO)	105	260	43.20N; 42.60E	1936–2010	P
Zugdidi (GEO)	125	160	42.50N; 41.90E	1961–2010	P

Ambolauri (GEO)	85	544	42.50N; 43.00E	1936–2010	P
Pasanauri (GEO)	130	1070	42.40N; 44.60E	1936–2010	P
Reanalysis CFSR	30 *			1979–2010	T, E, f, q **
Reanalysis ERA5	15 **			1979–2010	CAPE ***
GPCC				1891-2010	P

285 \* The distance between the drilling site and the nearest node of the reanalysis grid.

\*\* P-precipitation, T - temperature, E - vapor saturation pressure, f - relative humidity, q - cloud air mixture ratio

\*\*\* CAPE - convective available potential energy

RUS – Russia

GEO - Georgia

## 290 2.4.1 Seasonal calendar

A direct comparison of the ice-core reconstructed net accumulation/precipitation records with precipitation amounts measured at meteorological stations contains seasonal and annual dating uncertainty. The boundaries of warm and cold seasons at meteorological stations can vary significantly depending on the geographical location and altitude. Seasonal variations of ammonium for which the main source is vegetation in the active phenological phase corresponding to the warm half-year (April-September) were used for identification winter and summer layers in WP ice core. Ammonium concentration in the ice core is maximum during the period of active convection and the accumulation sum over this period corresponds to precipitation of the warm half of the year.

As a seasonality predictor, we used the convective available potential energy (CAPE) calculated from the ERA-Interim reanalysis data for 1979–2010 which has a pronounced seasonal variation and reflects convective movements in the atmosphere due to buoyancy forces. CAPE depends on the difference between the virtual temperature of the air particle ( $T_i$ ) and the virtual temperature of the air surrounding the particle ( $T$ ). Integrating this difference along the vertical ( $z$ ) and multiplying it by the gravitational acceleration  $g$ , we obtain the convective instability energy in  $\text{Jkg}^{-1}$ .

$$\int_{z_1}^{z_2} g \left( \frac{T_i - T}{T} \right) dz \quad (3)$$

305

A number of studies (Chen et al., 2008; Markowski and Richardson, 2010) formulate gradations of CAPE that correspond to a particular level of convective atmospheric instability. Most of these studies are aimed at identifying the relationship between the recurrence of hazardous weather events of convective origin and the CAPE value. For example, in (Chen et al., 2020) it is shown that most of the weather hazards in the temperate US climate zone occur when the CAPE is between 260-840  $\text{Jkg}^{-1}$ . Under these conditions there is a high probability of deep convection development that can transport biogenic particles and chemical ions from the ground layer of the atmosphere to the 5000 m level. At Caucasus latitudes the CAPE may exceed the

310

value of  $300 \text{ Jkg}^{-1}$  even in the cold half of the year, on some exceptional days but not systematically. We therefore assume that a warm half-year in the North Caucasus begins when the CAPE exceeds  $300 \text{ Jkg}^{-1}$  at least once every five days, i.e., during the natural synoptic period. Analysis of the annual variations of CAPE over the Caucasus (S7) shows CAPE is consistently less than  $300 \text{ Jkg}^{-1}$  in summer in the region only under long and pronounced anticyclonic conditions corresponding to blocking. Blocking anticyclones are characteristic of more northern latitudes, with blocking developing much less frequently in the subtropics and in the southern temperate climate zone (Bardin et al., 2019). In addition to deep convection, the advection of ammonium ions to the WP level may be regulated by the mountain-valley circulation, which is also most active in the summer months and is indirectly related to the CAPE value

The CAPE value depends on the heat content of the lower atmospheric layers and the intensity of heating of the underlying surface, so it has a pronounced seasonal variation. In spite of significant climate changes in the Caucasus, the CAPE value, which is one of the thermal characteristics of the atmosphere, did not show statistically significant increase neither in the duration of the warm period (number of days) nor in the seasonal mean value. For the period covered by the reanalysis data (1979-2015), a calendar of warm and cold seasons was compiled (S8).

## 3 Results and discussion

### 3.1 Net accumulation reconstruction

The graphical results of the accumulation reconstruction are presented in Figure 4. The uncertainty range for the reconstructed accumulation values is determined by multiple factors. Dating uncertainty varies with time, providing a precision of  $\pm 1$  year for the 2009-1912 CE period,  $\pm 2$  years for the period 1912-1825 CE, and decreasing to  $\pm 4$  years for the period 1825-1750 CE. Additionally, the uncertainty linked to the upstream effect correction is challenging to estimate precisely, depending on factors such as initial spatial accumulation distribution uncertainty, assumptions regarding its temporal persistence, and uncertainty in back trajectory modeling. To address potential errors, we estimated the uncertainty of the applied linear regression. Furthermore, for winter accumulation, the uncertainty increases due to lower sampling resolution. In particular, we added an uncertainty of 30% for winter accumulation values prior to 1865 CE, where the sampling resolution was insufficient, potentially leading to an underestimation of accumulation values. ~~The dating accuracy of the reconstruction varies, with a precision of  $\pm 1$  year for the years 2009–1912 CE,  $\pm 2$  years in the period 1912–1825 CE, and a decrease to  $\pm 4$  years in the period 1825–1750 CE for accumulation was.~~

The mean annual accumulation reconstructed at the WP from 1750 to 2009 CE is 1.2 m.w.e. During this same period, the mean summer and winter accumulations are 0.8 m.w.e and 0.4 m.w.e, respectively. Due to dating uncertainties this record is suitable for investigations of decadal, multidecadal and long term regional precipitation variations rather than interpretations of the accumulation for the exact years.




No statistically significant trends were observed in 20<sup>th</sup> century in WP. The record can be classified into five major periods based on accumulation amount and year-to-year variability. The period before 18320 was characterized by relatively high

~~accumulation and a high annual variability~~ increased summer accumulation and annual variability. This was followed by the relatively low summer and winter accumulation of the 1832-1860 CE period with less pronounced variations. From 1860 to 1935 CE, summer and winter accumulation increased by around a factor of two and strongly fluctuated. The period from 1935 to 1980 CE was again characterized by relatively low accumulation in summer and relatively high accumulation in winter. ~~was characterized by low accumulation in summer and relatively high accumulation in winter. Since around 1980 CE, an increase in summer and winter accumulation has been observed.~~

~~No statistically significant trends were observed in 20<sup>th</sup> century in WP. A slight positive trend (0.018 m w.e. per decade) was estimated for the annual accumulation over the record, attributable to a general increase in winter accumulation. However, this trend is likely due to the insufficient sampling resolution of the deepest layers, which failed to fully capture the winter layers for a certain year and resulted in an underestimation of winter accumulation below 110 mwe depth (corresponding to the year 1865 CE).~~

~~The record can be classified into five major periods based on accumulation amount and year to year variability. The period before 1830 was characterized by increased summer accumulation and annual variability. This was followed by the relatively low summer and winter accumulation of the 1830-1860 CE period with less pronounced variations. From 1860 to 1935 CE, summer and winter accumulation increased by a factor of two and strongly fluctuated. The period from 1935 to 1980 CE was characterized by low accumulation in summer and relatively high accumulation in winter. Since 1980 CE, an increase in summer and winter accumulation has been observed.~~

The WP site exhibits a seasonal distribution of precipitation typical of the Central Caucasus, with convective precipitation leading to a maximum in the summer months (Mikhalenko et al., 2020). ~~As discussed above winter accumulation before 1865 CE (110 m w.e.) is likely underestimated due to insufficient sampling resolution.~~ The mean share of summer accumulation in the total annual accumulation was 70.66% (STD=18) over the course of 144.260 years (1865-2009 CE), which is consistent with current measured precipitation data at weather stations (S9). ~~Over the entire period covered by the core data, there is a statistically significant decrease in the contribution of summer accumulation to the annual total ( $R^2 = 0.6$ ). The highest percentage of summer accumulation occurred between 1750 and 1830 CE, with an average contribution of 83% to the annual total, resulting in an anomaly of over 200% compared to the modern level in some years. In contrast, the lowest percentage of the summer component of the annual accumulation was observed in 1935-1980 (57%) and after 2004. This period was characterized by a prolonged negative anomaly of summer accumulation (Fig. 4b), the average value of which was -25%, surpassing the natural interannual variability of  $\pm 20\%$  for the Central Caucasus (Toropov et al., 2019).~~



375 **Figure. 4. Reconstructed annual (a), summer (b) and winter (c) snow accumulation at the WP. 10-year moving averages are shown in thick lines. Uncertainties are shown as shaded areas.**

### **3.2 Comparison with meteorological data**


#### **3.2.1 Spatial distribution of precipitation in the North Caucasus**

The interpretation of the reconstruction of accumulation relies on a comparison with meteorological information gathered during the period of instrumental observations, utilizing weather station data, global precipitation data sets and reanalyses.

380 This comparison enables us to assess the accuracy of annual and seasonal accumulation amounts, as well as to determine the area where this reconstruction holds significance for different timescales. Additionally, we can establish the statistical and physical relationship between resulting accumulation and precipitation amounts, while mitigating the potential impact of confounding factors such as wind drift, avalanche supply, evaporation, and sublimation rates.

385 The spatial correlation function for the observed annual precipitation amount was calculated relative to the Terskol meteorological station closest to the drilling site. The spatial distribution of the normalized correlation coefficients between the series of annual precipitation in Terskol and at other stations in the Caucasus is shown in Fig. 5.






390 **Figure 5. Spatial correlation function of the annual precipitation calculated relative to the Terskol meteorological station (red dot) (a): isolines of correlation coefficients calculated according to Student's t-test are given by white lines; dashed lines indicate the profiles for which the correlation function was calculated. Correlation function values along these profiles (b). ArcGIS World Imagery Basemap used as the background. Source: DigitalGlobe.**

The significance of the normalized correlation coefficient was assessed using Student's t-test, with a threshold value of 0.4. However, as a correlation coefficient exceeding this value not only has statistical but also physical meaning, the threshold was increased to 0.6. Figure 5 demonstrates that the precipitation field in the Caucasus is significantly isotropic, with correlation coefficient isolines appearing as elliptical shapes stretching along the Greater Caucasus. The correlation radius is larger in the WNW direction, owing to the influence of western air mass transport and the Black Sea, compared to the ESE, S, and N directions, as moisture-bearing air masses rarely reach the Greater Caucasus from the north. Consequently, annual year-to-year variability of precipitation in the northern foothills may differ from the precipitation in the high-mountain regions of the Caucasus. Despite the challenging orographic conditions, the radius of statistically significant correlation of annual precipitation ( $r > 0.4$ ) measured at the weather stations encompasses the majority of the Western and Central Caucasus, while a physically significant radius ( $r > 0.6$ ) extends to 50–100 km. Similar findings were reported by (Tashilova et al., 2019). The

highest correlation ( $r \geq 0.8$ ) occurs within an area of 10-50 km, ~~indicating that on the annual and sub-annual timescale ice core records likely provide description of precipitation within 50-100 km radius.~~

405 Statistical analysis of long-term series of annual precipitation (Sochi, Teberda, ~~Krasnaya Polyana~~) and reconstructed accumulation on WP reveal high similarity of empirical distribution functions (Fig. 6), which are asymmetric log-normal distributions (Fig. 6 b, c) shifted towards positive anomalies and decreasing sharply towards negative ones. The median value corresponds to a positive anomaly of 10-15%, which is typical for time series of precipitation. The root mean square (RMS) deviation of both precipitation and reconstructed accumulation is approximately 20%, which is typical for annual precipitation in temperate climatic zones. The range of variability of precipitation anomalies, based on averaged weather station data, ranges  
410 from a decrease of 50% to an increase of 80%. ~~In contrast, ice-core records show larger range of variability. This difference can be attributed to the length of the ice-core data, which spans 260 years, significantly longer than the longest continuous meteorological records (since 1875 CE, Sochi). Consequently, the ice-core data captures the full range of precipitation variability in the Caucasus, ranging from extremely dry conditions (one fifth of annual precipitation) to exceptionally wet conditions (double the annual average).~~



415


**Figure 6. Statistical analysis of the data time series on the longest-range weather stations and ice cores: (a) spectral analysis of the series, (b) empirical distribution function of annual precipitation anomalies compared to the period ~~1981-1951-2010-1980~~, averaged between data from the longest-row weather stations Sochi, ~~Krasnaya Polyana~~, Teberda and ~~Sulak-Vysokogornaya~~, (c) empirical**

420 **distribution function of annual precipitation anomaly on the West Elbrus plateau according to ice core data compared to the period 1951-1980.**

Spectral analysis showed that according to the accumulation data and the results of meteorological observations, the statistically significant peak of the spectral density function for all data series (ice core, Sochi, Teberda) falls on a period of 8-  
425 10 years, and 30-40 years. The second period is close to the characteristic scale of the North Atlantic Oscillation (NAO) and Atlantic Multi-decadal Oscillation (AMO) index fluctuations. The similarity of the distribution functions and spectral density from the data of weather stations and ice-corer records indicates that accumulation at Elbrus is primarily related to precipitation and not significantly affected by post-depositional factors (e.g. melt, wind erosion etc.). At least for temporal averages >5 years, the reconstructed record is dominated by the precipitation signal.  
430 ~~indicates that the reconstructed accumulation values at Elbrus are primarily related to precipitation rather than to other accumulation factors. In some years, they may play a decisive role, but at averaging scales of more than 5 years, the precipitation factor plays the predominant role.~~

### 3.2.2 Seasonal accumulation

Over the past 47 years, the period of summer identified by CAPE typically spans from May 1 to October 10, which nearly aligns with the hydrological year. However, our calendar displays considerable inter annual variability (Table S8). The earliest  
435 sustained transition through  $300 \text{ Jkg}^{-1}$  occurred on April 1, 2008, while the latest was on May 27, 2003. The earliest transition to winter convective instability took place on September 17, 1986, and the latest on November 1, 2005. The minimum period with steady CAPE  $> 300 \text{ Jkg}^{-1}$  was 135 days (in 2010), the maximum was 200 days (in 1995 and 2005), and the average was 164 days. The distribution of the warm period's duration by CAPE is close to normal, but at the  $2\sigma$  level, the deviation is  $\pm 1$  month. Therefore, accounting for this factor in determining seasonal ~~and annual~~ precipitation amounts is critical. ~~Failure to do~~  
440 ~~so can result in errors exceeding 10% of annual precipitation sums, as illustrated in Figure 7. This figure~~Figure 7 displays regression relations of the accumulation reconstructed from the core with precipitation amounts at the nearest weather stations before and after the use of CAPE calendar. for the 1979-2009 CE period.




**Figure 7. Regression relations between the accumulation layer on the West Elbrus plateau ( $\Sigma P_{core}$ , mm) and precipitation amounts at the nearest weather stations ( $\Sigma P_{ws}$ , mm) during 1979-2009 CE period: (a-d) for the cold season (a, c for October-March calendar period, b, d for the cold period highlighted by CAPE value thresholds); (e-h) for the warm season (e, f) for the April-September calendar period, (g, h) for the warm period highlighted by CAPE values).**

It is well seen that the  $R^2$  in the relations between precipitation and accumulation in the ice core with-when considering CAPE dates increases, and become statistically significant (more than 0.4,  $p < 0.001$ ). This indicates that accounting for the role of free convection (~~CAPE~~) significantly improves the relationship between accumulation on the WP and precipitation measured at weather stations. The improvement is most significant for the winter season. This is due to the greater spatial homogeneity of the precipitation field in the cold half of the year, which is expressed in larger values of the radii of significant correlation. In summer likely due to a significant contribution of local convective precipitation the improvement is not so significant. To facilitate further analysis, we have defined the cold period as the period from October to March and the warm period as the period from April to September.

### 3.2.3 Comparison with gridded data

Spatial comparison with the gridded GPCC v2020 0.25 precipitation dataset revealed a statistically significant correlation for both seasons between ice core accumulation and precipitation estimates (Fig. 8). Summer accumulation moderately correlates with precipitation observations over Ukraine, South Russia, and the Black Sea. Winter accumulation shows an even stronger correlation with gridded datasets over the entire Black Sea region and North Caucasus. We also found a strong correlation between winter snow accumulation at the WP and observed (GPCC) March precipitation over the North Caucasus, with the highest coefficients (0.5) concentrated close to Elbrus (S10). According to Terskol weather station data, around 30% of the December-March precipitation falls in March, which is likely reflected in the higher share of March snow in cold season accumulation recorded at WP. ~~The snow deposited in March is usually of higher density and is less prone to wind removal.~~



**Figure 8. Spatial correlation of the summer (a) (April-September) and winter (b) (October-March) accumulation with precipitation data of the GPCC v2020 data set in 1950-2009 period. Location of the Elbrus Mt. is shown by black triangle.**

470 The correlation between seasonal accumulation and gridded precipitation data GPCC v2020 0.25 remains below 0.4 for the majority of the territory. However, during the cold period in a substantial area of the North Caucasus steppe zone, the correlation significantly exceeds 0.4, reaching values ranging from 0.6 to 0.65.


475 Analysis of spatial correlation of gridded precipitation data and meteorological measurements at Terskol station further support our interpretation of the ice core record. Monthly correlation fields show that precipitation pattern is more homogenous in autumn and winter months, and area of correlation expands from east Europe to Turkey and Southern Russia. In summer (May-August) the correlation radii is reduced to on only 100-200 km (S11). Ice core data captured these patterns. A plausible explanation for the strong correlation observed between winter accumulation on the WP and precipitation levels in the foothill areas of the North Caucasus could be attributed to the relatively frequent occurrence of orographic occlusion (Mikhalenko et

al, 2020). This meteorological phenomenon arises when a wave of cold temperate or Arctic air interacts with the Caucasus Mountains. It commonly transpires in the rear sector of Mediterranean or Black Sea cyclones, or behind an Atlantic vortex that has shifted from Scandinavia to the Volga region. The presence of the Caucasus ridges impedes the progress of cold air, causing it to circumnavigate the mountains from the direction of the Black and Caspian Seas, thereby saturating the Transcaucasus region. Consequently, the warm and moist air is forced upwards towards the mountains, leading to an extended release of moisture. During summer, this process often manifests in the formation of localized cumulonimbus clouds, while in winter, it results in substantial frontal precipitation across a significant expanse. These dynamics are most pronounced in the high mountain zone and the foothill areas of the Central Caucasus, offering an explanation for the correlation observed between winter accumulation on the WP and precipitation in the steppe regions of the northern foothills.


### 3.3 Comparison with climate indexes

In many parts of Europe, quasi-decadal variations in atmospheric precipitation are associated with the internal nonlinear dynamics of the climate system. This is reflected in the interannual variability of indices such as NAO, AMO, East Atlantic/Western Russia teleconnection pattern (EA/RW), and others in temperate latitudes (e.g., Trigo et al., 2002; Hurrell and Deser, 2009; López-Moreno et al., 2011; Ionita, 2014).

Negative NAO values often correspond to a weakening of cyclogenesis over Iceland and the formation of anticyclones over northern Europe, which is atypical for the North Atlantic. As a result, cyclogenesis at the polar front over the Mediterranean and Black Seas becomes more active, leading to positive precipitation anomalies in the Caucasus. Such a large-scale circulation anomaly and the negative phase of the NAO and AMO prevailed during the cold seasons of the 1960-1970 CE. The positive winter precipitation anomaly recorded in the WP ice core reflects the large-scale pattern of precipitation anomalies. In 1960-1970 CE, a pronounced region of statistically significant precipitation anomalies covered the whole of the Mediterranean and most of the southern part of Eastern Europe. The value of anomalies reached 15-20 mm per month or 20-30 % of the seasonal amount for the period October-March (Fig. 9a). They corresponded to predominantly negative geopotential anomalies in the middle troposphere, particularly over Western Europe (Fig. 9b). Northern Europe experienced an extensive precipitation deficit due to a pronounced positive geopotential anomaly over the North Atlantic.



505 **Figure 9** An average winter precipitation anomalies in 1961 – 1970 (GPCC) compared to 1981 – 2010 (a) and average 500 hPa isobaric surface height anomalies (NCEP/NCAR) (b).



510 **Figure 10.** Spatial correlation of the detrended winter (October-March) annual accumulation at WP with ERA5 500 mb geopotential height data (a) and sea surface temperature (HadISST1) data in 1950-2009 (b). Ten-year moving average of winter

NAO (October-March) and detrended ice core winter accumulation anomalies (c). Location of the Elbrus Mt. is shown by black triangle.

515 The influence of the North Atlantic circulation on the cold season precipitation over the region is highlighted by the correlation of the WP winter accumulation with the Sea surface temperature and geopotential height anomalies over the North Atlantic (Fig. 9a, b). For the cold period (October-March) in 1930-2009 CE, the 10-year moving averaged WP accumulation anomalies show a statistically significant negative correlation with the NAO index ( $r=-0.74$ ,  $p<0.001$ ) (Fig. 10c).

520 A relationship between precipitation regime and the NAO index has been shown in previous studies (Deser et al., 2017; Vicente-Serrano and López-Moreno, 2008), particularly during winter months. Positive correlation coefficients between NAO and precipitation sums are characteristic for Northern Europe, while negative correlations are observed for the Mediterranean region and southern Europe, which is also evident in our case. Furthermore, a statistically significant inverse relationship between winter precipitation and the NAO index in Turkey (~~a region relatively close to the Central Caucasus~~) was revealed in the study by Türkeş and Ecmel (2005), with a coefficient of -0.4 over a significant portion of the territory. The substantial role of NAO in shaping the precipitation regime of southern Europe and Turkey ~~is~~ was also demonstrated using global circulation  
525 models for 15 mountain areas in Mediterranean in the work by (López-Moreno et al., 2011).~~,-~~

Opposite to the last ~80 years where the correlation is negative, a moderate positive correlation between NAO and accumulation in the cold season was found for the earlier period from 1880-1925 ( $r=0.6$ ,  $p < 0.001$ ). This period is also characterized by increased summer accumulation and generally high annual precipitation variability. The variation of the correlation between the reconstructed accumulation for Elbrus and the NAO index is in line with previous findings of  
530 instabilities in the connection of precipitation and large-scale atmospheric circulation at decadal timescales observed over southern and central Europe (Pauling et al., 2006).

We did not find any significant relationships between summer accumulation records and climate indexes. In summer, the NAO is significantly weaker due to the greater role of mesoscale processes in the formation of seasonal precipitation, as well as the proximity of the Black Sea. In particular, the increase in precipitation and the related rise in summer accumulation during  
535 2000-2009 may be a response to the increase in sea surface temperature in the eastern part of the Black Sea which could be a cause of the observed precipitation increase in southern Russia (Aleshina et al., 2018). The role of positive sea surface temperature anomalies in the Black Sea in the formation of synoptic situations leading to extreme precipitation in the Western Caucasus is clearly demonstrated in the study by (Meredith et al., 2015). Additionally, the increase in summer accumulation in the first decade of the 21st century is generally consistent with the overall positive trend of convective precipitation  
540 (Chernokulsky et al., 2019).

Our findings suggest that heat transport in the North Atlantic is of critical importance in determining the precipitation regime in cold season over the Black Sea and Northern Caucasus.



### 3.4 Comparison with paleo records

Reconstructions of the accumulation and precipitation from ice cores from mountain glaciers are relatively limited compared to the polar regions. Thus, in the Alps, the main problem is the influence of avalanche feeding and significant blizzard redistribution of snow at the drilling sites (Bohleber, 2019).

Several estimates of past precipitation were published for the Caucasus region. Martin-Benito et al. (2016) reconstructed May-June precipitation in Transcaucasia since 1752 based on the three ring width. Their reconstruction explains 51.2% of the variability of instrumental data. Verhaegen et al. (2020) used this reconstruction to calculate the annual amount of precipitation in Terskol. The analysis of precipitation frequency, magnitude, seasonality, as well as the source areas and trajectories of the air masses involved in the Colchis Lowland and near Mt. Elbrus, reveals substantial differences in the precipitation regimes of these regions. Consequently, the reliability of this reconstruction is questionable. Tree-ring width based reconstruction of spring (April–July) precipitation in Crimea over 1620–2002 was published (Solomina et al., 2005). The reconstruction accounts for 37% of the variance in observed precipitation over 1896–1988 although the difference in season duration does not allow the direct comparison with WP accumulation record.

We compared WP accumulation reconstruction with the available monthly resolved paleo-reanalysis EKF400 version 2 with a 2x2 degree resolution, covering the period 1603 to 2003 (Valler et al., 2022). In the EKF400 Version 2, the Kalman filtering technique was utilized to couple an ensemble of atmospheric general circulation models for assimilating early instrumental observations of temperature, surface pressure, and precipitation, in addition to temperature indices derived from historical documents and tree-ring measurements that are sensitive to temperature and moisture.


We observed a robust correlation between 10-year averaged summer and winter accumulation data and precipitation data from the EKF400 Version 2 (Fig. 4011). The region exhibiting a stronger correlation is situated to the south of the Eastern European plain for both summer and winter seasons. Which is consistent with the pattern observed for the spatial correlation analysis in Figure 8 and S11. This correlation remains consistently strong throughout the instrumental observational period since 1850 CE. However, prior to 1850 CE, the datasets do not align. The summer paleo precipitation records for the region display an unexpected and unsupported strong negative trend, with a sharp decrease from 170 mm a<sup>-1</sup> (1750-1830CE) to ~120 mm a<sup>-1</sup> (after 1830 CE). Other records do not corroborate support this such trend sharp change in precipitation and moisture neither in eastern Europe (Nagavciuc et al., 2022) nor in eastern Mediterranean (Touchan et al., 2005). Similarly, winter records before 1850 also display discrepancies. These discrepancies may be attributed to a higher uncertainty in the ice core winter accumulation record, as well as a reduction in the number of observations included in the EKF400 v.2 dataset. Similarly, winter records also exhibit discrepancies before 1850, possibly attributed to a decrease in the number of observations included in the EKF400 v.2 dataset. The WP accumulation record effectively captures the decadal and long-term variability for a larger region during both summer and winter seasons.

The most recent little ice age (LIA) glaciation maximum was observed in the Caucasus in the middle of the 19th century and is well reflected in the dendrochronological data from this region. The general glacier retreat in the North Caucasus began in

the late 1840s, which was interrupted 4-5 times in 1860-1880, as well as in the 1910s, 1920s, and 1970-80s (Solomina et al., 2016). According to our results, the period of glacier retreat was preceded by a long period of negative precipitation trend in summer, which was accompanied by an increase in summer temperature, reconstructed from the tree ring widths (Dolgova, 2016). The period of short readvances at the end of 19<sup>th</sup> century coincide with several intervals of increased summer and winter accumulation. The recent increase in both summer and winter accumulation, despite the continued rise in air temperature, caused the positioning of glacier fronts and their minor advance in the 1970s and 1980s (Solomina et al., 2016).

580 The primary driver of climatic changes during the late eighteenth and nineteenth centuries was the natural quasi-decadal variability of the thermohaline ocean circulation. The summer temperatures reconstructed for the Caucasus using tree rings are strongly correlated with AMO oscillations (Dolgova, 2016). Our findings support the role of North Atlantic oscillations not

585 only in temperature, but also in precipitation levels in this region especially during winter seasons.



590 **Figure 1011.** Spatial correlation of the 10-year moving average of summer (a) (April-September) and winter (b) (October-March) WP accumulation with EK400v2 paleo reanalysis ensemble mean precipitation data over 1900-2003 period, time series of the precipitation data averaged over the area indicated by the black box for summer (c) and winter (d) shown together with WP accumulation data. Location of the Elbrus Mt. is shown by black triangle.

#### 4 Conclusions

The ice core retrieved from Mt. Elbrus was dated back to 1750 CE using annual layer counting, with a focus on high-resolution oscillations of ammonium and succinic acid. A new depth-age scale was established by reference horizons associated with

known volcanic eruptions. The high accumulation rate and sampling resolution of the ice core facilitated the identification of both annual and seasonal layers. The reconstructed accumulation with the process of layers thinning and the upstream effect accounted for allowed for a meaningful comparison with meteorological data.

To distinguish between winter and summer seasons in meteorological data, a method based on Convective Available Potential Energy was ~~proposed~~applied, resulting in a significant improvement in detecting annual and seasonal precipitation. Comparisons of the distribution functions and spectral density of annual precipitation between meteorological stations in the Caucasus and ice-core records provided clear agreement, validating the reconstructed data as representative of precipitation. The comparison between the obtained ice core accumulation record from WP and gridded precipitation datasets, as well as paleo reanalysis data, revealed that the accumulation record represents the precipitation regime on decadal and long-term timescales in a larger region encompassing the Northern Caucasus, Black Sea, and South-Eastern Europe.

~~The analysis showed that precipitation variability in the WP was primarily influenced by the summer season, with a significant decrease in the contribution of summer precipitation over a span of 260 years, corresponding to the transition from the Little Ice Age to modern climatic conditions.~~

Variations in precipitation were identified with periodicities of 20 and 40 years, corresponding to the typical quasi-decadal variability associated with North Atlantic thermohaline circulation oscillations. Statistically significant relationship was observed between ice core accumulation pattern and fluctuations in the NAO. Similar variations were found in dendrochronological data on air temperature anomalies in the Caucasus, which aligned with the AMO index. This study supports the hypothesis that the quasi-decadal variations in temperature-~~moisture~~humidity-regime in the Caucasus have an oceanic nature.

## References

Ahrens, J., Geveci, B. and Law, C.: ParaView: An End-User Tool for Large-Data Visualization, in Visualization Handbook, pp. 717–731, Elsevier., 2005.

Aleshina, M. A., Toropov, P. A. and Semenov, V. A.: Temperature and Humidity Regime Changes on the Black Sea Coast in 1982-2014, Russ. Meteorol. Hydrol., 43(4), 235–244, doi:10.3103/S1068373918040040, 2018.

Barber, K., Zolitschka, B., Tarasov, P. and Lotter, A. F.: Atlantic to Urals – the Holocene climatic record of Mid-Latitude Europe, in Past Climate Variability through Europe and Africa, pp. 417–442, Springer Netherlands, Dordrecht., 2004.

Bardin, M. Y., Platova, T. V. and Samokhina, O. F.: VARIABILITY OF ANTI-CYCLONIC ACTIVITY IN THE NORTHERN EXTRATROPICS, Fundam. Appl. Climatol., 3, 32–58, doi:10.21513/0207-2564-2019-3-32-58, 2019.

Bintanja, R.: The contribution of snowdrift sublimation to the surface mass balance of Antarctica, Ann. Glaciol., 27, 251–259, doi:10.3189/1998AoS27-1-251-259, 1998.

- Bohleber, P., Wagenbach, D., Schöner, W. and Böhm, R.: To what extent do water isotope records from low accumulation  
625 [Alpine ice cores reproduce instrumental temperature series?](#), *Tellus, Ser. B Chem. Phys. Meteorol.*, 65(1), 1–17,  
[doi:10.3402/tellusb.v65i0.20148, 2013.](#)
- Borisova, O.: Environmental and climatic conditions of human occupation in the central East European Plain during the Middle  
Holocene: Reconstruction from palaeofloristic data, *Quat. Int.*, 516, 42–57, doi:10.1016/j.quaint.2018.05.025, 2019.
- Bunde, A., Büntgen, U., Ludescher, J., Luterbacher, J. and von Storch, H.: Is there memory in precipitation?, *Nat. Clim.*  
630 *Chang.*, 3(3), 174–175, doi:10.1038/nclimate1830, 2013.
- Büntgen, U., Urban, O., Krusic, P. J., Rybníček, M., Kolář, T., Kyncl, T., Ač, A., Koňasová, E., Čáslavský, J., Esper, J.,  
Wagner, S., Saurer, M., Tegel, W., Dobrovolný, P., Cherubini, P., Reinig, F. and Trnka, M.: Recent European drought extremes  
beyond Common Era background variability, *Nat. Geosci.*, 14(4), 190–196, doi:10.1038/s41561-021-00698-0, 2021.
- Chen, J., Del Genio, A. D., Carlson, B. E. and Bosilovich, M. G.: The spatiotemporal structure of twentieth-century climate  
635 variations in observations and reanalyses. Part I: Long-term trend, *J. Clim.*, 21(11), 2611–2633, doi:10.1175/2007JCLI2011.1,  
2008.
- Chernokulsky, A., Kozlov, F., Zolina, O., Bulygina, O., Mokhov, I. I. and Semenov, V. A.: Observed changes in convective  
and stratiform precipitation in Northern Eurasia over the last five decades, *Environ. Res. Lett.*, 14(4), 045001,  
doi:10.1088/1748-9326/aafb82, 2019.
- 640 Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker,  
T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. B. and P. M. M. (eds. )]: *Climate  
Change 2013 – The Physical Science Basis*, edited by Intergovernmental Panel on Climate Change, Cambridge University  
Press, Cambridge, United Kingdom and New York, NY, USA,, 2014.
- Cook, E. R., Seager, R., Kushnir, Y., Briffa, K. R., Büntgen, U., Frank, D., Krusic, P. J., Tegel, W., van der Schrier, G.,  
645 Andreu-Hayles, L., Baillie, M., Baittinger, C., Bleicher, N., Bonde, N., Brown, D., Carrer, M., Cooper, R., Cufar, K., Dittmar,  
C., Esper, J., Griggs, C., Gunnarson, B., Günther, B., Gutierrez, E., Haneca, K., Helama, S., Herzig, F., Heussner, K.-U.,  
Hofmann, J., Janda, P., Kontic, R., Köse, N., Kyncl, T., Levanić, T., Linderholm, H., Manning, S., Melvin, T. M., Miles, D.,  
Neuwirth, B., Nicolussi, K., Nola, P., Panayotov, M., Popa, I., Rothe, A., Seftigen, K., Seim, A., Svarva, H., Svoboda, M.,  
Thun, T., Timonen, M., Touchan, R., Trotsiuk, V., Trouet, V., Walder, F., Wazny, T., Wilson, R. and Zang, C.: Old World  
650 megadroughts and pluvials during the Common Era, *Sci. Adv.*, 1(10), e1500561–e1500561, doi:10.1126/sciadv.1500561,  
2015.
- Cook, E. R., Solomina, O., Matkovsky, V., Cook, B. I., Agafonov, L., Berdnikova, A., Dolgova, E., Karpukhin, A., Knysh,  
N., Kulakova, M., Kuznetsova, V., Kyncl, T., Kyncl, J., Maximova, O., Panyushkina, I., Seim, A., Tishin, D., Wazny, T. and  
Yermokhin, M.: The European Russia Drought Atlas (1400–2016 CE), *Clim. Dyn.*, 54(3–4), 2317–2335, doi:10.1007/s00382-  
655 019-05115-2, 2020.

- Dahl-Jensen, D., Johnsen, S. J., Hammer, C. U., Clausen, H. B. and Jouzel, J.: Past Accumulation rates derived from observed annual layers in the GRIP ice core from Summit, Central Greenland, in *Ice in the Climate System*, pp. 517–532, Springer Berlin Heidelberg, Berlin, Heidelberg., 1993.
- Dansgaard, W. and Johnsen, S. J.: A Flow Model and a Time Scale for the Ice Core from Camp Century, Greenland, *J. Glaciol.*, 660 8(53), 215–223, doi:10.3189/S0022143000031208, 1969.
- Deser, C., Hurrell, J. W. and Phillips, A. S.: The role of the North Atlantic Oscillation in European climate projections, *Clim. Dyn.*, 49(9–10), 3141–3157, doi:10.1007/s00382-016-3502-z, 2017.
- Dolgova, E.: June-September temperature reconstruction in the Northern Caucasus based on blue intensity data, *Dendrochronologia*, 39, 17–23, doi:10.1016/j.dendro.2016.03.002, 2016.
- 665 Gagliardini, O. and Meyssonier, J.: Flow simulation of a firn-covered cold glacier, *Ann. Glaciol.*, 24, 242–248, doi:10.3189/S0260305500012246, 1997.
- Fuentes-Franco, R., Docquier, D., Koenigk, T., Zimmermann, K. and Giorgi, F.: Winter heavy precipitation events over Northern Europe modulated by a weaker NAO variability by the end of the 21st century, *npj Clim. Atmos. Sci.*, 6(1), 72, doi:10.1038/s41612-023-00396-1, 2023.
- 670 Gagliardini, O., Zwinger, T., Gillet-Chaulet, F., Durand, G., Favier, L., de Fleurian, B., Greve, R., Malinen, M., Martín, C., Råback, P., Ruokolainen, J., Sacchettini, M., Schäfer, M., Seddik, H. and Thies, J.: Capabilities and performance of Elmer/Ice, a new-generation ice sheet model, *Geosci. Model Dev.*, 6(4), 1299–1318, doi:10.5194/gmd-6-1299-2013, 2013.
- Goodwin, B. P., Mosley-Thompson, E., Wilson, A. B., Porter, S. E. and Roxana Sierra-Hernandez, M.: Accumulation variability in the Antarctic Peninsula: The role of large-scale atmospheric oscillations and their interactions\*, *J. Clim.*, 29(7), 675 2579–2596, doi:10.1175/JCLI-D-15-0354.1, 2016.
- Henderson, K., Laube, A., Gäggeler, H. W., Olivier, S., Papina, T. and Schwikowski, M.: Temporal variations of accumulation and temperature during the past two centuries from Belukha ice core, Siberian Altai, *J. Geophys. Res.*, 111(D3), D03104, doi:10.1029/2005JD005819, 2006.
- Herren, P.-A., Eichler, A., Machguth, H., Papina, T., Tobler, L., Zapf, A. and Schwikowski, M.: The onset of Neoglaciation 680 6000 years ago in western Mongolia revealed by an ice core from the Tsambagarav mountain range, *Quat. Sci. Rev.*, 69, 59–68, doi:10.1016/j.quascirev.2013.02.025, 2013.
- Hörhold, M. W., Kipfstuhl, S., Wilhelms, F., Freitag, J. and Frenzel, A.: The densification of layered polar firn, *J. Geophys. Res. Earth Surf.*, 116(F1), n/a-n/a, doi:10.1029/2009JF001630, 2011.
- Hurrell, J. W. and Deser, C.: North Atlantic climate variability: The role of the North Atlantic Oscillation, *J. Mar. Syst.*, 78(1), 685 28–41, doi:10.1016/j.jmarsys.2008.11.026, 2009.
- Ionita, M.: The Impact of the East Atlantic/Western Russia Pattern on the Hydroclimatology of Europe from Mid-Winter to Late Spring, *Climate*, 2(4), 296–309, doi:10.3390/cli2040296, 2014.

- Kozachek, A., Mikhailenko, V., Masson-Delmotte, V., Ekaykin, A., Ginot, P., Kutuzov, S., Legrand, M., Lipenkov, V. and Preunkert, S.: Large-scale drivers of Caucasus climate variability in meteorological records and Mt El'brus ice cores, *Clim. Past*, 13(5), 473–489, doi:10.5194/cp-13-473-2017, 2017.
- 690 Kutuzov, S., Legrand, M., Preunkert, S., Ginot, P. and Mikhailenko, V.: The Elbrus (Caucasus, Russia) ice core record – Part 2: history of desert dust deposition, , 14133–14148, 2019.
- Lavrentiev, I. I., Mikhailenko, V. N. and Kutuzov, S. S.: Ice thickness and subglacial relief of the Western ice plateau of Elbrus., *Ice Snow*, 2, 2–18, 2010.
- 695 Lavrentiev, I. I., Kutuzov, S. S., Mikhailenko, V. N., Sudakova, M. S. and Kozachek, A. V.: Spatial and Temporal Variations of Snow Accumulation on the Western Elbrus Plateau, the Central Caucasus, *Water Resour.*, 49(S1), S1–S11, doi:10.1134/S0097807822070090, 2022.
- Licciulli, C., Bohleber, P., Lier, J., Gagliardini, O., Hoelzle, M. and Eisen, O.: A full Stokes ice-flow model to assist the interpretation of millennial-scale ice cores at the high-Alpine drilling site Colle Gnifetti, Swiss/Italian Alps, *J. Glaciol.*, 700 66(255), 35–48, doi:10.1017/jog.2019.82, 2019.
- Ligtenberg, S. R. M., Helsen, M. M. and van den Broeke, M. R.: An improved semi-empirical model for the densification of Antarctic firn, *Cryosph.*, 5(4), 809–819, doi:10.5194/tc-5-809-2011, 2011.
- Lim, S., Faïn, X., Ginot, P., Mikhailenko, V., Kutuzov, S., Paris, J. D., Kozachek, A. and Laj, P.: Black carbon variability since preindustrial times in the eastern part of Europe reconstructed from Mt. Elbrus, Caucasus, ice cores, *Atmos. Chem. Phys.*, 705 17(5), 3489–3505, doi:10.5194/acp-17-3489-2017, 2017.
- Loader, N. J., Young, G. H. F., McCarroll, D., Davies, D., Miles, D. and Bronk Ramsey, C.: Summer precipitation for the England and Wales region, 1201–2000 CE, from stable oxygen isotopes in oak tree rings, *J. Quat. Sci.*, 35(6), 731–736, doi:10.1002/jqs.3226, 2020.
- López-Moreno, J. I., Vicente-Serrano, S. M., Morán-Tejeda, E., Lorenzo-Lacruz, J., Kenawy, A. and Beniston, M.: Effects of 710 the North Atlantic Oscillation (NAO) on combined temperature and precipitation winter modes in the Mediterranean mountains: Observed relationships and projections for the 21st century, *Glob. Planet. Change*, 77(1–2), 62–76, doi:10.1016/j.gloplacha.2011.03.003, 2011.
- Maeno, N. and Ebinuma, T.: Pressure sintering of ice and its implication to the densification of snow at polar glaciers and ice sheets, *J. Phys. Chem.*, 87(21), 4103–4110, doi:10.1021/j100244a023, 1983.
- 715 Mariani, I., Eichler, A., Jenk, T. M., Brönnimann, S., Auchmann, R., Leuenberger, M. C. and Schwikowski, M.: Temperature and precipitation signal in two Alpine ice cores over the period 1961–2001, *Clim. Past*, 10(3), 1093–1108, doi:10.5194/cp-10-1093-2014, 2014.
- Markowski, P. and Richardson, Y.: *Mesoscale Meteorology in Midlatitudes*, Wiley., 2010.
- Martin-Benito, D., Ummenhofer, C. C., Köse, N., Güner, H. T. and Pederson, N.: Tree-ring reconstructed May–June 720 precipitation in the Caucasus since 1752 CE, *Clim. Dyn.*, 47(9–10), 3011–3027, doi:10.1007/s00382-016-3010-1, 2016.

- Meredith, E. P., Semenov, V. A., Maraun, D., Park, W. and Chernokulsky, A. V.: Crucial role of Black Sea warming in amplifying the 2012 Krymsk precipitation extreme, *Nat. Geosci.*, 8(8), 615–619, doi:10.1038/ngeo2483, 2015.
- Mikhaleiko, V., Sokratov, S., Kutuzov, S., Ginot, P., Legrand, M., Preunkert, S., Lavrentiev, I., Kozachek, A., Ekaykin, A., Fain, X., Lim, S., Schotterer, U., Lipenkov, V. and Toropov, P.: Investigation of a deep ice core from the Elbrus western plateau, the Caucasus, Russia, *Cryosphere*, 9(6), 2253–2270, doi:10.5194/tc-9-2253-2015, 2015.
- Mikhaleiko, V. N., Ed.: *Elbrus Glaciers and Climate*, Nestor-Historia Publications, Moscow-St. Petersburg, Russia., 2020.
- Mikhaleiko, V. N., Kutuzov, S. S., Lavrentiev, I. I., Toropov, P. A., Vladimirova, D. O., Abramov, A. A. and Matskovsky, V. V.: Glacioclimatological investigations of the Institute of Geography, RAS, in the crater of Eastern Summit of Mt. Elbrus in 2020, *Ice Snow*, 61(1), 149–160, doi:10.31857/S2076673421010078, 2021.
- 725 Nagavciuc, V., Ionita, M., Kern, Z., McCarroll, D. and Popa, I.: A ~700 years perspective on the 21st century drying in the eastern part of Europe based on  $\delta^{18}\text{O}$  in tree ring cellulose, *Commun. Earth Environ.*, 3(1), 277, doi:10.1038/s43247-022-00605-4, 2022.
- Palm, S. P., Kayetha, V., Yang, Y. and Pauly, R.: Blowing snow sublimation and transport over Antarctica from 11 years of CALIPSO observations, *Cryosph.*, 11(6), 2555–2569, doi:10.5194/tc-11-2555-2017, 2017.
- 735 Pauling, A., Luterbacher, J., Casty, C. and Wanner, H.: Five hundred years of gridded high-resolution precipitation reconstructions over Europe and the connection to large-scale circulation, *Clim. Dyn.*, 26(4), 387–405, doi:10.1007/s00382-005-0090-8, 2006.
- Pohjola, V. A., Martma, T. A., Meijer, H. A. J., Moore, J. C., Isaksson, E., Vaikmäe, R. and van de Wal, R. S. W.: Reconstruction of three centuries of annual accumulation rates based on the record of stable isotopes of water from Lomonosovfonna, Svalbard, *Ann. Glaciol.*, 35, 57–62, doi:10.3189/172756402781816753, 2002.
- Pomeroy, J. W. and Gray, D. M.: *Snowcover accumulation, relocation and management*, Saskatoon, Saskatchewan : National Hydrology Research Institute., 1995.
- Preunkert, S. and Legrand, M.: Towards a quasi-complete reconstruction of past atmospheric aerosol load and composition (organic and inorganic) over Europe since 1920 inferred from Alpine ice cores, *Clim. Past*, 9(4), 1403–1416, doi:10.5194/cp-9-1403-2013, 2013.
- 745 Preunkert, S., Legrand, M., Kutuzov, S., Ginot, P., Mikhaleiko, V. and Friedrich, R.: The Elbrus (Caucasus, Russia) ice core record-Part 1: Reconstruction of past anthropogenic sulfur emissions in south-eastern Europe, *Atmos. Chem. Phys.*, 19(22), 14119–14132, doi:10.5194/acp-19-14119-2019, 2019.
- Rasmussen, R., Baker, B., Kochendorfer, J., Meyers, T., Landolt, S., Fischer, A. P., Black, J., Thériault, J. M., Kucera, P., Gochis, D., Smith, C., Nitu, R., Hall, M., Ikeda, K. and Gutmann, E.: How well are we measuring snow: The NOAA/FAA/NCAR winter precipitation test bed, *Bull. Am. Meteorol. Soc.*, 93(6), 811–829, doi:10.1175/BAMS-D-11-00052.1, 2012.
- 750 Simkin, T. and Siebert, L.: *Volcanoes of the World*, Geoscience Press, Inc., Tucson, Arizona., 1994.

- Solomina, O., Davi, N., D'Arrigo, R. and Jacoby, G.: Tree-ring reconstruction of Crimean drought and lake chronology correction, *Geophys. Res. Lett.*, 32(19), n/a-n/a, doi:10.1029/2005GL023335, 2005.
- 755 Solomina, O., Bushueva, I., Dolgova, E., Jomelli, V., Alexandrin, M., Mikhailenko, V. and Matskovsky, V.: Glacier variations in the Northern Caucasus compared to climatic reconstructions over the past millennium, *Glob. Planet. Change*, 140, 28–58, doi:10.1016/j.gloplacha.2016.02.008, 2016.
- Solomina, O. N., Bushueva, I. S., Dolgova, E. A., Zolotokrylin, A. N., Kuznetsova, V. V., Kuznetsova, T. O., Kukhta, A. E.,  
760 Lazukova, L. I., Lomakin, N. A., Matskovsky, V. V., Matveev, S. M., Mikhailov, A. Y., Mikhailenko, V. N., Pozhidaeva, L. S., Rummyantsev, D. E., Sakulina, G. A., Semenov, V. A., Khasanov, B. F., Cherenkova, E. A. and Chernokulsky, A. V.: Droughts of the East European Plain according to hydrometeorological and tree-ring data, Nestor-Historia Publications, Moscow-St. Petersburg, Russia., 2017.
- Sun, Q., Miao, C., Duan, Q., Ashouri, H., Sorooshian, S. and Hsu, K.: A Review of Global Precipitation Data Sets: Data  
765 Sources, Estimation, and Intercomparisons, *Rev. Geophys.*, 56(1), 79–107, doi:10.1002/2017RG000574, 2018.
- Tashilova, A., Ashabokov, B., Kesheva, L. and Teunova, N.: Analysis of Climate Change in the Caucasus Region: End of the 20th–Beginning of the 21st Century, *Climate*, 7(1), 11, doi:10.3390/cli7010011, 2019.
- Tielidze, L. G., Nosenko, G. A., Khromova, T. E. and Paul, F.: Strong acceleration of glacier area loss in the Greater Caucasus between 2000 and 2020, *Cryosph.*, 16(2), 489–504, doi:10.5194/tc-16-489-2022, 2022.
- 770 Touchan, R., Akkemik, Ü., Hughes, M. K. and Erkan, N.: May-June precipitation reconstruction of southwestern Anatolia, Turkey during the last 900 years from tree rings, *Quat. Res.*, 68(2), 196–202, doi:10.1016/j.yqres.2007.07.001, 2007.
- Touchan, R., Xoplaki, E., Funkhouser, G., Luterbacher, J., Hughes, M. K., Erkan, N., Akkemik, Ü. and Stephan, J.: Reconstructions of spring/summer precipitation for the Eastern Mediterranean from tree-ring widths and its connection to large-scale atmospheric circulation, *Clim. Dyn.*, 25(1), 75–98, doi:10.1007/s00382-005-0016-5, 2005.
- 775 Valler, V., Franke, J., Brugnara, Y. and Brönnimann, S.: An updated global atmospheric paleo-reanalysis covering the last 400 years, *Geosci. Data J.*, 9(1), 89–107, doi:10.1002/gdj3.121, 2022.
- Verhaegen, Y., Huybrechts, P., Rybak, O. and Popovnin, V. V.: Modelling the evolution of Djankuat Glacier, North Caucasus, from 1752 until 2100 CE, *Cryosph.*, 14(11), 4039–4061, doi:10.5194/tc-14-4039-2020, 2020.
- Vicente-Serrano, S. M. and López-Moreno, J. I.: Nonstationary influence of the North Atlantic Oscillation on European  
780 precipitation, *J. Geophys. Res.*, 113(D20), D20120, doi:10.1029/2008JD010382, 2008.
- Winski, D., Osterberg, E., Ferris, D., Kreutz, K., Wake, C., Campbell, S., Hawley, R., Roy, S., Birkel, S., Introne, D. and Handley, M.: Industrial-age doubling of snow accumulation in the Alaska Range linked to tropical ocean warming, *Sci. Rep.*, 7(1), 17869, doi:10.1038/s41598-017-18022-5, 2017.
- Winstrup, M., Vallenga, P., Kjær, H. A., Fudge, T. J., Lee, J. E., Riis, M. H., Edwards, R., Bertler, N. A. N., Blunier, T.,  
785 Brook, E. J., Buizert, C., Ciobanu, G., Conway, H., Dahl-Jensen, D., Ellis, A., Emanuelsson, B. D., Hindmarsh, R. C. A., Keller, E. D., Kurbatov, A. V., Mayewski, P. A., Neff, P. D., Pyne, R. L., Simonsen, M. F., Svensson, A., Tuohy, A.,



Waddington, E. D. and Wheatley, S.: A 2700-year annual timescale and accumulation history for an ice core from Roosevelt Island, West Antarctica, *Clim. Past*, 15(2), 751–779, doi:10.5194/cp-15-751-2019, 2019.

790 Yao, T., Duan, K., Xu, B., Wang, N., Guo, X. and Yang, X.: Precipitation record since AD 1600 from ice cores on the central Tibetan Plateau, *Clim. Past*, 4(3), 175–180, doi:10.5194/cp-4-175-2008, 2008.

Zhang, R., Shang, H., Yu, S., He, Q., Yuan, Y., Bolatov, K. and Mambetov, B. T.: Tree-ring-based precipitation reconstruction in southern Kazakhstan, reveals drought variability since A.D. 1770, *Int. J. Climatol.*, 37(2), 741–750, doi:10.1002/joc.4736, 2017.

795 Zhang, W., Hou, S., Wu, S.-Y., Pang, H., Sneed, S. B., Korotkikh, E. V., Mayewski, P. A., Jenk, T. M. and Schwikowski, M.: A quantitative method of resolving annual precipitation for the past millennia from Tibetan ice cores, *Cryosph.*, 16(5), 1997–2008, doi:10.5194/tc-16-1997-2022, 2022.

Zwinger, T., Greve, R., Gagliardini, O., Shiraiwa, T. and Lyly, M.: A full Stokes-flow thermo-mechanical model for firn and ice applied to the Gorshkov crater glacier, Kamchatka, *Ann. Glaciol.*, 45, 29–37, doi:10.3189/172756407782282543, 2007.

#### 800 **Data availability:**

Accumulation reconstruction ~~data are available from the authors upon request~~ presented in the supplementary (Table S12) and will be accessible on PANGAEA. Reanalysis ERA5 data are available at <https://cds.climate.copernicus.eu/> (last access 16 May 2023), GPCC data are available at <https://psl.noaa.gov/data/gridded/data.gpcc.html> (last access: 16 May 2023). Ensemble Kalman Fitting Paleo-Reanalysis Version 2.0 (EKF400\_v2.0) is available at [https://www.wdc-climate.de/ui/entry?acronym=EKF400\\_v2.0](https://www.wdc-climate.de/ui/entry?acronym=EKF400_v2.0). (last access 16 May 2023).

805

#### **Author contributions:**

The paper was written by VM and SK with contributions from PT, ML, GC, and SS. The ice-core chemistry records were produced by SP and ML; the water isotope measurements were made by AK and VL. The layer-counted timescale was developed by ML, SP, SK, MV and AK. Volcanic identification and synchronization to time scale were performed by ML and 810 SP. Seasonal snow accumulation distribution was estimated by IL. GC calculated backward ice flow trajectories. SS constructed the thinning function. Interpretation of the accumulation rate history was performed by SK, VM and PT. All the authors read and discussed the manuscript and contributed to improving the final paper.

#### **Competing interests:**

The authors declare that they have no conflict of interest.

815 **Acknowledgements:**

This work was carried out within the framework of the Russian Science Foundation project 17-17-01270. The study was completed in the laboratory created within Megagrant project (agreement no. 075-15-2021-599, 08.06.2021) with the support of the FMGE-2019-0004 project. We thank all the people who provided support at all stages of the work, participated in the field work, carried out the sampling and analytical processing of the ice core. We are grateful to V. Matskovsky for his help  
820 in interpreting dendrochronological data for the Caucasus.