



# Drivers of late Holocene ice core chemistry in Dronning Maud Land: The context for the ISOL-ICE project

V. Holly L. Winton<sup>1,2</sup>, Robert Mulvaney<sup>1</sup>, Joel Savarino<sup>3</sup>, Kyle R. Clem<sup>4</sup>, Markus M. Frey<sup>1</sup>

<sup>1</sup>British Antarctica Survey, Cambridge, CB3 0ET, United Kingdom

5 <sup>2</sup>Antarctic Research Centre, Victoria University of Wellington, Wellington 6102, New Zealand

<sup>3</sup>University of Grenoble Alpes, CNRS, IRD, Grenoble INP, INRAE, IGE, F-38000 Grenoble, France

<sup>4</sup>School of Geography, Environment and Earth Sciences, Victoria University of Wellington, Wellington 6102, New Zealand

Correspondence to: V. Holly L. Winton ([holly.winton@vuw.ac.nz](mailto:holly.winton@vuw.ac.nz)) and Markus M. Frey ([maey@bas.ac.uk](mailto:maey@bas.ac.uk))

**Abstract.** Quantifying the natural variability of the stratospheric ozone layer and understanding the underlying factors that control natural total column ozone (TCO) variability are required to put modern observations into historical context and evaluate the effectiveness of climate and TCO protection policies. Within the framework of the Isotopic Constraints on Past Ozone Layer in Polar Ice (ISOL-ICE) project, we present initial ice core results from the new ISOL-ICE ice core covering the last millennium from the high-elevation Dronning Maud Land (DML) located under the Antarctic spring stratospheric TCO minimum, and discuss the implications for interpreting the stable isotopic composition of nitrogen in ice core nitrate ( $\delta^{15}\text{N}(\text{NO}_3^-)$ ) as a surface ultra-violet radiation (UV) and TCO proxy. To interpret the ice core  $\delta^{15}\text{N}(\text{NO}_3^-)$  record, an understanding of past snow accumulation changes, as well as aerosol source regions and present-day drivers of their variability are required. We therefore report here the ice core age-depth model, the snow accumulation and ice chemistry records, and correlation analysis of these records with climate variables over the observational era (1979-2016). The ISOL-ICE ice core covers the last 1349 years from 668 to 2017 C.E.  $\pm 3$  years extending previous ice core records from the region by two decades and shows excellent reproducibility with those records. The extended ISOL-ICE record of last two decades showed a continuation of the methanesulphonate (MSA) increase from  $\sim 1800$  to present while there were less frequent large deposition events of sea salts relative to the last millennium. The correlation analysis, combined with the finding that sea salts do not carry a sea ice signature to the site, highlight that sea salt and MSA aerosol concentrations are primarily related to atmospheric transport over the extended two-decade period and not to changes in sea ice source strength. Correlation of the snow accumulation record with climate variables over the observational era showed that precipitation at ISOL-ICE is predominately derived from the South Atlantic with onshore winds delivering marine air masses to the site. The snow accumulation rate was stable over the last millennium with no notable trends over last two decades relative to the last millennium. Interannual variability in the accumulation record, ranging between 2 and 20  $\text{cm a}^{-1}$  (w.e.), would influence the ice core  $\delta^{15}\text{N}(\text{NO}_3^-)$  record. The mean snow accumulation rate of  $6.5 \pm 2.4 \text{ cm a}^{-1}$  (w.e.) falls within the range suitable for reconstructing surface mass balance from ice core  $\delta^{15}\text{N}(\text{NO}_3^-)$  highlighting that the ISOL-ICE ice core  $\delta^{15}\text{N}(\text{NO}_3^-)$  can be used to reconstruct either the snow accumulation rate or surface UV if the ice core  $\delta^{15}\text{N}(\text{NO}_3^-)$  is corrected for the snow accumulation influence.



## 1 Introduction

The stratospheric ozone O<sub>3</sub> layer shields all land-based life forms from harmful ultraviolet (UV-B) radiation (wavelength 290-315 nm). Satellites and ground-based instruments measure total column ozone (TCO) to which the largest contribution is the stratospheric O<sub>3</sub> layer (e.g. Bodeker et al., 2021). Yet, no direct observations of TCO exists prior to 1926 (Brönnimann et al., 2003). Assessing the natural variability of TCO before the instrumental era is critical to put modern observations into historical context but this has proven difficult owing to the lack of a well-constrained proxy with the exception of a UV biomarker in lake sediments (Lomax et al., 2008), and a chemical signature of UV-B absorbing compounds in *Pinus* pollen grains (Jardine et al., 2020). To address the lack of suitable UV radiation proxies, the Isotopic Constraints on Past Ozone Layer in Polar Ice (ISOL-ICE) project drilled a new 120 m ice core from the East Antarctic Plateau in 2017 to apply a new ice core proxy based on the fractionation observed in stable nitrogen ( $\delta^{15}\text{N}$ ) and oxygen ( $\Delta^{17}\text{O}$ ) isotopes of nitrate ( $\text{NO}_3^-$ ) and to develop the first millennial-timescale reconstruction of past surface UV-radiation (TCO) and atmospheric oxidation processes. Interpretation of the nitrate stable isotope record ( $\delta^{15}\text{N}$  and  $\Delta^{17}\text{O}$ ) in terms of past UV (TCO) and atmospheric oxidation processes requires a thorough understanding of the local air-transfer function and its driving parameters. The stable isotope of nitrogen  $^{15}\text{N}$  in  $\text{NO}_3^-$  ( $\delta^{15}\text{N}(\text{NO}_3^-)$ ) in surface snow is highly enriched compared to atmospheric  $\text{NO}_3^-$  (e.g., Erbland et al., 2013; Cao et al., 2022; Winton et al., 2020) due to  $\text{NO}_3^-$  loss and redistribution from snow, which is driven by UV-photolysis (Frey et al., 2009; Berhanu et al., 2014; Shi et al., 2019).  $\text{NO}_3^-$  isotopic fractionation is strongest at sites with very low snow accumulation rates (2-3 cm a<sup>-1</sup> (w.e.), such as Dome A and Dome C on the East Antarctic Plateau, due to enhanced  $\text{NO}_3^-$  post-depositional recycling which erases the source signature of  $\delta^{15}\text{N}(\text{NO}_3^-)$  due to longer exposure of surface snow layers to incoming UV radiation before burial (Shi et al., 2022a; Frey et al., 2009). Increases of snow accumulation rate to 6 cm a<sup>-1</sup> (w.e.) and above is sufficient to preserve the seasonal cycle of  $\delta^{15}\text{N}(\text{NO}_3^-)$  in the snowpack (Winton et al., 2020) and enrichments in  $\delta^{15}\text{N}(\text{NO}_3^-)$  by the effects of snow accumulation superimpose those due to stratospheric ozone depletion (Cao et al., 2022; Shi et al., 2022b). The inverse relationship between the snow accumulation rate and  $\delta^{15}\text{N}(\text{NO}_3^-)$  is well constrained across Antarctica so that for ice core sites within the transition zones between the dome summits and the coast with a snow accumulation rate between 4 and 20 cm a<sup>-1</sup> (w.e.), ice core  $\delta^{15}\text{N}(\text{NO}_3^-)$  has been proposed as proxy for surface mass balance (Akers et al., 2022b).

As a first part of the ISOL-ICE project, an air-snow transfer study was carried out at the ice core drill site (Winton et al., 2020). At the ISOL-ICE ice core site located on the high East Antarctic Plateau at Kohnen Station (75°00'06"S, 0°04'04"E, 2892 m amsl) in Dronning Maud Land (DML), the sensitivity of  $\delta^{15}\text{N}(\text{NO}_3^-)$  to changes in snow accumulation (and seasonality), TCO and snow e-folding depth (light attenuation) was assessed using atmospheric, surface snow and snow pit observations of  $\delta^{15}\text{N}(\text{NO}_3^-)$  supported by air-snow transfer modelling (Winton et al., 2020). The dominant factors controlling the archived  $\delta^{15}\text{N}(\text{NO}_3^-)$  signature at the ISOL-ICE site are the e-folding depth and snow accumulation rate, followed by TCO. Taking past changes in snow accumulation into account, it may be possible to reconstruct past UV and TCO on longer timescales from the  $\delta^{15}\text{N}(\text{NO}_3^-)$  signal in the ISOL-ICE ice cores, provided other factors such as the e-folding depth have remained the same within



65 the target resolution time. Thus, in order to interpret the ice core  $\delta^{15}\text{N}(\text{NO}_3^-)$  record, the accumulation rate needs to be independently derived to account for post-depositional isotope fractionation not related to surface-UV changes.

Furthermore, constraining the aerosol source regions for the ISOL-ICE ice core has implications for interpreting the ice core record of  $\text{NO}_3^-$  and its stable isotopes. The net primary sources of  $\text{NO}_3^-$  to the Antarctic atmosphere and snowpack are methyl  
70 nitrate ( $\text{CH}_3\text{NO}_3$ ) from open ocean and sea ice, and stratospheric  $\text{NO}_3^-$  (Burger et al., 2022; Burger et al., 2023; Mulvaney and Wolff, 1993; Savarino et al., 2007; Jacobi et al., 2000; Jones et al., 1999; Beyersdorf et al., 2010). Ice core proxies of open ocean and sea ice aerosol include biogenic sulfur species and sea salts (Frey et al., 2020; Abram et al., 2013; Thomas et al., 2019; Thomas et al., 2023). At some ice core sites, atmospheric transport strength, rather than aerosol source, is the dominant  
75 factor that determines the chemical signal preserved in ice cores. For example, methanesulfonic acid (MSA) is derived from the oxidation of dimethylsulfide (DMS,  $(\text{CH}_3)_2\text{S}$ ) from marine phytoplankton emissions to the atmosphere, and reflects atmospheric transport rather than sea ice conditions in coastal Weddell Sea and Dronning Maud Land ice cores (e.g. Fundel et al., 2006; Abram et al., 2007). Thus, evaluating the ISOL-ICE ice chemistry derived from oceanic sources is necessary to understand the relationships between ice core  $\text{NO}_3^-$  and its sources.

Within the framework of the ISOL-ICE project, we present the full ice core record of annual snow accumulation and ice core  
80 chemistry, and the age-depth model. The ISOL-ICE ice core extends the European Program for Ice Coring in Antarctica (EPICA) Dronning Maud Land (EDML) record by two decades towards the present. We discuss the aerosol source regions and links to climate variability to provide context for potential  $\text{NO}_3^-$  sources. These initial ice core results, together with the ISOL-ICE air-snow transfer study (Winton et al., 2020) set the framework for the interpretation of the 1000-year ice core record of stable nitrogen and oxygen isotopes from the high-elevation DML which will be presented in a future study.


## 85 **2 Methods**

### **2.1 Ice core drilling**

A new ice core was drilled as part of the ISOL-ICE project at Kohnen Station, located in DML on the high-elevation East Antarctic Plateau (2892 m a.s.l.) approximately 550 km from the coast. The ISOL-ICE ice core site was located within the clean air sector ~1 km from the deep EDML core, which was drilled in 2001-2006 to a depth of 2761 m (Wilhelms et al.,  
90 2017). The 120 m deep ISOL-ICE ice core (core A; 74.9961 °S, 0.09472 °E; Figure 1) was drilled in January 2017 using the British Antarctic Survey (BAS) shallow electromechanical ice core drill (Mulvaney et al., 2002), which produces a core with a 105 mm internal diameter in a dry borehole. Two additional 10 m firm cores (cores B and C) were recovered from the site using a stainless-steel drill with an internal diameter of 96 mm. A fourth 2 m firm core (core CF) was recovered using a carbon fiber tube with an internal diameter of 98 mm. The core sections were cut into 80 cm lengths and shipped frozen to the BAS  
95 in Cambridge, United Kingdom and stored at -25 °C until core processing. In addition, two snow pits were sampled for soluble



ions and nitrate stable isotopes. Core C was analysed for perfluorocarboxylate. Results from the snow pits and firn core C can be found in Winton et al. (2020) and Garnett et al. (Garnett et al., 2022) respectively.



100 **Figure 1: Map of the ISOL-ICE ice core drilling site in the clean air sector of Kohnen Station in Dronning Maud Land (DML) highlighting the predominate wind direction, deep EDML ice core site, and the ISOL-ICE “ice core” and “flux” sites. The latter is described in Winton et al. (2020). (b) ISOL-ICE “ice core site” showing the main ice core (core A) and shallow firn core (cores B, C and CF) locations. Modified from Winton et al. (2020).**

## 2.2 Ice core processing and glacio-chemical analysis

105 At the BAS, the main ice core (core A) was logged, analysed by dielectric profiling (DEP), and then cut into sections for continuous flow analysis (CFA), analysis of nitrate stable isotopic composition, and archival (Figure A1). Firn cores B and CF were also logged and analysed by DEP. Here we present initial results from the CFA and DEP analysis as a basis for interpreting the nitrate concentration, stable nitrate isotope, and insoluble particle records which will be presented in subsequent studies. The ISOL-ICE ice chemistry data are available through the Polar Data Centre (Winton et al., 2019).

110

Dielectric conductivity of the full ice core was determined by DEP (Oerter et al., 2000; Moore and Paren, 1987; Wilhelms et al., 1998; Hofstede et al., 2004). The DEP data were taken at 250 kHz in 5 mm increments with a 10 mm long measuring



electrode. Cores were measured only after temperature equilibration in the cold laboratory to reduce the effects of temperature-dependent activation energy on the DEP and corrected for minor differences in core temperature measured by a probe inserted  
115 between the electrodes and the ice core. Repeat measurements were made on the main core A between 35 and 39 m and showed excellent reproducibility ( $R^2=0.93$ ,  $P < 0.001$ ;  $RMSE=0.28 \mu S cm^{-1}$ ). Mean background differences between the DEP values in the 3 cores also reflect the different core diameters since the dielectric response measured by the DEP is proportional to the volume of ice between the measurement plates; this is evident around 40 m depth in Figure 2c where several meters of ice were cut with the core cutters set to a smaller diameter. The rising trend observed in Figure 2c reflects the increasing density  
120 of the ice with depth.

A stainless-steel bandsaw was used to cut a 32 x 32 mm ice stick for CFA of trace elements, soluble anions, hydrogen peroxide ( $H_2O_2$ ), insoluble particles and liquid conductivity. CFA sticks were melted vertically on a heated, gold-plated copper melt head, which consists of two rings separating the inner and outer section of the meltwater. The inner section of the CFA  
125 meltwater was used for chemical analysis ensuring no contact of the meltwater with exposed ice core surfaces. The CFA system used was an earlier version of that described by Grieman et al. (2022).

Soluble anions were measured using fast ion chromatography (FIC; ICS-3000, Dionex™). A suite of anions, including  $NO_3^-$ , chloride ( $Cl^-$ ), methanesulfonic acid (MSA;  $CH_3SO_3H$ ) and sulfate ( $SO_4^{2-}$ ), were determined using an AS15-5  $\mu m$  and  
130 AERS500-4 mm suppressor. The sampling resolution of the FIC measurements was 5 cm or 0.3 years at the top of the core and 4 cm or 0.5 years at the bottom. A certified reference material (CRM; European Reference Material ERM-CA408 simulated rainwater) containing  $NO_3^-$ ,  $SO_4^{2-}$  and  $Cl^-$  was measured at the beginning, end and during each laboratory analysis day to determine the accuracy and precision of the soluble anion measurements which are reported in Table A1.


High-resolution measurement of trace elements, including sodium (Na) and magnesium (Mg), were made using inductively coupled plasma-mass spectrometry (ICP-MS; 7700 series, Agilent). A CRM (European Reference Material, ERM-CA616 groundwater) containing Na and Mg was measured at the beginning, end and during the course of each day to determine the accuracy and precision of the trace element measurements which are reported in Table A1. Na and Mg measured by ICP-MS is assumed to represent their ionic form  $Na^+$  and  $Mg^{2+}$  (Grieman et al., 2022).  
140

Liquid conductivity was measured in the CFA fluid line using a digital conductivity meter (Dionex™ 2010i) with a flow through cell (Amber Science, Model 1056).

Insoluble particle counts were performed using a laser particle detector (Abakus, Fa.Klotz, Bad Liebenzell, Germany) which  
145 was connected to the meltwater flow with a flow rate of  $1.6 mL min^{-1}$ . The instrument was set to measure insoluble particles in 32 individual diameter size bins ranging between 0.9 and 50  $\mu m$ . The dust mass was calculated from the measured volume



assuming a spherical particle density of  $2.5 \text{ g cm}^{-3}$ . The sampling resolution of the ICP-MS, liquid conductivity and insoluble particle measurements was  $<1 \text{ mm}$  or  $0.01$  years.



150

**Figure 2.** Volcanic markers identified in the ISOL-ICE core using (a) sulfate concentrations, (b) liquid conductivity, and (c) ice conductivity (dielectric profile; DEP). The unknown volcanic eruptions are documented in the literature for ice cores in the Dronning Maud Land region. DEP data were corrected for temperature but not density. Note the diameter of the core reduced to 98 mm between 38 and 42 m which is reflected in the lower DEP values in the repeated section of the main core A.

### 155 2.3. Ice core dating

The age-depth model of the ISOL-ICE ice core was constructed using annual layer counting constrained by well-dated volcanic eruption horizons documented in the literature based on ice cores in the Dronning Maud Land region (e.g. Sigl et al., 2022). Annual layer counting was based on the high-resolution  $\text{Na}^+$  and  $\text{Mg}^{2+}$  concentrations following previous ice core and aerosol studies in DML (Göktas et al., 2002; Garnett et al., 2022; Ming et al., 2020). Multi-year aerosol observations from Kohnen  
160 Station exhibit a narrow seasonal minimum in  $\text{Na}^+$  and  $\text{Mg}^{2+}$  concentrations during austral summer and a broad peak in late winter/spring (Weller and Wagenbach, 2007). Annual markers (January) were positioned at the sharp, well defined summer  $\text{Na}^+$  and  $\text{Mg}^{2+}$  minima. Volcanic tie points were identified using  $\text{SO}_4^{2-}$  concentrations, liquid conductivity and DEP (Cole-Dai



and Mosley-Thompson, 1999; Zielinski et al., 1994; Langway et al., 1995; Traufetter et al., 2004) and were placed at the depths reported in Table A2 and illustrated in Figure 2.

#### 165 **2.4. Snow accumulation rate and density**

The density of each 80 cm ice core bag was calculated using the ice core mass, diameter and length. Snow accumulation rates were derived from the density measurements and annual layer thickness of the core. Plastic deformation associated with thinning of deeper ice layers was accounted for by using the thinning model of Nye (1963) which resulted in a 10 % correction of the accumulation rate at the base of the core.

#### 170 **2.5. Relationships between accumulation rate, glacio-chemistry and large-scale climate variability**

To understand the transport mechanisms and source regions of moisture, snowfall and aerosols to the ISOL-ICE site, the snow accumulation and glacio-chemistry ice core records were correlated with climate variables over the observational era (1979-2016). Correlations were performed on annual averaged data (January-December), following Thomas et al. (2023), to correspond with the annually-resolved accumulation and MSA records, and seasonally averaged data resulting from the sub-annually-resolved Na<sup>+</sup> and Mg<sup>2+</sup> measurements. Seasonal means were grouped into austral summer-autumn (December to May average) and austral winter-spring (June to November average) for each year using linear interpolation between annual markers. Despite extreme Antarctic precipitation events and nonlinear snowfall throughout a year (Turner et al., 2019), the climatology of monthly mean Na<sup>+</sup> and Mg<sup>2+</sup> concentrations exhibits the same seasonal trend as aerosol observations at Kohnen Station, i.e., a narrow seasonal minimum in austral summer and a broad peak in late winter/spring (Weller and Wagenbach, 2007). These seasonal grouping reflects the known aerosol Na<sup>+</sup> concentration peak in late winter/spring (Weller and Wagenbach, 2007), the distinct ice core Na<sup>+</sup> concentration peak between June and September, the relatively coherent seasonal Na<sup>+</sup> and Mg<sup>2+</sup> concentration patterns and background climate patterns, e.g., the winter transition and sea ice advance season and the summer transition and sea ice retreat season, respectively. Correlations with NO<sub>3</sub><sup>-</sup> concentrations were not calculated due to missing NO<sub>3</sub><sup>-</sup> concentration data in the top 8 m of the record corresponding to the observational era.

185

The records were correlated with select climate variables from ERA5, the fifth generation European Centre for Medium Range Weather Forecast (ECMWF) atmospheric reanalysis data (Hersbach et al., 2020), during the modern satellite observing period 1979-2016. The variables analysed include mean sea level pressure (MSLP), 10 m meridional wind (V10M), 500-hPa geopotential height (Z500) and 500-hPa meridional winds (V500) to investigate regional and large-scale circulation patterns associated with air mass and aerosol transport. The 10 m meridional wind level was chosen to investigate regional scale circulation patterns associated with surface air mass and marine aerosol transport. The 500 hPa level was chosen to capture broader-scale circulation and is reflective of the near-surface winds over the higher elevations of the Antarctic continent, such as at the high-elevation ISOL-ICE site which has an average surface pressure of 675 hPa between 1997 and 2017 (Utrecht University Automatic Weather Station (AWS) at DML05/Kohnen (AWS9;



195 [https://www.projects.science.uu.nl/iceclimate/aws/files\\_oper/oper\\_20632](https://www.projects.science.uu.nl/iceclimate/aws/files_oper/oper_20632), last access: 29 March 2017), but is less influenced  
by localised near-surface boundary layer features such as katabatic winds. The records were also correlated with sea ice  
concentration from the Met Office Hadley Centre Sea Ice and Sea Surface Temperature data set (Rayner et al., 2003; Titchner  
and Rayner, 2014) to investigate relationships between atmospheric transport pathways to the site and areas of sea ice  
anomalies. Lastly, the ISOL-ICE ice core records were correlated with several large-scale climate mode indices, including the  
200 observation-based Southern Annular Mode (SAM) index (Marshall, 2003), and for the El Niño-Southern Oscillation (ENSO)  
we employ the Niño 3.4 index and Southern Oscillation Index (SOI). The SAM index reflects the difference in mean sea level  
pressure anomalies between the southern middle latitudes and Antarctica, whereby positive (negative) SAM index values  
reflect positive (negative) pressure anomalies in middle latitudes, negative (positive) pressure anomalies over Antarctica, and  
a stronger/poleward (weaker/equatorward) mid-latitude jet. The Niño 3.4 index is the sea surface temperature anomaly in the  
205 east-central equatorial Pacific area-averaged over the region 5°N-5°S, 170-120°W, with positive (negative) values associated  
with El Niño (La Niña) conditions. The SOI reflects the atmospheric component of ENSO, and is the difference in mean sea  
level pressure anomalies between Tahiti (central tropical Pacific) and Darwin, Australia (west Pacific warm pool), whereby  
positive (negative) SOI values reflect La Niña (El Niño) conditions. The SAM index was accessed freely online at  
<http://www.nerc-bas.ac.uk/icd/gjma/sam.html>, and the Niño 3.4 index and SOI are from the Climate Prediction Center and  
210 were accessed freely online at <https://www.cpc.ncep.noaa.gov/data/indices> (all last accessed 22 July 2022). Annual averages  
of the SAM, Niño 3.4, and SOI were calculated from the monthly-mean values and are based on the standard 12-month calendar  
year (January-December). The SAM index begins in 1957, and so the correlations were calculated over the common period of  
1957-2016 along with the reanalysis period of 1979-2016.

### 3. Results

#### 215 3.1. Age-depth model

As with previous ice core dating studies in the high-elevation DML region, the ISOL-ICE ice core displayed clear seasonality  
and distinct summer  $\text{Na}^+$  and  $\text{Mg}^{2+}$  troughs throughout the record enabling annual layer counting for the full 120 m record.  
The high-resolution  $\text{Na}^+$  and  $\text{Mg}^{2+}$  annual signals were evaluated simultaneously. The lower resolution soluble anion  
measurements by FIC did not resolve seasonal signals capable for annual layer counting. The  $\text{H}_2\text{O}_2$  data was also evaluated  
220 for summer peaks or winter minima, however the seasonal signal was smoothed below 2 m depth and was not suitable for  
annual layer counting at the low accumulation rate site. Nineteen volcanic time markers were identified from the  $\text{SO}_4^{2-}$   
concentrations, liquid conductivity and DEP. All volcanic horizons were well aligned in the three records (Figure 2). The  
difference between the conductivity measurements of the ice (DEP) and meltwater (liquid conductivity), i.e., the increasing  
trend in the firn section of the DEP record, reflects the change in density with depth compared to the meltwater conductivity  
225 measurements where there is no density effect. The conductivity values from the ice and meltwater measurements agree well  
below ~75 m depth. Annual layer counting of  $\text{Na}^+$  and  $\text{Mg}^{2+}$  concentrations constrained by the volcanic horizons indicates that






the ISOL-ICE ice core spans the last 1349 years from 2017-668 C.E. extending the EDML core by two decades towards the present (Ruth et al., 2007). An age uncertainty of  $\pm 3$  years is estimated at the base of the ice core (Figure A2).

### 3.2. Snow accumulation record

230 The ISOL-ICE snow accumulation record is presented Figure 3a. The mean accumulation rate for the entire core is  $6.5 \pm 2.4$  cm a<sup>-1</sup> (w.e.) consistent with previous estimates of 6.0–7.1 cm a<sup>-1</sup> (w.e.) from ice cores from the high-elevation DML region (Sommer et al., 2000; Oerter et al., 2000; Hofstede et al., 2004). The mean of the 1995-2017 extension of snow accumulation record is  $6.0 \pm 2.7$  cm a<sup>-1</sup> (w.e.) indicating that the accumulation rate has been relatively stable over the last millennium. While the record does not display centennial scale variability in the snow accumulation rate, there is significant interannual variability  
235 in the record ranging between 2 and 20 cm a<sup>-1</sup> (w.e.).

### 3.4. Ice chemistry

The ISOL-ICE ice chemistry records are displayed in Figure 3. Na, Mg and Cl in Antarctic ice cores are derived from marine sea salt either originating from the open ocean or from sea ice sources such as blowing snow or frost flowers (e.g. Abram et al., 2013). The concentrations of Na, Mg and Cl marine aerosols have been stable over the last millennium with mean  
240 concentration values of  $25.4 \pm 18.9$ ,  $2.5 \pm 2.1$  and  $39.0 \pm 23.8$  ng g<sup>-1</sup> respectively. The extended ISOL-ICE record showed that there were less frequent sea salt deposition events over the most recent two decades relative to the last millennium. Sulfate deposited to ice cores is derived from mixed sources including marine primary production, sea salt and volcanic eruptions (e.g. Sigl et al., 2014; Lin et al., 2022). Sulfate concentrations, with a mean value of  $70.6 \pm 37.2$  ng g<sup>-1</sup>, also did not display shift in  
245 the baseline. The ISOL-ICE SO<sub>4</sub><sup>2-</sup> record shows excellent reproducibility with other DML ice cores (Traufetter et al., 2004; Sigl et al., 2022) where the large SO<sub>4</sub><sup>2-</sup> peaks are attributed to volcanic sources (Figure 2). The sole source of MSA is from the oxidation of dimethylsulfide (DMS, (CH<sub>3</sub>)<sub>2</sub>S) from marine phytoplankton emissions to the atmosphere. While sea salt deposition was relatively constant over the last millennium, MSA and NO<sub>3</sub><sup>-</sup> were the only analytes that displayed centennial scale variability with a gradual decrease in concentrations from 668 to 900, stable concentrations from 900 to 1500 and a  
250 notable increase from 1800 to present for MSA and from 1500 to present for NO<sub>3</sub><sup>-</sup>. The centennial scale temporal variability of MSA and NO<sub>3</sub><sup>-</sup> largely covaried in the ISOL-ICE ice core except between 1600 and 1800 when MSA concentrations decreased while NO<sub>3</sub><sup>-</sup> concentrations increased. Concentrations of MSA continued to increase over the extended two decades towards the present of the record. The mean MSA and NO<sub>3</sub><sup>-</sup> values were  $6.7 \pm 2.3$  ng g<sup>-1</sup> and  $47 \pm 13$  ng g<sup>-1</sup> respectively.



255

**Figure 3. Variability in annual snow accumulation rate (a) anion concentrations (b) nitrate ( $\text{NO}_3^-$ ), (c) methanesulfonic acid (MSA), (d) sulfate ( $\text{SO}_4^{2-}$ ), and (e) chloride ( $\text{Cl}^-$ ), and trace element concentrations (f) sodium (Na), and (g) magnesium (Mg) over the last millennium from the ISOL-ICE ice core. Black lines indicate smoothed data using a 0.01 loess regression model (Cleveland and Devlin, 1988). Grey box: two-decade extension of the ice chemistry record from the Dronning Maud Land site. Water equivalent: w.e. Common era: C.E.**

260



### 3.4. Meteorological conditions and spatial correlations

For the period 1957-2016, there are weak, but statistically significant at the 10 % level ( $P < 0.10$ ), annual correlations found with SAM and  $\text{Na}^+$  ( $R = -0.32$ ) and accumulation ( $R = 0.27$ ), and for the SOI and  $\text{SO}_4^{2-}$  ( $R = -0.25$ ), i.e., La Niña (El Niño) is associated with lower (higher)  $\text{SO}_4^{2-}$  concentrations. The SAM relationships weaken and become insignificant after 1979, while the ENSO relationship with  $\text{SO}_4^{2-}$  strengthens after 1979 (Table A3). There were no other statistically significant annual relationships with the ice core record and SAM or ENSO.

Figure 4a shows the annual correlations for snow accumulation with 500 hPa geopotential height and meridional winds, 10 m meridional wind, mean sea level pressure and sea ice concentration for the period 1979-2016. Higher annual snow accumulation is associated with anomalous low pressure (anomalous cyclonic/clockwise circulation) over the Weddell Sea (significant in mean sea level pressure field) and northerly flow from the South Atlantic (significant at the 10 m level) bringing marine airmasses to the site (i.e., the negative correlation between meridional wind and accumulation implies positive accumulation anomalies are associated with northerly wind anomalies). The anomalous northerly flow corresponds with negative sea ice concentration anomalies in the outer edge of the sea ice field, with significant correlations in the northern Weddell Sea. While the northerly flow would directly contribute to a decrease in sea ice in the northern Weddell Sea through melt and/or convergence, it is also possible that the increase in open water could locally enhance atmospheric moisture content and contribute to higher snowfall.


Enhanced MSA concentrations are associated with an anomalous high in the South Atlantic, an offshore low near  $15^\circ\text{E}$ , and an offshore high near  $90^\circ\text{E}$ . In between the latter two pressure anomalies there is significant northerly flow onto the ice sheet between  $30\text{--}45^\circ\text{E}$  that circulates clockwise and flows offshore over the ISOL-ICE site due to the strong high-pressure anomaly in the South Atlantic. This onshore/northerly flow to the northeast of the site cuts across a local region of below average sea ice concentrations within the pack ice near Prydz Bay ( $\sim 70^\circ\text{E}$ ), with mostly above average sea ice concentrations over the Weddell Sea where southerly flow prevails.

Seasonal correlations for  $\text{Mg}^{2+}$  and  $\text{Na}^+$  are similar (Figure 4c-f) as they are strongly positively correlated ( $R = 0.65$  for summer-autumn and  $R = 0.97$  for winter-spring). In winter, strong correlation and similar correlation maps in Figures 5e-f) suggest that  $\text{Na}^+$  and  $\text{Mg}^{2+}$  are sea salt aerosol tracers from the open ocean or the sea ice and are advected with the same air masses (e.g. Wagenbach et al., 1998; Vega et al., 2018), whereas in summer the weaker correlation could be due to mineral dust contributions. During the summer-autumn period high  $\text{Mg}^{2+}$  and  $\text{Na}^+$  is associated with a circumpolar zonal wave 3 pattern (Raphael, 2004) with three high-low pressure pairs encircling Antarctica. The zonal wave 3 pattern is most pronounced for Mg. Within this pattern, there is anomalous northerly/onshore flow between  $15\text{--}30^\circ\text{E}$  bringing marine airmasses to the site. For Na, the onshore flow is tied to a local anticyclone just off the Enderby Land coast (near  $45^\circ\text{E}$ ), and for Mg the onshore



295 flow is tied to a local cyclone just off the DML coast (near 0°E). There is also significant northerly flow over the South Pacific associated with a cyclone north of the Ross Sea. The northerly flow from the South Pacific appears to cut across the West Antarctic Ice Sheet and flow offshore (turns to southerly wind anomalies/positive correlations) over the eastern Ronne-Filchner Ice Shelf (near 45°W), especially for Mg. This transport pathway aligns with reduced sea ice concentrations in the Amundsen Sea which could contribute to enhanced Mg concentrations in summer-autumn.

300 During winter-spring, the large-scale circulation pattern and regional pressure anomalies associated with Na<sup>+</sup> and Mg<sup>2+</sup> are weaker and less coherent compared to summer-autumn. There appears to be two potential sources of onshore flow that bring marine air masses to the site in winter-spring. First, high Mg<sup>2+</sup> and Na<sup>+</sup> concentrations are associated with northerly flow over the Bellingshausen Sea/Antarctic Peninsula that cuts across Ellsworth Land and then flows off the continent along 0°E, which is weakly tied to an anticyclone north of the Weddell Sea and slightly lower-than-average pressure over the Amundsen Sea  
305 region. Second, similar to the annual MSA correlations, there is a local cyclonic/clockwise circulation anomaly centred on the East Antarctic plateau near 10°E, which produces onshore northerly flow near 45°E that travels clockwise around the cyclone across the South Pole and reaches the ISOL-ICE site on southerly winds. The local southerly flow is enhanced by the weak positive pressure/anticyclonic circulation anomaly north of the Weddell Sea. There is a negative correlation between Mg<sup>2+</sup> and Na<sup>+</sup> concentrations and sea ice concentrations in the Weddell Sea around 0° yet this sea salt source region would not be tied  
310 to the ISOL-ICE ice core site as it is associated with offshore/southerly flow.



315 **Figure 4.** Annual and seasonal correlations during the observational era (1979–2016) between the ISOL-ICE ice core record and sea ice concentration (SIC), 500-hPa geopotential height (Z500), 500-hPa meridional wind (V500), mean sea level pressure (MSLP), and 10 m meridional wind (V10M). Correlations are for (a) annual accumulation, (b) annual MSA concentration, (c) summer-autumn sodium concentration, (d) summer-autumn magnesium concentration, (e) winter-spring sodium concentration, and (f) winter-spring magnesium concentration. Coloured shading shows Pearson's correlation coefficient values as indicated by the colour bar at the bottom. The bold black contours denote correlations that are significant at the 10 % level based on a two tailed Students t-test.

## 4. Discussion

### 320 4.1. Aerosol source regions

The winter sea salt peak in the ISOL-ICE ice core is dominated by a northerly atmospheric transport signal, that is northerly flow directly to the site or northerly flow in other Antarctic sectors with transport across the continent to the site, rather than



an open water/sea ice source region signal. In the case of the ISOL-ICE site, direct transport of sea salt aerosol is most likely as wet deposition of aerosol is connected to lifting and adiabatic cooling of the air mass. Comparison of ice core  $\text{SO}_4^{2-}/\text{Na}^+$  ratios with the reference sea water ratio of 0.252 (Millero et al. (2008)) and those from fractionated sea ice sources ( $< 0.252$ ) may allow attribution of sea salt aerosol source regions assuming there are two sea salt sources from open ocean and sea ice regions (Abram et al., 2013). This approach is limited because only annual  $\text{SO}_4^{2-}/\text{Na}^+$  ratios are retrieved from the ISOL-ICE core, yet the seasonally resolved snow pit reported by Winton et al. (2020) shows that the winter  $\text{Na}^+$  maximum at the ISOL-ICE site is associated with lower  $\text{SO}_4^{2-}/\text{Na}^+$  ratios (Figure A3). At the same time, nss- $\text{SO}_4^{2-}$  concentrations remain positive (Figures A3 and 6a). Thus, the ISOL-ICE winter  $\text{SO}_4^{2-}/\text{Na}^+$  ratios  $> 0.252$  and the positive nss- $\text{SO}_4^{2-}$  values both indicate enrichment compared to seawater. However, in winter  $\text{SO}_4^{2-}/\text{Na}^+$  values decrease due to contributions from sea ice sources of sea salt aerosol such as blowing snow in the Antarctic (Figure A3) (Frey et al., 2020). At coastal Antarctic sites, such as Halley Station, the winter peak in  $\text{Na}^+$  originates from sea salt aerosol production from blowing snow above sea ice in the Weddell Sea with minor contributions from sea-ice production and frost flowers in coastal leads in the Weddell Sea (Frey et al., 2020; Yang et al., 2019). The evidence for a sea ice source of sea salt at inland, high-elevation sites is harder to substantiate due to the higher nss- $\text{SO}_4^{2-}$  background levels from other sources relative to sea salt concentrations and thus negative values of nss- $\text{SO}_4^{2-}$  are often not observed (Abram et al., 2013). Analysis of year-round  $\text{SO}_4^{2-}$  and  $\text{Na}^+$  aerosol data from Kohnen Station also could not differentiate between open ocean and sea ice sources based on the fractionation of winter sea salt aerosol and its origin from a potential sea ice source, i.e., winter sea salt- $\text{SO}_4^{2-}$  makes a minor contribution to the annual total which is dominated by large summer nss- $\text{SO}_4^{2-}$  (Weller and Wagenbach, 2007). Therefore, the sea salt fractionation cannot be used as a chemical signature of sea ice source strength at the ISOL-ICE site because the nss- $\text{SO}_4^{2-}$  contribution to the annual mean is so large. Whereas recent changes in the proximity of the ice front to ice core sites in coastal DML highlighted the important role of elevation of an ice core site on the sea salt record as observed from the calving of Trolltunga in 1950 which increased coastal sea salt concentrations and shifted nss- $\text{SO}_4^{2-}$  towards more negative values in a low elevation ice core (Vega et al., 2018). Yet the changing sea salt source region was not reflected in a higher elevation ice rise site in the region showing that, even proximal to the coast, elevation plays a role in the detection of a sea salt fractionation signal in the DML region. While the source of winter  $\text{Na}^+$  at ISOL-ICE may be less certain, the correlation analysis can explain the northerly transport to the site.

In summer, sea salt deposition at the ISOL-ICE site is explained by a transport signal associated with northerly marine air masses from Enderby Land and DML and an open water aerosol source in the Amundsen Sea with atmospheric transport across West Antarctica. The low sea salt concentrations in summer occur with more enriched  $\text{SO}_4^{2-}/\text{Na}^+$  values, and higher nss- $\text{SO}_4^{2-}$  and MSA concentrations ( $R=0.15$ ,  $P < 0.01$ ) indicating biogenic  $\text{SO}_4^{2-}$  aerosol from open water (Figure A3). The observed correlation is weak but statistically significant, meaning that the parameters are somehow related but the predictive skill to estimate one parameter from the other at a given point in time is very low. Taken together, at the seasonal scale, the correlation analysis and high-resolution ice chemistry provide evidence for sea salt variability driven by year-round northerly



atmospheric flow, either directly to the site or from other regions of the continent that are transported to the site, with an additional contribution from an open ocean derived sea salt in summer rather than a sea ice source.

360 Over the last millennium, unfractionated sea salt from open ocean sources was relatively constant;  $\text{SO}_4^{2-}/\text{Na}^+$  ratios remained  
enriched with a mean  $\text{SO}_4^{2-}/\text{Na}^+$  of  $3.7 \pm 2.7$  above the seawater value of 0.252 (Figure 5). A lack of  $\text{SO}_4^{2-}$  depletion in the  
ISOL-ICE ice core from fractionated sea ice sources, likely dominated by blowing snow and contributions from frost flowers,  
is similarly not observed at other inland, high-elevation Antarctic ice core sites which is attributed to the low accumulation  
365 rate and low temporal resolution smoothing the winter signal, and high summer biogenic  $\text{SO}_4^{2-}$  overriding the depleted  $\text{SO}_4^{2-}$   
signal (Rankin et al., 2000; Rankin et al., 2004). Furthermore, nss- $\text{SO}_4^{2-}$  from the open ocean does reach Kohlen Station in  
winter (Weller and Wagenbach, 2007). There are three notable periods in the ISOL-ICE sea salt chemistry record (Figures 4  
and 6). The first is around 1200 where there are no extreme sea salt deposition events and  $\text{SO}_4^{2-}/\text{Na}^+$  ratios become more  
enriched. The second is around 1600 where the frequency of sea salt deposition events increase. The third is from 1800 to  
370 present where there is an increase in the frequency of sea salts deposition events (Na, Cl and Mg concentrations) at the same  
time as the increase in MSA concentrations. Based on the sea salt-northerly wind relationship for the 1979-2016 period (Figure  
4), the increase in extreme sea salt deposition years in the ISOL-ICE record likely reflects stronger sea salt transport rather  
than changing open water/sea ice conditions. We acknowledge that these relationships may change over time. For example,  
the SAM influence on Antarctic temperature and wind changed between 1957-1979 compared to 1979-present (Marshall et  
al., 2022). While the large-scale drivers of northerly wind may change, the sea salt-northerly wind relationship may hold true  
375 over the last millennium. Fundel et al. (2006) investigated MSA variability in another core from the high-elevation Dronning  
Maud Land (DML05) in relation to atmospheric patterns. The authors found that over the reanalysis period (1969-1997), the  
Antarctic Dipole plays an important role during years of particularly high MSA concentrations, while over the last 2000 years,  
periods of high MSA concentrations are related to periods of higher sea-salt aerosol reflecting the influence of transport on  
both species despite the different seasonality. Atmospheric transport strength, rather than sea ice conditions, was also found  
380 by Abram et al. (2007) as the dominant factor determining the MSA signal preserved in ice cores in the Weddell Sea sector.  
While longer-term trends in MSA concentrations are not coupled with  $\text{SO}_4^{2-}/\text{Na}^+$  ratios or sea salt variations throughout the  
ISOL-ICE record, transport clearly plays a dominant role in both records with the Antarctic Dipole likely aiding in the efficient  
atmospheric transport of both sea salt and MSA to the site during periods for example, from 1200 to 1600 C.E.

385 The ISOL-ICE core reproduces the MSA and Cl records from existing DML cores (Fundel et al., 2006) and extends these  
records by two decades from 2000 to 2017. The striking feature of the extended MSA record is the continued increase in MSA  
concentration over the last two decades starting around 1800 (Figure 3). While MSA concentrations continue to increase over  
the last two decades, years of extreme sea salt deposition do not. Both changes in the seasonality and extent of regional sea  
ice, which impacts the source region and development of seasonal phytoplankton blooms in Antarctic waters (e.g. Abram et  
390 al., 2011), a change in DMS oxidation pathway, and a change in atmospheric circulation strength and pathway could contribute



to the MSA increase over the last two centuries. During this same period of MSA increase, reconstructions of sea ice conditions from Antarctic ice cores show that sea ice extent declined in the Bellingshausen-Amundsen Sea region since 1850 C.E. (Dalaiden et al., 2021; Abram et al., 2010; Thomas et al., 2019) and the northernmost latitude of sea ice edge declined in the Weddell Sea sector (50°W–20°E) over the reconstructed period from 1900 to present (Yang et al., 2021). These recent changes  
395 call for an updated analysis of MSA-sea ice relationships over the past two decades towards the present.

While previous studies found that MSA variability in the high-elevation DML region was associated with transport strength rather than the sea ice conditions (Fundel et al., 2006; Abram et al., 2007), we investigated whether the additional two decades of MSA ice core data from ISOL-ICE captures local sea ice changes. Recently, Isaacs et al. (2021) and Clem et al. (2018) found  
400 that over the period 1979–2018, ENSO impacted summer and autumn sea ice concentration around coastal DML (10–70°E) where sea ice concentration increased during a cold ENSO phase and decreased during a warm ENSO phase. Here, sea surface temperature anomalies in the tropical Pacific Ocean influenced regional circulation over the South Atlantic and encouraged meridional airflow to DML affecting sea ice thermodynamically, by altering local heat transport and hence sea ice formation and melt. In the region where sea ice concentration is significantly correlated with ENSO, the sea ice concentration anomaly  
405 increased from 1979 to 2016 consistent with the pan Antarctic sea ice concentration trend, and also demonstrated that interannual variability was strongest in austral winter and spring where it was more subdued in summer and autumn when sea ice concentration was most strongly correlated with ENSO. Our correlation analysis (1979–2016) shows that years of higher annual ice core MSA concentrations are associated with reduced sea ice concentrations west of Prydz Bay and this MSA aerosol is delivered to the ISOL-ICE ice core site by northerly distal winds. We did not find evidence for a locally derived  
410 MSA aerosol source in coastal DML (Figure 4b). Nor did we find a relationship between ENSO and MSA in the ISOL-ICE record. Thus, our updated record shows that years of increased ice core MSA were likely driven by the transport direction with meridional winds that flow across a reduced sea ice region in the Prydz Bay region acting to synergistically reduce sea ice concentration while also increasing the delivery of MSA to the ISOL-ICE ice core site. Assuming the MSA-wind direction relationship holds true for the last millennium, the increase in MSA since 1880 likely reflects stronger northerly transport over  
415 that period. Our correlation analysis, combined with our finding that sea salts do not carry a fractionated sea ice signature over the last 1300 years, shows that sea salt aerosol and MSA concentrations continue to be primarily related to atmospheric transport over the extended two-decade period.


While the variability of  $\text{NO}_3^-$  concentration in the ISOL-ICE ice core will be discussed in a separate study, the significance of  
420 constraining aerosol source regions has implications for ice core  $\text{NO}_3^-$ . While  $\text{NO}_3^-$  sources at remote high elevation Antarctic sites are not well constrained, the net primary sources in the Southern Ocean and Antarctica are peroxyacetyl nitrate (PAN) and methyl nitrate ( $\text{CH}_3\text{NO}_3$ ) from open ocean and sea ice and stratospheric  $\text{NO}_3^-$  (Burger et al., 2022; Burger et al., 2023; Mulvaney and Wolff, 1993; Jones et al., 2011). The similar temporal trends of  $\text{NO}_3^-$  and MSA compared to sea salts over the last millennium suggests a potential link between MSA and  $\text{NO}_3^-$  through the contribution of oceanic  $\text{CH}_3\text{NO}_3$  to ice core  $\text{NO}_3^-$





425 , i.e., the weak but statistically significant correlation between ice core  $\text{NO}_3^-$  and MSA ( $R=0.10$ ,  $P < 0.01$ ) is similar to  $\text{SO}_4^{2-}$  and MSA ( $R=0.15$ ,  $P < 0.01$ ) where the latter relationship is traditionally used to assess biogenic sulfur contributions from the ocean.

Previous studies from the Antarctic coast and Weddell Sea region provide mixed evidence for a  $\text{CH}_3\text{NO}_3$  source. For example, recent studies attribute some of the atmospheric  $\text{NO}_3^-$  observed in the Weddell Sea boundary layer to oceanic alkyl nitrates including  $\text{CH}_3\text{NO}_3$  (Burger et al., 2023) in addition to alkyl amide emissions from algae in the marginal sea ice zone around the Antarctic Peninsula (Brean et al., 2021; Dall'osto et al., 2019; Dall'osto et al., 2017). At Halley Station, Jones et al. (2011) observed atmospheric organic  $\text{NO}_3^-$  (PAN and  $\text{CH}_3\text{NO}_3$ ), dominate  $\text{NO}_y$  in winter, and with a seasonality consistent with an oceanic source. Weller et al. (2002) showed similar seasonality of  $\text{NO}_y$  at Neumayer Station, yet they did not find any significant contribution from the marine boundary layer based on trajectory analyses and radioisotope measurements. However, at Halley Station, year-round  $\text{NO}_3^-$  in surface snow correlated with inorganic  $\text{NO}_y$  and not with  $\text{CH}_3\text{NO}_3$  (Jones et al., 2011). The correlation may be different further from the source on the Antarctic plateau as suggested by the MSA and  $\text{NO}_3^-$  correlation in the ISOL-ICE ice core (full record:  $R=0.10$ ,  $P < 0.01$ ; record excluding 1600-1800:  $R=0.22$ ,  $P < 0.01$ ). As there are no  $\text{NO}_y$  observations in high-elevation Droning Maud Land, further field measurements and modelling studies are required to understand the relationship, to confirm an oceanic alkyl nitrate/ $\text{CH}_3\text{NO}_3$  contribution to ice core  $\text{NO}_3^-$  and whether these organic source changes are imprinted in the  $\text{NO}_3^-$  concentration record, and to understand why the ice core MSA and  $\text{NO}_3^-$  relationship is intermittent i.e., between 1600-1800 when MSA and  $\text{NO}_3^-$  do not covary ( $R=0.24$ ,  $P < 0.01$ ). While sea salt fractionation at the ISOL-ICE ice core site cannot be used as a chemical signature of sea ice source strength, sea salt concentrations show that oceanic aerosol sources have remained stable over the last millennium and the relationship between sea salts and reanalysis data products implies that there have been no large changes in atmospheric transport over the period. Therefore, trends in ice core  $\text{NO}_3^-$  and its stable isotopic composition likely reflect changes in atmospheric processes such as photochemical recycling of  $\text{NO}_3^-$  between the air-snow interface.



450 **Figure 5.** Variability in ice core anions (a) non sea salt sulfate ( $\text{nss-SO}_4^{2-}$ ) concentration, and (b)  $\text{SO}_4^{2-}/\text{Na}^+$  ratio over the last  
460 millennium from the ISOL-ICE ice core. The orange dashed line indicates the  $\text{SO}_4^{2-}/\text{Na}^+$  ratio of reference seawater. Above this line  
ice core  $\text{SO}_4^{2-}/\text{Na}^+$  values represent open ocean sources in dark blue and below the line represents frost flower/sea ice sources of sea  
salt (Rankin et al., 2000). Positive  $\text{nss-SO}_4^{2-}$  values in light blue indicate contributions from open ocean and sea ice margins, and  
465 negative values indicate contributions from sea ice sources from blowing snow and frost flowers. The  $\text{nss-SO}_4^{2-}$  contains all the  
volcanic peaks identified earlier for dating the ice core. Common era: C.E.

#### 4.2 Accumulation rate and implications for the nitrate isotope record

Snow accumulation at ISOL-ICE originates predominantly from the South Atlantic with onshore winds delivering marine air masses to the site (Figure 4) and the majority of precipitation occurring from frontal clouds (Reijmer and Oerlemans, 2002; Welker et al., 2014). This is consistent with air mass back trajectory studies by Reijmer et al. (2002) who show that air parcels and precipitation events at the Kohnen Station site originate in the Southern Atlantic Ocean around four days prior to their arrival on the DML plateau. Assuming the relationship between annual snow accumulation and winds persisted over the last millennium, years of high snow accumulation in the ISOL-ICE record were driven by northerly atmospheric flow from the South Atlantic. While the accumulation rate has been relatively stable over the last millennium and the record does not display centennial scale variability, the significant interannual variability in the record ranging between 2 and 20  $\text{cm a}^{-1}$  (w.e.) needs to be accounted for when interpreting the ice core  $\delta^{15}\text{N}(\text{NO}_3^-)$  record which in turn is sensitive to changes in the accumulation rate and seasonality at the site (Winton et al., 2020). Air snow transfer modelling indicates that  $\delta^{15}\text{N}(\text{NO}_3^-)$  preserved in the ISOL-ICE ice core will be less sensitive to changes in surface-UV than that retrieved at the lower snow accumulation site of



470 Dome C. However, the higher snow accumulation rate and more accurate dating of the ISOL-ICE ice core allows for higher-resolution  $\delta^{15}\text{N}(\text{NO}_3^-)$  based reconstructions of past surface-UV (TCO) (Winton et al., 2020). Given the snow accumulation rate falls within the range proposed for reconstructing surface mass balance from ice core  $\delta^{15}\text{N}(\text{NO}_3^-)$  (Akers et al., 2022a), future work will investigate the possibility of removing the accumulation signature of the  $\delta^{15}\text{N}(\text{NO}_3^-)$  record to reconstruct past surface-UV radiation at the site.

## 5. Summary

475 In this study, we present the depth-age scale, accumulation rate and initial ice chemistry data from the 120 m deep ISOL-ICE ice core drilled at Kohnen Station in DML on the high-elevation Plateau in January 2017 to set the framework for the interpretation of the ISOL-ICE ice core record of stable nitrogen and oxygen isotopes in nitrate. The age scale was derived using well established dating techniques for ice cores in the region. The ISOL-ICE ice core spans the last 1349 years from 668 to 2017 C.E.  $\pm 3$  years extending the EDML core by 20 years. The mean snow accumulation rate of  $6.5 \pm 2.4 \text{ cm a}^{-1}$  (w.e.) was stable over the last millennium, consistent with other ice cores in the region, with no notable changes over the last two decades. 480 While the snow accumulation record does not display longer temporal trends over the last millennium, the significant interannual variability in the record needs to be accounted for when interpreting the nitrate stable isotope record which is sensitive to changes in the accumulation rate and seasonality at the site. Correlation of the snow accumulation record with climate variables over the observational era (1979-2016) showed that snow accumulation at ISOL-ICE is predominately derived from the South Atlantic with onshore winds delivering marine air masses to the site.

485 The ISOL-ICE core replicated other high-elevation DML ice chemistry records over the last millennium that were recovered decades earlier highlighting the reproducibility of climate and aerosol records in the region. Deposition of marine sea salts (Na, Mg and Cl) to the ISOL-ICE ice core site have remained stable over the last millennium. While the centennial scale variability of MSA and  $\text{NO}_3^-$  concentrations largely covaried and increased over the last few centuries. The extended ISOL-ICE record of last two decades showed that MSA continued to increase while there were less frequent deposition events of sea salts relative to the last millennium. Our correlation analysis, combined with our finding that sea salts do not carry a fractionated sea ice signature over the last 200 years, shows that sea salt aerosol and MSA concentrations continue to be primarily related to atmospheric transport over the extended two-decade period. A potential link with MSA and  $\text{NO}_3^-$  ice core variability warrants future work to investigate an oceanic alkyl nitrate/ $\text{CH}_3\text{NO}_3$  contribution to ice core  $\text{NO}_3^-$ . As both sea salt 495 deposition, atmospheric transport and snow accumulation rates remained stable over the last millennium, these factors are unlikely to drive the temporal variability on the observed 1300-year ISOL-ICE ice core record of  $\text{NO}_3^-$  and its stable isotopic composition.



**Data availability** The datasets for the ISOL-ICE age-depth model, accumulation rate, DEP, and ice chemistry are available through the Polar Data Centre at <https://doi.org/10.5285/9c972cfb-0ffa-4144-a943-da6eb82431d2> (Unreleased data) (Winton et al., 2019).

**Author contribution** Conceptualisation, M.M.F., J.S. and V.H.L.W.; methodology, M.M.F., R.M. and V.H.L.W.; validation, V.H.L.W.; formal analysis, V.H.L.W.; investigation, all authors.; resources, M.M.F. and R.M.; data curation, V.H.L.W.; writing - original draft preparation, V.H.L.W.; writing - review and editing, all authors; funding acquisition, M.M.F., J.S. and R.M. All authors have read and agreed to the published version of the manuscript.

**Competing interests** The authors declare that they have no conflict of interest.

**Acknowledgments** We would like to thank the British Antarctic Survey (BAS) and Alfred Wegener Institute (AWI) staff for their field and logistical support at Halley Station and Kohnen Station, respectively. Technical support for ice core measurements at BAS were provided by Rebecca Tuckwell, Lisa Hauge, Julius Rix, Catriona Sinclair, Emily Ludlow, and Shaun Miller. The ISOL-ICE ice core project was funded by a Natural Environment Research Council (NERC) Standard Grant (NE/N011813/1) to M.M.F. V.H.L.W. was supported by the NERC Standard Grant, and a Rutherford Foundation Postdoctoral Fellowship (RTF-VUW1801-PD) and Rutherford Discovery Fellowship (RDF-VUW2203) administered by the Royal Society Te Apārangi. K.R.C. acknowledges support from the Royal Society Te Apārangi Marsden Fund (MFP-VUW2010). JS thanks the French polar institute IPEV for its continuous support of field deployments and ice, snow and atmospheric sampling under the CAPOXI program (grant # 1177) and the European Union (grant 101054558 DOC-PAST). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or European Research Council (ERC). Neither the European Union nor the ERC can be held responsible for them.



## References

- Abram, N. J., Mulvaney, R., and Arrowsmith, C.: Environmental signals in a highly resolved ice core from James Ross Island, Antarctica, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- 525 Abram, N. J., Wolff, E. W., and Curran, M. A.: A review of sea ice proxy information from polar ice cores, *Quaternary Science Reviews*, 79, 168-183, 2013.
- Abram, N. J., Mulvaney, R., Wolff, E. W., and Mudelsee, M.: Ice core records as sea ice proxies: An evaluation from the Weddell Sea region of Antarctica, *J. Geophys. Res.*, 112, D15101, 10.1029/2006jd008139, 2007.
- Abram, N. J., Thomas, E. R., McConnell, J. R., Mulvaney, R., Bracegirdle, T. J., Sime, L. C., and Aristarain, A. J.: Ice core evidence for a  
20th century decline of sea ice in the Bellingshausen Sea, Antarctica, *Journal of Geophysical Research: Atmospheres*, 115, 2010.
- 530 Akers, P. D., Savarino, J., Caillon, N., Servettaz, A. P., Le Meur, E., Magand, O., Martins, J., Agosta, C., Crockford, P., and Kobayashi, K.:  
Sunlight-driven nitrate loss records Antarctic surface mass balance, 2022a.
- Akers, P. D., Savarino, J., Caillon, N., Servettaz, A. P., Le Meur, E., Magand, O., Martins, J., Agosta, C., Crockford, P., and Kobayashi, K.:  
Sunlight-driven nitrate loss records Antarctic surface mass balance, *Nature Communications*, 13, 4274, 2022b.
- Berhanu, T. A., Meusinger, C., Erbland, J., Jost, R., Bhattacharya, S., Johnson, M. S., and Savarino, J.: Laboratory study of nitrate photolysis  
535 in Antarctic snow. II. Isotopic effects and wavelength dependence, *The Journal of chemical physics*, 140, 244306, 2014.
- Beyersdorf, A. J., Blake, D. R., Swanson, A., Meinardi, S., Rowland, F., and Davis, D.: Abundances and variability of tropospheric volatile  
organic compounds at the South Pole and other Antarctic locations, *Atmospheric Environment*, 44, 4565-4574, 2010.
- Bodeker, G. E., Nitzbon, J., Tradowsky, J. S., Kremser, S., Schwertheim, A., and Lewis, J.: A global total column ozone climate data record,  
*Earth System Science Data*, 13, 3885-3906, 2021.
- 540 Brean, J., Dall'Osto, M., Simó, R., Shi, Z., Beddows, D. C. S., and Harrison, R. M.: Open ocean and coastal new particle formation from  
sulfuric acid and amines around the Antarctic Peninsula, *Nature Geoscience*, 14, 383-388, 10.1038/s41561-021-00751-y, 2021.
- Brönnimann, S., Staehelin, J., Farmer, S., Cain, J., Svendby, T., and Svenøe, T.: Total ozone observations prior to the IGY. I: A history,  
*Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical  
oceanography*, 129, 2797-2817, 2003.
- 545 Burger, J. M., Joyce, E., Hastings, M. G., Spence, K. A., and Altieri, K. E.: A seasonal analysis of aerosol NO<sub>3</sub>– sources and NO<sub>x</sub> oxidation  
pathways in the Southern Ocean marine boundary layer, *Atmospheric Chemistry and Physics*, 23, 5605-5622, 2023.
- Burger, J. M., Granger, J., Joyce, E., Hastings, M. G., Spence, K. A., and Altieri, K. E.: The importance of alkyl nitrates and sea ice emissions  
to atmospheric NO<sub>x</sub> sources and cycling in the summertime Southern Ocean marine boundary layer, *Atmospheric Chemistry and Physics*,  
22, 1081-1096, 2022.
- 550 Cao, Y., Jiang, Z., Alexander, B., Cole-Dai, J., Savarino, J., Erbland, J., and Geng, L.: On the fingerprint of the Antarctica ozone hole in ice  
core nitrate isotopes: a case study based on a South Pole ice core, *Atmospheric Chemistry and Physics Discussions*, 1-23, 2022.
- Clem, K. R., Renwick, J. A., and McGregor, J.: Autumn cooling of western East Antarctica linked to the tropical Pacific, *Journal of  
Geophysical Research: Atmospheres*, 123, 89-107, 2018.
- Cleveland, W. S. and Devlin, S. J.: Locally weighted regression: an approach to regression analysis by local fitting, *Journal of the American  
555 Statistical Association*, 83, 596-610, 1988.
- Cole-Dai, J. and Mosley-Thompson, E.: The Pinatubo eruption in South Pole snow and its potential value to ice-core paleovolcanic records,  
*Annals of Glaciology*, 29, 99-105, 1999.
- Dalaiden, Q., Goosse, H., Rezsöhazy, J., and Thomas, E. R.: Reconstructing atmospheric circulation and sea-ice extent in the West Antarctic  
over the past 200 years using data assimilation, *Climate Dynamics*, 57, 3479-3503, 2021.
- 560 Dall'Osto, M., Airs, R. L., Beale, R., Cree, C., Fitzsimons, M. F., Beddows, D., Harrison, R. M., Ceburnis, D., O'Dowd, C., and Rinaldi,  
M.: Simultaneous detection of alkylamines in the surface ocean and atmosphere of the Antarctic sympagic environment, *ACS Earth and  
Space Chemistry*, 3, 854-862, 2019.
- Dall'Osto, M., Ovadnevaite, J., Paglione, M., Beddows, D. C., Ceburnis, D., Cree, C., Cortés, P., Zamanillo, M., Nunes, S. O., and Pérez,  
G. L.: Antarctic sea ice region as a source of biogenic organic nitrogen in aerosols, *Scientific Reports*, 7, 1-10, 2017.
- 565 Erbland, J., Vicars, W., Savarino, J., Morin, S., Frey, M., Frosini, D., Vince, E., and Martins, J.: Air–snow transfer of nitrate on the East  
Antarctic Plateau–Part 1: Isotopic evidence for a photolytically driven dynamic equilibrium in summer, *Atmospheric Chemistry and Physics*,  
13, 6403-6419, 2013.
- Frey, M. M., Savarino, J., Morin, S., Erbland, J., and Martins, J.: Photolysis imprint in the nitrate stable isotope signal in snow and atmosphere  
of East Antarctica and implications for reactive nitrogen cycling, *Atmospheric Chemistry and Physics*, 9, 8681-8696, 2009.
- 570 Frey, M. M., Norris, S. J., Brooks, I. M., Anderson, P. S., Nishimura, K., Yang, X., Jones, A. E., Nerentorp Mastromonaco, M. G., Jones,  
D. H., and Wolff, E. W.: First direct observation of sea salt aerosol production from blowing snow above sea ice, *Atmospheric Chemistry  
and Physics*, 20, 2549-2578, 2020.
- Fundel, F., Fischer, H., Weller, R., Traufetter, F., Oerter, H., and Miller, H.: Influence of large-scale teleconnection patterns on methane  
sulfonate ice core records in Dronning Maud Land, *Journal of Geophysical Research: Atmospheres*, 111, 2006.



- 575 Garnett, J., Halsall, C., Winton, H., Joerss, H., Mulvaney, R., Ebinghaus, R., Frey, M., Jones, A., Leeson, A., and Wynn, P.: Increasing Accumulation of Perfluorocarboxylate Contaminants Revealed in an Antarctic Firm Core (1958–2017), *Environmental Science & Technology*, 10.1021/acs.est.2c02592, 2022.
- Göktas, F., Fischer, H., Oerter, H., Weller, R., Sommer, S., and Miller, H.: A glacio-chemical characterization of the new EPICA deep-drilling site on Amundsenisen, Dronning Maud Land, Antarctica, *Annals of Glaciology*, 35, 347-354, 2002.
- 580 Grieman, M. M., Hoffmann, H. M., Humby, J. D., Mulvaney, R., Nehrbass-Ahles, C., Rix, J., Thomas, E. R., Tuckwell, R., and Wolff, E. W.: Continuous flow analysis methods for sodium, magnesium and calcium detection in the Skytrain ice core, *Journal of Glaciology*, 68, 90-100, 2022.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., and Schepers, D.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999-2049, 2020.
- 585 Hofstede, C. M., van de Wal Roderik, S., Kaspers, K. A., Van Den Broeke, M. R., Karlöf, L., Winther, J.-G., Isaksson, E., Lappégard, G., Mulvaney, R., and Oerter, H.: Firm accumulation records for the past 1000 years on the basis of dielectric profiling of six cores from Dronning Maud Land, Antarctica, *Journal of Glaciology*, 50, 279-291, 2004.
- Isaacs, F. E., Renwick, J. A., Mackintosh, A. N., and Dadic, R.: ENSO modulates summer and autumn sea ice variability around Dronning Maud Land, Antarctica, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033140, 2021.
- 590 Jacobi, H.-W., Weller, R., Jones, A., Anderson, P., and Schrems, O.: Peroxyacetyl nitrate (PAN) concentrations in the Antarctic troposphere measured during the photochemical experiment at Neumayer (PEAN'99), *Atmospheric Environment*, 34, 5235-5247, 2000.
- Jardine, P. E., Fraser, W. T., Gosling, W. D., Roberts, C. N., Eastwood, W. J., and Lomax, B. H.: Proxy reconstruction of ultraviolet-B irradiance at the Earth's surface, and its relationship with solar activity and ozone thickness, *The Holocene*, 30, 155-161, 2020.
- 595 Jones, A., Weller, R., Minikin, A., Wolff, E., Sturges, W., McIntyre, H., Leonard, S., Schrems, O., and Bauguitte, S.: Oxidized nitrogen chemistry and speciation in the Antarctic troposphere, *Journal of Geophysical Research: Atmospheres*, 104, 21355-21366, 1999.
- Jones, A. E., Wolff, E. W., Ames, D., Bauguitte, S.-B., Clemishaw, K., Fleming, Z., Mills, G., Saiz-Lopez, A., Salmon, R. A., and Sturges, W.: The multi-seasonal NO<sub>y</sub> budget in coastal Antarctica and its link with surface snow and ice core nitrate: results from the CHABLIS campaign, *Atmospheric Chemistry and Physics*, 11, 9271-9285, 2011.
- 600 Langway, C., Osada, K., Clausen, H., Hammer, C., and Shoji, H.: A 10-century comparison of prominent bipolar volcanic events in ice cores, *Journal of Geophysical Research: Atmospheres*, 100, 16241-16247, 1995.
- Lin, J., Svensson, A., Hvidberg, C. S., Lohmann, J., Kristiansen, S., Dahl-Jensen, D., Steffensen, J. P., Rasmussen, S. O., Cook, E., and Kjær, H. A.: Magnitude, frequency and climate forcing of global volcanism during the last glacial period as seen in Greenland and Antarctic ice cores (60–9 ka), *Climate of the Past*, 18, 485-506, 2022.
- 605 Lomax, B. H., Fraser, W. T., Sephton, M. A., Callaghan, T. V., Self, S., Harfoot, M., Pyle, J. A., Wellman, C. H., and Beerling, D. J.: Plant spore walls as a record of long-term changes in ultraviolet-B radiation, *Nature Geoscience*, 1, 592-596, 2008.
- Marshall, G. J.: Trends in the Southern Annular Mode from observations and reanalyses, *Journal of climate*, 16, 4134-4143, 2003.
- Marshall, G. J., Fogt, R. L., Turner, J., and Clem, K. R.: Can current reanalyses accurately portray changes in Southern Annular Mode structure prior to 1979?, *Climate Dynamics*, 59, 3717-3740, 2022.
- 610 Millero, F. J., Feistel, R., Wright, D. G., and McDougall, T. J.: The composition of Standard Seawater and the definition of the Reference-Composition Salinity Scale, *Deep Sea Research Part I: Oceanographic Research Papers*, 55, 50-72, 2008.
- Ming, A., Winton, V. H. L., Keeble, J., Abraham, N. L., Dalvi, M. C., Griffiths, P., Caillon, N., Jones, A. E., Mulvaney, R., Savarino, J., Frey, M., and Yang, X.: Stratospheric ozone changes from explosive tropical volcanoes: Modeling and ice core constraints, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD032290, 2020.
- 615 Moore, J. and Paren, J.: A new technique for dielectric logging of Antarctic ice cores, *Le Journal de Physique Colloques*, 48, C1-155-C151-160, 1987.
- Mulvaney, R. and Wolff, E. W.: Evidence for winter/spring denitrification of the stratosphere in the nitrate record of Antarctic firm cores, *Journal of Geophysical Research: Atmospheres*, 98, 5213-5220, 1993.
- Mulvaney, R., Bremner, S., Tait, A., and Audley, N.: A medium-depth ice core drill, 2002.
- 620 Nye, J.: Correction factor for accumulation measured by the thickness of the annual layers in an ice sheet, *Journal of Glaciology*, 4, 785-788, 1963.
- Oerter, H., Wilhelms, F., Jung-Rothenhäusler, F., Göktas, F., Miller, H., Graf, W., and Sommer, S.: Accumulation rates in Dronning Maud Land, Antarctica, as revealed by dielectric-profiling measurements of shallow firm cores, *Annals of Glaciology*, 30, 27-34, 2000.
- Plummer, C. T., Curran, M. A., van Ommen, T. D., Rasmussen, S. O., Moy, A. D., Vance, T. R., Clausen, H. B., Vinther, B. M., and Mayewski, P. A.: An independently dated 2000-yr volcanic record from Law Dome, East Antarctica, including a new perspective on the dating of the 1450s CE eruption of Kuwae, Vanuatu, *Climate of the Past*, 8, 1929-1940, 2012.
- 625 Rankin, A., Auld, V., and Wolff, E.: Frost flowers as a source of fractionated sea salt aerosol in the polar regions, *Geophysical Research Letters*, 27, 3469-3472, 2000.
- Rankin, A. M., Wolff, E. W., and Mulvaney, R.: A reinterpretation of sea-salt records in Greenland and Antarctic ice cores?, *Annals of Glaciology*, 39, 276-282, 2004.
- 630 Raphael, M.: A zonal wave 3 index for the Southern Hemisphere, *Geophysical Research Letters*, 31, 2004.



- Rayner, N., Parker, D. E., Horton, E., Folland, C. K., Alexander, L. V., Rowell, D., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, *Journal of Geophysical Research: Atmospheres*, 108, 2003.
- 635 Reijmer, C. and Oerlemans, J.: Temporal and spatial variability of the surface energy balance in Dronning Maud Land, East Antarctica, *Journal of Geophysical Research: Atmospheres*, 107, ACL 9-1-ACL 9-12, 2002.
- Reijmer, C., Van den Broeke, M., and Scheele, M.: Air parcel trajectories and snowfall related to five deep drilling locations in Antarctica based on the ERA-15 dataset, *Journal of climate*, 15, 1957-1968, 2002.
- Ruth, U., Barnola, J. M., Beer, J., Bigler, M., Blunier, T., Castellano, E., Fischer, H., Fundel, F., Huybrechts, P., Kaufmann, P., Kipfstuhl, S., Lambrecht, A., Morganti, A., Oerter, H., Parrenin, F., Rybak, O., Severi, M., Udisti, R., Wilhelms, F., and Wolff, E.: "EDML1": a  
640 chronology for the EPICA deep ice core from Dronning Maud Land, Antarctica, over the last 150 000 years, *Clim. Past*, 3, 475-484, 10.5194/cp-3-475-2007, 2007.
- Savarino, J., Kaiser, J., Morin, S., Sigman, D., and Thiemens, M.: Nitrogen and oxygen isotopic constraints on the origin of atmospheric nitrate in coastal Antarctica, *Atmospheric Chemistry and Physics*, 7, 1925-1945, 2007.
- Shi, G., Hu, Y., Ma, H., Jiang, S., Chen, Z., Hu, Z., An, C., Sun, B., and Hastings, M. G.: Snow nitrate isotopes in central Antarctica record  
645 the prolonged period of stratospheric ozone depletion from~ 1960 to 2000, *Geophysical Research Letters*, e2022GL098986, 2022a.
- Shi, G., Chai, J., Zhu, Z., Hu, Z., Chen, Z., Yu, J., Ma, T., Ma, H., An, C., and Jiang, S.: Isotope fractionation of nitrate during volatilization in snow: a field investigation in Antarctica, *Geophysical Research Letters*, 46, 3287-3297, 2019.
- Shi, G., Li, C., Li, Y., Chen, Z., Ding, M., Ma, H., Jiang, S., An, C., Guo, J., and Sun, B.: Isotopic constraints on sources, production, and  
650 phase partitioning for nitrate in the atmosphere and snowfall in coastal East Antarctica, *Earth and Planetary Science Letters*, 578, 117300, 2022b.
- Sigl, M., Toohey, M., McConnell, J. R., Cole-Dai, J., and Severi, M.: Volcanic stratospheric sulfur injections and aerosol optical depth during the Holocene (past 11 500 years) from a bipolar ice-core array, *Earth System Science Data*, 14, 3167-3196, 2022.
- Sigl, M., McConnell, J. R., Toohey, M., Curran, M., Das, S. B., Edwards, R., Isaksson, E., Kawamura, K., Kipfstuhl, S., and Krüger, K.:  
655 Insights from Antarctica on volcanic forcing during the Common Era, *Nature Climate Change*, 4, 693-697, 2014.
- Sommer, S., Appenzeller, C., Röthlisberger, R., Hutterli, M. A., Stauffer, B., Wagenbach, D., Oerter, H., Wilhelms, F., Miller, H., and Mulvaney, R.: Glacio-chemical study spanning the past 2 kyr on three ice cores from Dronning Maud Land, Antarctica: 1. Annually resolved accumulation rates, *Journal of Geophysical Research: Atmospheres*, 105, 29411-29421, 10.1029/2000jd900449, 2000.
- Thomas, E. R., Allen, C. S., Etourneau, J., King, A. C., Severi, M., Winton, V. H. L., Mueller, J., Crosta, X., and Peck, V. L.: Antarctic sea ice proxies from marine and ice core archives suitable for reconstructing sea ice over the past 2000 years, *Geosciences*, 9, 506, 2019.
- 660 Thomas, E. R., Vladimirova, D. O., Tetzner, D. R., Emanuelsson, B. D., Chellman, N., Dixon, D. A., Goose, H., Grieman, M. M., King, A. C. F., Sigl, M., Udy, D. G., Vance, T. R., Winski, D. A., Winton, V. H. L., Bertler, N. A. N., Hori, A., Laluraj, C. M., McConnell, J. R., Motizuki, Y., Takahashi, K., Motoyama, H., Nakai, Y., Schwanck, F., Simões, J. C., Lindau, F. G. L., Severi, M., Traversi, R., Wauthy, S., Xiao, C., Yang, J., Mosely-Thompson, E., Khodzher, T. V., Golobokova, L. P., and Ekaykin, A. A.: Ice core chemistry database: an Antarctic compilation of sodium and sulfate records spanning the past 2000 years, *Earth Syst. Sci. Data*, 15, 2517-2532, 10.5194/essd-15-2517-2023,  
665 2023.
- Titchner, H. A. and Rayner, N. A.: The Met Office Hadley Centre sea ice and sea surface temperature data set, version 2: 1. Sea ice concentrations, *Journal of Geophysical Research: Atmospheres*, 119, 2864-2889, 2014.
- Traufetter, F., Oerter, H., Fischer, H., Weller, R., and Miller, H.: Spatio-temporal variability in volcanic sulphate deposition over the past 2 kyr in snow pits and firn cores from Amundsenisen, Antarctica, *Journal of Glaciology*, 50, 137-146, 2004.
- 670 Turner, J., Phillips, T., Thamban, M., Rahaman, W., Marshall, G. J., Wille, J. D., Favier, V., Winton, H., Thomas, E., and Wang, Z.: The Dominant Role of Extreme Precipitation Events in Antarctic Snowfall Variability, *Geophysical Research Letters*, 2019.
- Vega, C. P., Isaksson, E., Schlosser, E., Divine, D., Martma, T., Mulvaney, R., Eichler, A., and Schwikowski-Gigar, M.: Variability of sea salts in ice and firn cores from Fimbul Ice Shelf, Dronning Maud Land, Antarctica, *The Cryosphere*, 12, 1681-1697, 2018.
- Wagenbach, D., Ducroz, F., Mulvaney, R., Keck, L., Minikin, A., Legrand, M., Hall, J., and Wolff, E.: Sea-salt aerosol in coastal Antarctic regions, *Journal of Geophysical Research: Atmospheres*, 103, 10961-10974, 1998.
- 675 Welker, C., Martius, O., Froidevaux, P., Reijmer, C. H., and Fischer, H.: A climatological analysis of high-precipitation events in Dronning Maud Land, Antarctica, and associated large-scale atmospheric conditions, *Journal of Geophysical Research: Atmospheres*, 119, 11,932-911,954, 2014.
- Weller, R. and Wagenbach, D.: Year-round chemical aerosol records in continental Antarctica obtained by automatic samplings, *Tellus B: Chemical and Physical Meteorology*, 59, 755-765, 2007.
- 680 Weller, R., Jones, A., Wille, A., Jacobi, H. W., McIntyre, H., Sturges, W., Huke, M., and Wagenbach, D.: Seasonality of reactive nitrogen oxides (NO<sub>y</sub>) at Neumayer Station, Antarctica, *Journal of Geophysical Research: Atmospheres*, 107, ACH 2-1-ACH 2-11, 2002.
- Wilhelms, F., Kipfstuhl, J., Miller, H., Heinloth, K., and Firestone, J.: Precise dielectric profiling of ice cores: a new device with improved guarding and its theory, *Journal of Glaciology*, 44, 171-174, 1998.
- 685 Wilhelms, F., Miller, H., Gerasimoff, M. D., Drücker, C., Frenzel, A., Fritzsche, D., Grobe, H., Hansen, S. B., Hilmarsson, S. Æ., Hoffmann, G., Hörnby, K., Jaeschke, A., Jakobsdóttir, S. S., Juckschat, P., Karsten, A., Karsten, L., Kaufmann, P. R., Karlin, T., Kohlberg, E., Kleffel,




- G., Lambrecht, A., Lambrecht, A., Lawer, G., Schärmeli, I., Schmitt, J., Sheldon, S. G., Takata, M., Trenke, M., Twarloh, B., Valero-Delgado, F., and Wilhelms-Dick, D.: The EPICA Dronning Maud Land deep drilling operation, *Annals of Glaciology*, 55, 355-366, 10.3189/2014AoG68A189, 2017.
- 690 Winton, V. H. L., Ming, A., Caillon, N., Hauge, L., Jones, A. E., Savarino, J., Yang, X., and Frey, M. M.: Deposition, recycling, and archival of nitrate stable isotopes between the air–snow interface: comparison between Dronning Maud Land and Dome C, Antarctica, *Atmospheric Chemistry and Physics*, 20, 5861-5885, 2020.
- Winton, V. H. L. W., Caillon, N., Hauge, L., Mulvaney, R., Rix, J., Tuckwell, R., Savarino, J., and Frey, M.: Ice core chemistry, density, conductivity, dust, snow accumulation rate, and stable nitrate isotopic composition of the 120 m ISOL-ICE ice core, Dronning Maud Land, Antarctica (Version 1.0) [Data set] [dataset], 10.5285/9c972cfb-0ffa-4144-a943-da6eb82431d2, 2019.
- 695 Yang, J., Xiao, C., Liu, J., Li, S., and Qin, D.: Variability of Antarctic sea ice extent over the past 200 years, *Science Bulletin*, 2021.
- Yang, X., Frey, M. M., Rhodes, R. H., Norris, S. J., Brooks, I. M., Anderson, P. S., Nishimura, K., Jones, A. E., and Wolff, E. W.: Sea salt aerosol production via sublimating wind-blown saline snow particles over sea ice: parameterizations and relevant microphysical mechanisms, *Atmospheric Chemistry and Physics*, 19, 8407-8424, 2019.
- 700 Zielinski, G. A., Mayewski, P. A., Meeker, L. D., Whitlow, S., Twickler, M. S., Morrison, M., Meese, D. A., Gow, A. J., and Alley, R. B.: Record of volcanism since 7000 BC from the GISP2 Greenland ice core and implications for the volcano-climate system, *Science*, 264, 948-952, 1994.






## Appendices




705

**Figure A1.** Cross section of the ISOL-ICE ice core cut plan.



710 **Figure A2.** Age depth model of the ISOL-ICE ice core and comparison to the EDML (EDML1 chronology (Ruth et al., 2007)) and M1 ice core (Hofstede et al., 2004). EDML and ISOL-ICE ice cores have the same age-depth scale as the drilling sites are located 1 km apart. Whereas, M1 was drilled in a slightly higher accumulation zone, and thus captures fewer years for the same depth.



**Figure A3.** Seasonal cycles in (a) methanesulfonic acid (MSA), (b) non sea salt-sulfate (nss-SO<sub>4</sub><sup>2-</sup>), (c) SO<sub>4</sub><sup>2-</sup>/Na<sup>+</sup> ratio, and (d) sodium (Na) concentrations in ISOL-ICE snow pit A (Winton et al., 2020).



715 **Table A1. Accuracy and precision of soluble anion and trace element concentrations determined from repeated measurements of two CRM (European Reference Material ERM-CA408 simulated rainwater and European Reference Material, ERM-CA616 groundwater). Accuracy and precision are not reported for MSA as the CRM does not contain that analyte. The number of CRM measurements used for the accuracy and precision calculations are also listed (n).**

Analyte	n	Accuracy (%)	Precision (%)
Na	20	94	4
Mg	20	113	4
NO <sub>3</sub> <sup>-</sup>	20	107	10
SO <sub>4</sub> <sup>2-</sup>	20	96	10



**Table A2.** Tie points for annual layer counting. No data: n.d.

<b>Tie points</b>	<b>Date (AD)</b>	<b>Depth (m)</b>		
Drilling start	2017	0		
<b>Eruption</b>	<b>Eruption date (AD)</b>	<b>Arrival date (AD)</b>	<b>Depth (m)</b>	<b>Reference</b>
Pinatubo	1991	n.d.	4.91	(Cole-Dai and Mosley-Thompson, 1999)
Krakatoa	1883	1884	19.75	(Zielinski et al., 1994)
Tambora	1815	1815	27.38	(Zielinski et al., 1994)
"1809"	1809	1810	28.02	(Zielinski et al., 1994)
Komaga-Take	1694	n.d.	39.68	(Zielinski et al., 1994)
Huaynaputina	1600	1602	48.13	(Zielinski et al., 1994)
Kuwaie	1450	1458	61.01	(Zielinski et al., 1994; Plummer et al., 2012)
"1285"	1285	1285	75.12	(Langway et al., 1995)
"1277"	1277	1277	75.77	(Langway et al., 1995)
"1269"	1269	1269	76.41	(Langway et al., 1995)
"1259"	1259	1259	77.12	(Langway et al., 1995)
"1228/30"	1229	n.d.	79.33	(Langway et al., 1995)
"1166"	1166	1168	83.99	(Zielinski et al., 1994; Langway et al., 1995)
"1108"	1108	n.d.	88.15	(Traufetter et al., 2004)
"1050"	1050	1040	91.28	(Traufetter et al., 2004)
"975"	975	n.d.	98.45	(Traufetter et al., 2004)
"961"	961	n.d.	99.52	(Traufetter et al., 2004)
"719"	719	n.d.	116.40	(Traufetter et al., 2004)
"685"	685	n.d.	118.90	(Traufetter et al., 2004)



**Table A3. Annual correlations of ice core chemistry and snow accumulation rate with the SAM, Niño 3.4, and SOI indices over the period 1957-2016 and 1979-2016. Correlations significant at  $p < 0.10$  are boldface. The significance is calculated based on a two-tailed Students t-test.**

	Na <sup>+</sup>	Mg <sup>2+</sup>	MSA	Cl <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>	Accumulation
1957-2016						
SAM	<b>-0.32</b>	-0.2	-0.08	-0.17	-0.02	0.27
Niño 3.4	-0.03	0.05	0.05	0.12	0.12	0.19
SOI	-0.02	-0.07	-0.02	-0.1	<b>-0.25</b>	-0.07
1979-2016						
SAM	-0.14	0.08	0.03	-0.16	-0.08	0.21
Niño 3.4	0	0.15	0.21	0.07	<b>0.36</b>	0.15
SOI	-0.15	-0.24	-0.22	-0.13	<b>-0.55</b>	-0.01