

1 **Contrasting the Penultimate and Last Glacial Maxima (140 and 21 ka** 2 **BP) using coupled climate-ice sheet modelling**

3 Violet L. Patterson¹, Lauren J. Gregoire¹, Ruza F. Ivanovic¹, Niall Gandy², Jonathan Owen¹, Robin S.
4 Smith³, Oliver G. Pollard¹, Lachlan C. Astfalck⁴, Paul J. Valdes⁵

5 ¹School of Earth and Environment, University of Leeds, Leeds, UK

6 ²Department of the Natural and Built Environment, Sheffield Hallam University, Sheffield, UK

7 ³NCAS, Department of Meteorology, University of Reading, Reading, UK

8 ⁴School of Physics, Mathematics and Computing, University of Western Australia, Perth, Australia

9 ⁵School of Geographical Sciences, University of Bristol, Bristol, UK

10 *Correspondence to:* Violet L. Patterson (ee17vp@leeds.ac.uk)

11 **Abstract.** The configuration of the Northern Hemisphere ice sheets during the Penultimate Glacial Maximum differed to the
12 Last Glacial Maximum. However, the reasons for this are not yet fully understood. These differences likely contributed to the
13 varied deglaciation pathways experienced following the glacial maxima and may have had consequences for the interglacial
14 sea level rise. To understand the differences between the North American Ice Sheet at the Last and Penultimate Glacial Maxima
15 (21 and 140 ka BP), we perform two perturbed-physics ensembles of 62 simulations using a coupled atmosphere-ice sheet
16 model, FAMOUS-ice, with prescribed surface ocean conditions, in which the North American and Greenland ice sheets are
17 dynamically simulated with the Glimmer ice sheet model. We apply an implausibility metric to find ensemble members that
18 match reconstructed ice extent and volumes at the Last and Penultimate Glacial Maxima. We use a resulting set of ‘plausible’
19 parameters to perform sensitivity experiments to decompose the role of climate forcings (orbit, greenhouse gases) and initial
20 conditions on the final ice sheet configurations. This confirms that the initial ice sheet conditions used in the model are
21 extremely important in determining the difference in final ice volumes between both periods due to the large effect of the ice-
22 albedo feedback. In contrast to evidence of a smaller Penultimate North American Ice Sheet, our results show that the climate
23 boundary conditions at these glacial maxima, if considered in isolation, imply a larger Penultimate Glacial Maximum North
24 American Ice Sheet than at the Last Glacial Maximum, of around 6 meters sea level equivalent. This supports the notion that
25 the growth of the ice sheet prior to the glacial maxima is key in explaining the differences in North American ice volume.

26 **1 Introduction**

27 The Penultimate Glacial Maximum (PGM) occurred around 140,000 years ago, within Marine Isotope Stage 6 (MIS 6).
28 Greenhouse gas (GHG) concentrations and global average insolation were similar to the Last Glacial Maximum (LGM; ~21
29 ka BP) (Berger and Loutre, 1991; Loulergue et al., 2008; Bereiter et al., 2015) but the orbital configuration differed, affecting
30 the seasonal and latitudinal distribution of incoming shortwave radiation (Berger, 1978; Colleoni et al., 2011). The global total
31 ice sheet volume, and thus the global mean sea level, was likely similar between the two glacial maxima (~120-130 m below

32 present), with larger uncertainty at the PGM (Rabineau et al., 2006; Masson-Delmotte et al., 2010; Rohling et al., 2017). Both
33 geological evidence and numerical modelling suggest that despite the similarities in total ice volume between the PGM and
34 the LGM, the configurations of the Northern Hemisphere ice sheets differed significantly (e.g. Svendsen et al., 2004; Colleoni
35 et al., 2016; Batchelor et al., 2019).

36 Some reconstructions suggest the Eurasian Ice Sheet (EIS) may have been up to ~50 % larger during the Penultimate Glacial
37 Cycle (MIS 6: ~190-130 ka BP) than during the Last Glacial Cycle (~115-12 ka BP) (Svendsen et al., 2004). However,
38 evidence of multiple advances and uncertainties in dating proxy records means that the maximum extent mapped at 140 ka BP
39 could correspond to previous advances during MIS 6. Similarly, the timing of the maximum extent of the EIS at the LGM is
40 also uncertain and areas of the ice margin likely reached their maximum extents at different times throughout the glacial cycle
41 (Svendsen et al., 2004; Margari et al., 2014; Colleoni et al., 2016; Ehlers et al., 2018). The extent of the North American Ice
42 Sheet (NAIS) during the PGM is even less well constrained due to a lack of glaciological evidence (e.g. moraines and till).
43 The scarcity of empirical data in itself suggests that it was smaller in most areas than at the LGM because the subsequent larger
44 ice sheet could have largely erased the evidence of prior glaciations (Dyke et al., 2002; Rohling et al., 2017). Additionally,
45 evidence of reduced ice rafted debris (IRD) discharge from the Hudson Strait in the North Atlantic IRD belt (e.g. Hemming,
46 2004; Naafs et al., 2013; Obrochta et al., 2014), relative sea level assessment studies (e.g. Rohling et al., 2017) and climate,
47 ice sheet and glacial isostatic adjustment modelling (e.g. Colleoni et al., 2016; Dyer et al., 2021) all point to a smaller volume
48 PGM NAIS. For example, assuming a similar global mean sea level fall (and Antarctic ice sheet volume) at the PGM as at the
49 LGM but with a larger volume EIS at the PGM (estimated at 33-53 m sea level equivalent (SLE) versus 14-29 m SLE at the
50 LGM), this follows that the NAIS must have been smaller than at the LGM to compensate (39-59 m SLE versus 51-88 m SLE)
51 (Rohling et al., 2017).

52 The reason for these differences is likely complex and is not yet fully understood. The evolution and surface mass balance
53 (SMB) of ice sheets depends on many factors such as; background climate, climate and ice sheet histories, dust deposition,
54 vegetation, ice albedo and sea surface temperatures, as well as the interactions and feedbacks between them all (Kageyama et
55 al., 2004; Krinner et al., 2006; 2011; Colleoni et al., 2009a; 2011; Liakka et al., 2012; Stone and Lunt, 2013). The ice sheets
56 themselves also strongly influence the climate through their interactions with atmospheric and oceanic circulation and the
57 energy balance. This alters global and local temperature and precipitation patterns which in turn affects ice sheet ablation and
58 accumulation (i.e. SMB) (e.g. Kageyama and Valdes, 2000; Abe-Ouchi et al., 2007; Beghin et al., 2014; 2015; Ullman et al.,
59 2014; Liakka et al., 2016; Gregoire et al., 2015; 2018; Snoll et al., 2022; Izumi et al., 2023). These interactions between the
60 vast ice sheets and other components of the climate system exerted an important control on the initial climate state for the
61 deglaciations, and hence on the subsequent chain of events, thus impacting the climate, ocean and sea level evolution during
62 deglaciation. Thus, the contrasting configurations of the Northern Hemisphere ice sheets at the glacial maxima may have
63 contributed to the different deglaciation pathways that followed. In this context, it is important to examine the complex physical
64 interactions between the climate and the ice sheets to better understand why the last two glacial maxima had different ice sheet

65 configurations and evaluate the ice sheets' sensitivities to changes in climate in relation to different orbits and greenhouse gas
66 concentrations. To achieve this, numerical simulations of these periods are required using a coupled climate-ice sheet model
67 that capture these complex, non-linear interactions. Previous studies on glacial-interglacial cycles, have relied on the coupling
68 of relatively fast, low resolution and simplified Earth system Models of Intermediate Complexity (EMICs) to an ISM (e.g.
69 Bonelli et al., 2009; Ganopolski et al., 2010; Fyke et al., 2011; Heinemann et al., 2014; Beghin et al., 2014; Ganopolski and
70 Brovkin, 2017; Quiquet et al., 2021; Poppelmeier et al., 2023; Willeit et al., 2024) or one-way forcing of an ice sheet model
71 with climate forcing output by stand-alone climate simulations (e.g. Abe-Ouchi et al., 2013; Stone and Lunt, 2013; Gregoire
72 et al., 2015; 2016). These computationally efficient techniques advanced our understanding of the roles of orbit and CO₂ in ice
73 sheet evolution and proposed plausible reconstructions of past ice sheets (e.g. Robinson et al., 2011; Stone et al, 2013). They
74 also highlighted important earth system interactions (e.g. Stone and Lunt, 2013; Willeit et al., 2024) such as with vegetation,
75 dust, albedo, glacial isostatic adjustment, disparate ice sheets (Beghin et al., 2015) as well as internal ice sheet instabilities
76 (Gregoire et al., 2012; Quiquet et al., 2021). However, the accuracy of these results has been limited by the simplified
77 representation of climate processes, atmospheric circulation and/or surface mass balance. A combination of increased computer
78 power, the development of more computationally efficient, lower resolution General Circulation Models (GCMs) and sub-grid
79 scale schemes translating ice sheet relevant atmospheric processes onto the higher resolution ice sheet grid, has made bi-
80 directional, coupled climate-ice sheet simulations over longer timescales, and in large ensembles, feasible (Vizcaino et al.,
81 2013; Ziemen et al., 2014; Sellevold et al., 2019; Smith et al., 2021). These coupled models have been used to simulate the
82 climate-ice sheet interactions during past glacial periods including; glacial inception (Gregory et al., 2012); the LGM and the
83 build up to it (Ziemen et al., 2014; Gandy et al., 2023; Sherriff-Tadano et al., 2023; Nui et al., 2024) and MIS 13 (Niu et al.,
84 2021).


85 To better understand the differences between the Penultimate and Last Glacial Maxima ice sheet configurations, we seek to
86 establish how the differences in climate forcings (such as orbit and greenhouse gases) between the two periods affected ice
87 sheet surface mass balance and in turn their geometry. To this end, this study uses a coupled climate-ice sheet model
88 (FAMOUS-ice; Smith et al., 2021), to perform ensembles of simulations of the PGM and LGM to explore input climate and
89 ice sheet parameter uncertainties and their effects on the North American ice sheet volume during each period. We identify
90 simulations that match volume and extent constraints and use these to perform a factorial decomposition of the effects of
91 climate forcing and initial conditions on ice volume difference between the two Glacial Maxima.

92 **2 Methods**

93 **2.1 Model description**

94 FAMOUS is a fast, low resolution AOGCM that is based on Hadley Centre coupled model HadCM3 and therefore retains all
95 the complex processes represented in an AOGCM but uses only half the spatial resolution and a longer time step. Since it

96 requires only 10 % of the computational costs of HadCM3, it has been successfully used for long transient palaeo simulations
97 (Smith and Gregory, 2012; Gregory et al., 2012; Gregoire et al., 2012; Roberts et al., 2014; Dentith et al., 2019) and large
98 ensembles for uncertainty quantification (Gregoire, 2010; Gandy et al., 2023). This study uses the atmospheric component,
99 which is a hydrostatic, primitive equation grid point model with a horizontal resolution of 7.5° longitude by 5° latitude with
100 11 vertical levels and a 1-hour time step (Williams et al., 2013). Land processes are modelled using the MOSES2.2 land surface
101 scheme (Essery et al., 2003), which uses a set of sub-gridscale tiles in each grid box to represent fractions of nine different
102 surface types, including land ice (Smith et al., 2021). Whilst this study prescribes sea surface temperatures and sea ice
103 concentrations, FAMOUS can also be run fully coupled with a dynamical ocean (e.g. Dentith et al., 2019).
104 FAMOUS now allows the direct two-way coupling to an ice sheet model in the configuration FAMOUS-ice (Smith et al.,
105 2021). Here, we use FAMOUS in combination with Glimmer to interactively simulate the North American and Greenland ice
106 sheets at 40km resolution. Glimmer is a fast running, 3D thermomechanical ice sheet model which uses the shallow ice
107 approximation. This allows it to model ice sheet evolution over long timescales as it is more computationally efficient, and
108 therefore has been used to simulate continental ice sheets over glacial-interglacial cycles (Rutt et al., 2009; Gregoire et al.,
109 2016).
110 FAMOUS-ice accounts for the mismatch between atmosphere and ice sheet grid sizes by using a multilayer surface snow
111 scheme to calculate SMB on ‘tiles’ at 10 set elevations within each grid box that contains land ice in FAMOUS. This SMB is
112 then downscaled from the coarse FAMOUS grid to the much finer Glimmer grid at each model year (Smith et al., 2021).
113 Glimmer uses this SMB field to calculate ice flow and surface elevation and passes this back to FAMOUS in which orography
114 and ice cover is updated. In this study, to reduce computational costs further, FAMOUS-ice runs at 10 times ice sheet
115 acceleration: for every year of climate integrated in FAMOUS, the simulated SMB field forces 10 years of ice sheet integration
116 in Glimmer. Figure 1 shows a simplified diagram of this coupling process and full details can be found in Smith et al.,
117 (2021). The current computational cost of this set up is around 50 decades (of climate years) per wallclock day using 8
118 processors (~ 192 core hours).
119 FAMOUS-ice has been shown to perform well in simulations of past and future ice sheets including Greenland and North
120 America (Gregory et al., 2020; Smith et al., 2021; Gandy et al., 2023). In particular, the LGM North American Ice Sheet study
121 of Gandy et al., (2023) was able to utilise the useful constraints of the LGM to infer the importance of parameters controlling
122 ice sheet albedo on ice sheet configuration in this model.



123
124 **Figure 1. Schematic illustrating the calculation of SMB along a specific transect across the ice sheet (blue line) at different elevations**
125 **on the FAMOUS grid followed by downscaling onto the Glimmer grid.**

126 **2.2 Experiment design**

127 **2.2.1 Climate boundary conditions**

128 With the exception of including dynamic North American and Greenland ice sheets, our FAMOUS-ice simulations are set up
129 following the Paleoclimate Modelling Intercomparison Project Phase 4 (PMIP4) protocols for the LGM (Kageyama et al.,
130 2017) and PGM (Menviel et al., 2019). These protocols prescribe climatic boundary conditions, including orbital parameters
131 and GHG concentrations, the values of which can be found in Table 1. Concentrations of CO_2 , CH_4 and N_2O are very similar
132 between the LGM and PGM but orbital parameters are significantly different. The larger eccentricity at the PGM enhances the
133 effect of precession compared to the LGM which affects the seasonal and latitudinal distribution of insolation. These changes
134 are important for ice sheet surface mass balance since melting is particularly sensitive to spring and summer temperatures

135 (Huybers, 2006; Niu et al., 2019). The PGM received lower insolation in the Northern Hemisphere in late winter to early
 136 summer but higher levels in late summer to early winter, compared to the LGM (Fig. 2a). Subsequent to the completion of this
 137 work, it was discovered that the equation for the role of eccentricity on solar insolation was incorrect in the model code. The
 138 magnitude of the error is larger for periods with higher eccentricity values and so a sensitivity test was run to determine the
 139 effect this correction has on SMB and ice volume at the PGM. Details of this error and the results of the sensitivity test can be
 140 found in Appendix A, but the impact was shown to be minimal (Fig. A1).

141

142 **Table 1. Climate boundary conditions used in the LGM and PGM experiments as prescribed by the PMIP4 protocols for each period**
 143 **(Kageyama et al., 2017; Menviel et al., 2019).**

	<i>Eccentricity</i>	<i>Obliquity</i> (°)	<i>Perihelion –</i> <i>180 (°)</i>	<i>Solar</i> <i>Constant</i> (Wm^{-2})	<i>CO₂</i> (<i>ppm</i>)	<i>CH₄</i> (<i>ppb</i>)	<i>N₂O</i> (<i>ppb</i>)	<i>Orography and</i> <i>ice extent</i>
<i>LGM</i> <i>(21 ka)</i>	0.019	22.949	114	1360.7	190	375	200	GLAC-1D (Tarasov et al., 2012; Briggs et al., 2014; Ivanovic et al., 2016)
<i>PGM</i> <i>(140 ka)</i>	0.033	23.414	73	1360.7	191	385	201	Combined reconstruction (Abe-Ouchi et al 2013; Briggs et al 2014; Tarasov et al 2012)






144

145 In the climate model, the global orography (including the Eurasian and Antarctic ice sheets) and land-sea mask for the LGM
 146 are calculated from the GLAC-1D 21 ka BP reconstruction (Tarasov et al., 2012; Briggs et al., 2014; Ivanovic et al., 2016),
 147 which is one of three recommendations in the PMIP4 protocol (Kageyama et al., 2017). For the PGM simulations we used the
 148 140 ka BP combined reconstruction (Tarasov et al., 2012; Abe-Ouchi et al., 2013; Briggs et al., 2014) detailed in the PGM
 149 PMIP4 protocol (Menviel et al., 2019). Vegetation is prescribed based on a pre-industrial distribution and kept constant. As
 150 ice cover changes, the fractions of grid cells that are land ice versus other surface types changes proportionally, altering albedo.
 151 However, since there is no dynamical vegetation component, some important climate-ice-vegetation feedbacks are neglected,
 152 which could have a significant impact on ice sheet evolution (Stone and Lunt, 2013).

153 Because of the low resolution of the FAMOUS model, using a dynamical ocean and sea ice can introduce large biases in the
154 simulated climate (Dentith et al. 2019). By prescribing Sea Surface Temperature (SST) and sea ice, we are able to limit the
155 amplification of climate biases arising from atmosphere-ocean-sea ice interactions. Thus, SSTs and sea ice concentration are
156 also prescribed and constant and are taken from higher resolution HadCM3 simulations of 21 ka BP (Fig. B1a; see details in
157 Izumi et al., 2022) and 140 ka BP (Fig. B1b). The 140 ka BP simulation is part of a suite of simulations covering the last
158 140,000 years (Allen et al., 2020). It was performed using a version of HadCM3 (specifically HadCM3B-M2.1aD, see Valdes
159 et al., (2017), which was the same version as used by Izumi et al., (2022) for the LGM and Davies-Barnard et al., (2017)). The
160 simulation was forced with 140 ka BP orbital configuration (Berger and Loutre, 1991) and greenhouse gases (Petit et al., 1999;
161 Spahni et al., 2005; Loulergue et al., 2008). Ice sheet forcing and land sea mask were from DeBoer et al., (2013) who modelled
162 the evolution of all the major ice sheets. It was run as a “snapshot” simulation for 3070 years which allowed the deeper ocean
163 to attain near equilibrium.

164 FAMOUS atmosphere-ocean GCM has not been run for the PGM, and we lack sufficient data density for precisely dated PGM
165 SSTs and sea-ice to produce statistically varied reconstructions, as in Gandy et al., (2023). Thus, for physical consistency
166 between the LGM and PGM periods, HadCM3 output was used for the surface ocean boundary conditions. Of all possible
167 options, HadCM3 output is the most appropriate choice for this because it is the parent model for FAMOUS; they share the
168 same physics, differing mainly in their resolutions, and HadCM3 was used as the tuning target for FAMOUS during model
169 development (Smith et al., 2008). We take the multi-year monthly mean “climatology” of SSTs and sea ice concentrations
170 from the final 100 years of the simulations. These 12-month climatologies are repeated throughout the duration of the
171 simulations to provide a seasonal forcing with no long-term trend and no interannual variability.

172 The modelled annual average SSTs are cooler at the LGM than at the PGM, everywhere, except in the North Atlantic due to
173 less sea ice cover in this region (Fig. 2b). However, the summer SSTs are warmer in the Northern Hemisphere at the LGM
174 compared to the PGM (Fig. 2c). The HadCM3 LGM SSTs are colder on average than the reconstruction in Gandy et al., (2023),
175 with the largest differences, of up to 6 °C, occurring in the tropics and mid-latitudes (Fig. B1c).



176

177

178

Figure 2. Difference between the LGM and PGM (a) latitudinal distribution of incoming top of the atmosphere shortwave radiation each month (b) modelled annual sea surface temperatures and (c) modelled summer (JJA) sea surface temperatures.

179

2.2.2 Ice sheet boundary and initial conditions

180


In all our simulations, the ice sheet extent is set to the PMIP4 boundary conditions for the LGM and PGM as described in Table 1, except in the interactive ice sheet model domain, which covers North America and Greenland. Here, we describe how the ice extent and elevation is initialised in FAMOUS and Glimmer over the interactive domain in our ensemble of PGM and LGM simulations and sensitivity experiments.

184

In our ensemble of LGM and PGM simulations, Glimmer is initiated from an 18.2 ka BP NAIS taken from a previous Last Deglaciation ensemble (Gregoire et al., 2016). This smaller intermediate (MIS 3-like) ice sheet was used in Gandy et al., (2023) as an approximate pre-glacial maximum extent from which to grow the ice sheet towards an equilibrium ice volume. For consistency, we used the same initial ice sheet conditions as in Gandy et al. (2023) when running our ensembles of LGM and PGM simulations. The coupling between the models passes this orography field from Glimmer to FAMOUS, updating the PMIP4 boundary condition that FAMOUS was initiated from. However, due to the technical formulation of the coupling, where entire gridboxes were initialised as covered in ice at all elevations in FAMOUS, the tiles in such gridboxes would not

190

191 subsequently update to reflect the existence of any non-glaciated fractions that might exist in the Glimmer state. This means
192 that when the initial conditions are radically different in FAMOUS and Glimmer (as in our ensemble of simulations), the
193 FAMOUS ice extent over the North American continent is not updated to match the Glimmer initial conditions. Thus, in our
194 ensemble of LGM simulations, the albedo remains high throughout the saddle region (the area between the Laurentide and
195 Cordilleran ice sheets) because the FAMOUS ice extent remains as large as the atmospheric model's initial conditions (i.e. the
196 GLAC-1D 21 ka BP reconstruction) for the duration of the simulations (Fig. 3). The different ice sheet configurations used in
197 FAMOUS and Glimmer in the ensembles, are outlined in our table of experiments, Table 2 (experiments 1 and 2). The impact
198 of this set-up compared to an ice sheet configuration matched in FAMOUS and Glimmer is explored in Sect. 3.2 and Appendix
199 C.
200



201 **Figure 3. Topography anomaly from present day used as the initial condition in FAMOUS and the ice masks (red lines) for (a) the**
202 **LGM and (b) the PGM.**
203

204 We perform two sets of sensitivity experiments to understand the relative impact of the initial ice sheet conditions and the
205 climate forcing on the resulting LGM and PGM NAIS volumes. The first set of experiments uses matching ice sheet
206 configurations in FAMOUS and Glimmer, set either to the LGM GLAC-1D reconstruction or to the end of one of our PGM
207 coupled simulations (Table 2; experiments 3 – 6). The second set uses the same initial ice sheet configurations as in the
208 ensemble, i.e. GLAC-1D and PMIP4 reconstructions in FAMOUS and the 18.2 ka ice sheet in Glimmer (Table 2; experiments
209 7 - 10). A full description of the initial conditions and methods used in these sensitivity experiments can be found in Sect. 2.5.
210

211 **Table 2.** Table of experiments performed in this study detailing the ‘climate forcing’ (orbital configuration, trace gases and global
 212 orography as outlined in Table 1 and SSTs/sea ice from HadCM3), initial ice extent set in FAMOUS over Greenland and North
 213 America, initial Glimmer ice sheet conditions and input parameter values. NROY are the simulations that are ‘Not Ruled Out Yet’
 214 after applying the implausibility metric described in Sect. 2.4.

<i>Experiments</i>	<i>Climate forcing</i>	<i>FAMOUS initial ice extent</i>	<i>Glimmer initial condition</i>	<i>Input parameter values</i>
1) LGM ensemble	LGM	PMIP4 LGM (GLAC-1D)	18.2 ka ice sheet	Randomly sampled from Table 3 ranges (See Sect. 2.3)
2) PGM ensemble	PGM	PMIP4 PGM	18.2 ka ice sheet	Randomly sampled from Table 3 ranges (See Sect. 2.3)
3) V_{-1} (full LGM)	LGM	PMIP4 LGM (GLAC-1D)	PMIP4 LGM GLAC-1D	Matching NROYa simulation xpken/xpkyn (See Sect. 2.4 and 3.1)
4) V_{c-1}	PGM	PMIP4 LGM (GLAC-1D)	PMIP4 LGM GLAC-1D	
5) V_{i-1}	LGM	PGM NROYa (xpkyn)	PGM NROYa (xpkyn)	
6) V_{ci-1} (full PGM)	PGM	PGM NROYa (xpkyn)	PGM NROYa (xpkyn)	
7) V_{-2} (NROYa LGM)	LGM	PMIP4 LGM (GLAC-1D)	18.2 ka ice sheet	
8) V_{c-2}	PGM	PMIP4 LGM (GLAC-1D)	18.2 ka ice sheet	
9) V_{i-2}	LGM	PMIP4 PGM	18.2 ka ice sheet	
10) V_{ci-2} (NROYa PGM)	PGM	PMIP4 PGM	18.2 ka ice sheet	

215

216 **2.3 Ensemble design**

217 The ensemble by Gandy et al., (2023) showed that uncertainty in parameters controlling SMB, ice sheet dynamics and climatic
 218 conditions over the ice sheets had a significant influence on the extent and volume of the LGM NAIS, with albedo parameters
 219 explaining the majority of the variation in model output. Since these parameters needed re-tuning from simulations of the
 220 present day Greenland ice sheet to produce an acceptable LGM NAIS configuration in FAMOUS-ice under LGM climate
 221 conditions, the PGM may also show different sensitivities to the uncertain parameters. Therefore, we ran new ensembles of
 222 the LGM and PGM in order to explore uncertainties and identify combinations of climate and ice sheet parameters that perform
 223 well for both periods.

224 Following on from Gandy et al., (2023), a second wave of simulations was performed and compared to reconstructions of ice
 225 sheet extent and volume to identify ‘Not Ruled Out Yet’ (NROY) parameter combinations (see methodology in Appendix D),
 226 the results of which formed the basis of the ensemble design in this study. We reran the LGM ensemble to allow for slight
 227 changes in the experiment design compared to Gandy et al., (2023): we use orbital parameters for 21 ka BP rather than 23 ka
 228 BP and HadCM3 SSTs instead of a statistical reconstruction (see Sect. 2.2.1). Table 3 details the 13 parameters that were

229 varied in these simulations. Out of the 176 NROY parameter combinations from the Wave 2, a representative subset of 62
 230 were selected which provided adequate coverage of the NROY space (see Appendix D for details). Each was run for 1000
 231 climate years (10,000 ice sheet years) for both the LGM and PGM experiments until the majority of the ice sheet reached close
 232 to equilibrium. Despite differences in the model set up between this study and Gandy et al., (2023), we expect the 62 samples
 233 chosen from their design to be a good estimate to an optimal parameter design for our experiment design (Appendix D).
 234

235 **Table 3. Description of parameters varied in the ensembles. Adapted from Gandy et al., (2023).**


<i>Parameter</i>	<i>Range</i>	<i>Description</i>
<i>Lapse Rate</i>	-0.01 – -0.002 K km ⁻¹	Prescribed lapse rate for air temperature used to downscale FAMOUS near-surface ice sheet climate onto surface elevation tiles. Down welling longwave radiation is also adjusted for consistency. More negative values lead to stronger lapse rate effects (Smith et al., 2021).
<i>Daice</i>	-0.4 - 0 K ⁻¹	Sensitivity of bare-ice albedo to surface air temperatures once the surface is in a melt regime. Albedo reduced to as low as 0.15 with minimum value (Smith et al., 2021)
<i>Fsnow</i>	350 – 800 kg m ⁻³	The threshold in surface snow density at which the FAMOUS albedo scheme switches from a scattering paradigm appropriate for a conglomeration of snow grains to one more appropriate for a solid surface. Higher values correspond to using brighter albedos for denser snow, increasing ice sheet albedo (Smith et al., 2021)
<i>AV_GR</i>	0 – 0.01 μ m ⁻¹	Sensitivity of the snow albedo to variation in surface grain size. Higher values enhance the darkening of snow over time, decreasing the albedo (Smith et al., 2021).
<i>RHcrit</i>	0.6 - 0.9 Pa ⁻¹	The threshold of relative humidity for cloud formation (R. Smith, 1990). A higher value means clouds can form less easily.
<i>VFI</i>	1 – 2 m s ⁻¹	The precipitating ice fall-out speed (Heymsfield, 1977).
<i>CT</i>	5x10 ⁻⁵ - 4x10 ⁻⁴ s ⁻¹	The conversion rate of cloud liquid water droplets to precipitation (R. Smith, 1990).
<i>CW</i>	1x10 ⁻⁴ – 2x10 ⁻³ kg m ⁻³	The threshold values of cloud liquid water for formation of precipitation (R. Smith, 1990). Only the value for the land is varied.
<i>Entrainment Coefficient</i>	1.5 - 6	Rate of mixing between environmental air and convective plume. Higher values enhance mixing of convective plumes with ambient dry air.
<i>Alpham</i>	0.2 - 0.65	The sea ice lowest albedo (Crossley & Roberts, 1995).
<i>Basal Sliding</i>	0.5 - 20 mm yr -1	The basal sliding rate. A higher value allows increased ice velocity.

<i>Mantle Relaxation Time</i>	300 – 9000 yrs	The relaxation time of the mantle, a lower value making the mantle less viscous, thus allowing a quicker topographic rebound.
<i>Flow Enhancement Factor</i>	1 - 10	Glen’s Flow Law enhancement factor. Increasing the factor makes the ice softer and more deformable (Rutt et al., 2009).

236

237 **2.4 Implausibility criteria**

238 To filter out implausible ice sheet configurations in the results, a set of constraints, based on southern ice sheet extent and
239 volume, were applied to the LGM ensemble. Both ensembles were filtered based on the LGM results since the extent of the
240 NAIS is very well constrained by geological data and there are more estimates of ice volume for the LGM than the PGM. This
241 is because there is a lack of empirical data (over both space and time) on ice sheet configuration at the PGM due to destruction
242 of evidence by subsequent glaciations and difficulties with dating what is available (Parker et al., 2022). Thus, most of the
243 reconstructions of NAIS PGM extent are actually the maximum extent reached over the whole of MIS 6 (190-132 ka BP) and
244 are mostly based on numerical modelling combined with this scarce proxy data (e.g. Colleoni et al., 2016; Batchelor et al.,
245 2019). This leaves a set of plausible or ‘Not Ruled Out Yet’ (NROY) LGM simulations that can then be compared to the
246 corresponding PGM simulations to determine whether parameters that performed well for the LGM also give plausible PGM
247 results. LGM ice extent was assessed against the reconstruction by Dalton et al. (2020). We focus our evaluation of ice extent
248 on the southern NAIS area and chose to disregard regions of known model bias. This includes marine margins that are subject
249 to processes not included in Glimmer and the Alaskan regions where small climate model biases lead to ice sheet overgrowth
250 (e.g. Ganopolski et al., 2010; Ziemen et al., 2014; Gregoire et al. 2016, Sherriff-Tadano et al., 2023). Additionally, ice lobes
251 are not well captured in many models as they are likely to be transient, short-lived features that may be caused by complex ice
252 dynamics (e.g. Zweck and Huybrechts, 2005). Therefore, we do not expect our simulations to perfectly match the reconstructed
253 Southern NAIS extent. To account for the expected mismatch between model and data, we applied a tolerance on the Southern
254 ice sheet area of $1.79 \times 10^6 \text{ km}^2$, equivalent to three-times the area of the lobes (Fig. 4). We thus calculate the Southern NAIS
255 ice area as the integrated area within the large box shown in Fig. 4 at the end of each LGM simulation and selected simulations
256 that matched the reconstructed area from Dalton et al. (2020) within plus or minus $1.79 \times 10^6 \text{ km}^2$. The volume of the NAIS is
257 not as well constrained by proxy data and so estimates rely on ice sheet, glacial isostatic adjustment and sea level modelling
258 studies. Based on a number of these studies (Marshall et al., 2002; Tarasov and Peltier, 2002; 2004; Tarasov et al., 2012;
259 Lambeck et al., 2014; Peltier et al., 2015; Rohling et al., 2017; Batchelor et al., 2019; Gowan et al., 2021), a minimum NAIS
260 (including Greenland) volume of 70 m SLE ($2.8 \times 10^7 \text{ km}^3$) was applied to the ensemble. The translation of ice volumes into
261 meters of sea level equivalent are calculated based on present day ocean area.



262

263

264

265

Figure 4. Outline of the LGM North American Ice Sheet by Dalton et al. (2020). The large red box shows the region used to calculate reconstructed and modelled Southern NAIS area. The small red box shows the region used to calculate the area of the lobes from which we set the upper and lower target bounds for southern ice extent (See Sect. 2.4).

266

2.5 Sensitivity analysis

267

268

269

270

271

272

273

We choose one of the resulting NROY parameter combinations, NROYa (specifically experiments xpken/xpkyn), which has LGM and PGM ice volumes lying in the middle of estimated ranges and the least excess ice growth over Alaska, to investigate the relative impact of the initial conditions versus the climate on the resulting ice sheet configurations. This is achieved through a sensitivity analysis along with factorisation based on the method used in Lunt et al., (2012) and Gregoire et al. (2015). We divided the differences in inputs between LGM and PGM into two factors; the initial ice sheet configurations used in FAMOUS and Glimmer and the climate boundary conditions (orbital parameters, greenhouse gases and SSTs/sea ice). Thus, the total difference in final ice volume (ΔV) between the LGM and the PGM can be written as Eq. (1):

274

$$\Delta V = dV_{ice} + dV_{climate} , \quad (1)$$

275

276

where dV_{ice} is the difference in final ice volume due to the different initial ice sheet configurations and $dV_{climate}$ is the difference due to the difference climate boundary conditions used.

277

278

279

280

The factorisation method requires 2^N simulations (where N is the number of different components) to determine the contribution of each component to ice volume difference, therefore $2^2 = 4$ experiments are needed that systematically change one variable. These experiments are listed in Table 2. The relative contributions of the initial conditions and climate can be calculated by Eqs. (2) and (3):

$$281 \quad dV_{ice} = \frac{1}{2}((V_i - V) + (V_{ci} - V_c)), \quad (2)$$

$$282 \quad dV_{climate} = \frac{1}{2}((V_c - V) + (V_{ci} - V_i)), \quad (3)$$


283 To properly understand the effect of the initial conditions, we performed two sets of sensitivity experiments. In the first set,
 284 labelled $V_{_1}$, $V_{c_{_1}}$, $V_{i_{_1}}$ and $V_{ci_{_1}}$ (Table 2; experiments 3 – 6), both the topography and ice cover are set to be consistent between
 285 the climate and ice sheet model components. Specifically, for the LGM, the Glimmer initial bedrock topography and ice surface
 286 elevation was prescribed from the GLAC-1D reconstruction used in the FAMOUS LGM boundary condition. For the PGM,
 287 the ice thickness data needed for the PMIP4 reconstruction to be converted to the Glimmer initial condition were not available.
 288 Instead, both Glimmer and FAMOUS were initialised with the final timestep of the NROYa PGM (xpkyn) experiment since
 289 it closely resembles the PMIP4 reconstruction. Experiment $V_{_1}$ corresponds to a full LGM simulation and $V_{ci_{_1}}$ corresponds to
 290 a full PGM simulation. In the second set of sensitivity experiments, we use the initial Glimmer ice sheet used in the ensembles,
 291 i.e. the 18.2 ka mid-size ice sheet, only varying the FAMOUS initial ice sheets to see how this difference in orography between
 292 the climate and ice sheet models may have impacted the result. These experiments are labelled $V_{_2}$, $V_{c_{_2}}$, $V_{i_{_2}}$ and $V_{ci_{_2}}$ (Table
 293 2; experiments 7 – 10), with $V_{_2}$ corresponding to the LGM NROYa (xpken) and $V_{ci_{_2}}$ corresponding to the PGM NROYa
 294 (xpkyn).

295 **3 Results**

296 **3.1 Ensembles**

297 Our ensembles of 62 North American Ice Sheet configurations spans uncertainty in model parameters and reveals the wide
 298 range of possible modelled ice sheet evolutions. Over the full ensembles, we find that the set-up of the original Wave 2 meant
 299 that the albedo values were too high and so the use of more realistic albedos in these ensembles led to many of the runs
 300 deglaciating to very low volumes as shown in Fig. 5 (see Appendix D for more detail).

301



302

303 **Figure 5.** (a) Ice volume evolution over modelled time, and (b) density distribution of final ice volumes for the full LGM and PGM
 304 ensembles. Percentage of simulations with ice cover for (c) LGM (with the Dalton et al., (2020) reconstructed margin shown in red);
 305 (d) PGM (with the PMIP4 PGM modelled margin shown in solid red and the Batchelor et al., (2019) reconstructed maximum MIS
 306 6 margin shown in dashed red), and (e) the difference between the LGM and PGM, at the end of the simulations.


307 **Table 4.** Average volumes (NAIS + Greenland) and southern NAIS areas and their standard deviations (SD) of the NROY LGM and
 308 PGM simulations. Also shown are estimated values from literature for comparison.

309

	<i>Mean Total Volume (SD), m SLE</i>	<i>Estimated Total Volume, m SLE</i>	<i>Mean Southern Area (SD), x 10⁶ km²</i>	<i>Estimated Southern Area, x 10⁶ km²</i>
LGM	82.1 (8.29)	61-98 (Rohling et al., 2017)	5.55 (0.33)	6.28 (Dalton et al., 2020)
PGM	62.3 (10.3)	49-69 (Rohling et al., 2017)	3.64 (0.82)	3.32 (Menviel et al., 2019)

310


311 After applying our implausibility criteria (Sect. 2.4), six non-implausible or NROY LGM simulations remained. Table 4 gives
 312 the average volumes and areas of these six simulations and the corresponding six PGM ice sheets compared to estimated values
 313 from empirical and model data. All six LGM simulations show an overgrowth of ice in Alaska of varying magnitudes, as a
 314 result of the previously mentioned climate model bias. However, in other regions the simulations display a very similar ice
 315 extent, with the southern area only varying by $9.7 \times 10^5 \text{ km}^2$. None of the simulations form ice lobes, as expected, but they do
 316 show a close match to reconstructed ice extent in our target area, although towards the lower end of the plausible range, and
 317 in the marine regions (Fig. 6a and 7a). There is a minimum ice volume of 73.9 m SLE and a maximum of 97.1 m SLE. The
 318 maximum ice thickness varies by around 300 m but the overall shapes of the ice sheets remain the same, with the thickest ice
 319 towards the east of the ice sheet over Hudson Bay.



320
 321 **Figure 6. (a) The relationship between final ice volume and southern area for the LGM ensemble, and the relationship between the**
 322 **LGM and PGM (b) final ice volume, and (c) final southern areas. The filled in blue dots represent the six NROY LGM simulations**
 323 **and the solid lines on panel (a) show the minimum volume and area constraints applied to the ensemble. The ensemble member**
 324 **chosen as NROYa is outlined in red (Sect 2.5).**

325 All the PGM ice sheets were smaller in volume than their LGM counterpart (Figs. 6 and 7) and displayed a smaller extent in
 326 the southern margin and the saddle region between the western Cordilleran Ice Sheet and eastern Laurentide Ice Sheet.
 327 However, the PGM simulations also displayed more variability in their ice extent and volumes. The ice volumes range from
 328 53.4 m SLE to 83.37 m SLE and the southern extent varies by $2.44 \times 10^6 \text{ km}^2$. The range in maximum ice thickness is also
 329 over double the LGM, varying by around 613 m. These PGM configurations also look plausible compared to the less well
 330 constrained extent data available, including previous empirical and modelled reconstructions of the PGM/MIS 6 extent
 331 (Menviel et al., 2019; Batchelor et al., 2019; Fig. 7b). For example, all the simulations maintain an ice-free corridor between
 332 the Laurentide and Cordilleran ice sheets which is a common feature in these PGM reconstructions. In addition, the excess
 333 Alaskan ice seen in LGM simulations is also present at the PGM, however the growth is not as excessive.

334



335

336

337

338

339

Figure 7. Percentage of simulations with ice cover for (a) LGM with the Dalton et al., (2020) reconstructed margin shown in red; (b) PGM with the PMIP PGM modelled margin shown in solid red and the Batchelor et al., (2019) reconstructed maximum MIS 6 margin shown in dashed red, and (c) the difference between the LGM and PGM, at the end of the simulations for the six NROY ensemble members.

340

3.2 Impact of initial ice sheet vs climate

341

342

343

344

345

346

347

348

349


350

Out of our six NROY model configurations, we selected the parameters of a pair of LGM and PGM experiments xpken/xpkyn (NROYa; Fig. 6) to perform two sets of four sensitivity experiments to decompose the effects of climate forcing and initial conditions on the final ice sheet volume. This included repeating xpken and xpkyn using matching FAMOUS and Glimmer LGM and PGM initial conditions respectively (Table 2, experiments 3 and 6). For both glacial maxima, using the matching initial conditions resulted in more excess ice over Alaska (Fig. C1), though the southern ice extents are relatively similar between the two sets of experiments. Overall, for the LGM, using the GLAC-1D reconstruction in Glimmer ($V_{i,1}$) resulted in an ice sheet 9.7 m SLE larger than if the 18.2 ka ice sheet was used ($V_{i,2}$) (Table 5; Fig. C1a). For the PGM, the matching initial conditions ($V_{ci,1}$) resulted in only 0.45 m SLE increase from the NROYa simulation ($V_{ci,2}$) due to a decrease in ice volume over the Laurentide ice sheet (Table 5; Fig. C1b).

351 **Table 5. Final ice volumes of the four sensitivity experiments performed with matching climate model and ice sheet model ice sheets**
 352 **and the equivalent four performed with different initial ice sheets in each model**

<i>Experiment</i>	<i>Final ice volume (m SLE)</i>	<i>Experiment</i>	<i>Final ice volume (m SLE)</i>
$V_{_1}$ (full LGM)	100.3	$V_{_2}$	90.6
V_{c_1} (LGM ice , PGM climate)	104.2	V_{c_2}	97.1
V_{i_1} (PGM ice, LGM climate)	64.7	V_{i_2}	63.0
V_{ci_1} (full PGM)	68.6	V_{ci_2}	68.1


353
 354 The final ice sheet volumes from the first set of four sensitivity experiments (Table 2; experiments 3 – 6) are displayed in
 355 Table 5 and shown in Fig. 8. The results of the second set of four experiments (Table 2; experiments 7 – 10) are also included
 356 in Table 5. The results of the factor decomposition analysis show that the simulated ice volume at the PGM was 31.7 m SLE
 357 ($1.25 \times 10^7 \text{ km}^3$) lower than at the LGM ($dV_{_1}$). The initial ice sheet configuration (dV_{i_1}) alone caused a 35% decrease in
 358 volume, but this was partially offset by the climatic conditions (dV_{c_1}), which resulted in an increase in volume of 4%. The
 359 result was similar for the second set of experiments, with the initial ice sheet configuration (dV_{i_2}) causing a decrease of 31%
 360 in ice volume at the PGM compared to the LGM, but the climate (dV_{c_2}) caused a 6% increase in volume.



361
 362 **Figure 8. Final ice thickness in the sensitivity tests using (a) LGM ice sheets and LGM climate; (b) LGM ice sheets and PGM climate;**
 363 **(c) PGM ice sheets and LGM climate, and (d) PGM ice sheets and PGM climate.**

364 The PGM climate is conducive to growing a larger ice sheet (Fig. 9a) because the orbital configuration results in the Northern
 365 Hemisphere receiving less incoming solar radiation in spring and early summer (Table 1; Fig 2a). This reduces the melting of
 366 snow that has accumulated in winter (Fig. 9b). The winter snow accumulation is also higher at the PGM than at the LGM (Fig.
 367 9c) due to the PGM having warmer air temperatures in autumn and winter, because of the orbital forcing, leading to a wetter
 368 climate. Summer SSTs are also cooler at the PGM (Fig. 2c) due to lower spring insolation, further contributing to reduced

369 runoff. In contrast, the Greenland ice sheet decreases in size due to PGM climate conditions (Fig. 9a), likely due to higher sea
370 ice concentration south of Greenland reducing the moisture source available for precipitation.




371
372 **Figure 9. Difference between experiment $V_{ci,1}$ (full PGM) and $V_{i,1}$ (PGM ice sheet with LGM climate) isolating the effect of LGM**
373 **climate vs PGM climate on (a) final ice thickness simulated by Glimmer and (b) spring (MAM) runoff and (c) winter (DJF) snowfall**
374 **over the first 10 years.**

375 3.3 Uncertainty due to model parameters

376 Due to the sampling strategy, this ensemble does not have an optimal design for analysing the sensitivity of the ice sheets
377 during the two time periods to the different model parameter values because our ensemble of simulations does not uniformly
378 span the uncertain parameter space. For this, we refer the reader to the studies of Gandy et al., (2023) and Sherriff-Tadano et
379 al., (2023), which present larger ensembles of experiments. Here, we first evaluate if our results are consistent with these two
380 studies before examining if the difference between the PGM and LGM ice sheets is sensitive to specific model parameters.

381 Based on correlations between the parameters and ice sheet area and volume, we find that the LGM and PGM behave similarly
382 across the parameter ranges (Figs. E1 and E2) and most of the uncertainty in the results for both periods can be explained by
383 parameters that affect the surface albedo of the ice sheet; $Daice$, AV_GR and to a lesser extent, $Fsnow$. Higher values of $Daice$
384 and $Fsnow$ and lower values of AV_GR cause higher albedos and lead to larger ice sheets (Table 3). *Basal sliding* also

385 influences the volume of the ice sheet, with less impact on the area, with lower values and thus lower ice velocities causing
 386 larger volume ice sheets. The cloud parameter CW also shows a relatively high positive correlation for the PGM (Fig. 10).
 387 This is consistent with the findings of previous studies and current understanding on the importance of albedo for ice sheet
 388 evolution (Willeit and Ganopolski, 2018; Sherriff-Tadano et al., 2023; Gandy et al., 2023).
 389 Additionally, there is a negative correlation between the difference in ice volume and area between the LGM and PGM and
 390 the parameters AV_GR , $basal\ sliding$, and $RHCrit$. Conversely, there is a positive correlation between the LGM-to-PGM
 391 difference in ice volume/area and $Daice$ (Fig. E3). This suggests that lower values of AV_GR and higher values of $Daice$ and
 392 thus a higher albedo, as well as lower ice sheet velocity and more cloud, make the ice sheet more sensitive to changes in
 393 radiative forcings from the orbital boundary conditions.
 394



395
 396 **Figure 10. Relationship between LGM southern area and the four most influential parameters. The green shaded region shows the**
 397 **southern area constraint applied with the dotted line showing the exact area of the reconstruction and the solid line the minimum**

398 **bound applied. The colour scale represents ice volume and the dots outlined in red are the six NROY LGM simulations with the red**
399 **line on the colour bar showing the volume constraint.**


400 **4 Discussion**

401 After constraining our ensembles based on the available empirical and model data for the LGM, we find that the model was
402 able to successfully simulate the ice sheet at both periods under different LGM and PGM climate boundary conditions (orbital
403 parameters, SSTs and global orography) and initial ice sheets. However, the southern extents of the constrained LGM
404 simulations all fall towards the lower end of the plausible range, which is a common feature seen in other simulations using a
405 low resolution atmosphere model due to biases that cause a reduced stationary wave effect over this region (Ziemen et al.,
406 2014; Sherriff-Tadano et al., 2023; Gandy et al., 2023). Additionally, the ice lobes that are present over the Great Lakes are
407 not captured in these simulations. Again, this is common in ice sheet models and is likely a result of missing subglacial
408 processes or the low resolution of the climate and ice sheets models.

409 Analysis of the behaviour of the modelled ice sheets across the parameter spaces reveals that both the LGM and PGM ice
410 volume and extent have similar sensitivities to parameter uncertainties. We therefore conclude that parameters that produce a
411 good LGM NAIS also produce a plausible PGM NAIS under PGM boundary conditions and thus similar model parameters
412 are appropriate for use when modelling both periods. Our simulations can thus be compared and analysed to understand the
413 causes of the different configurations between the two periods. However, since the ice volume is most sensitive to surface
414 albedo and most simulations deglaciade under low values of *Daice*, this suggests that the value of bare ice albedo in the model
415 may need to be increased for future work.

416 The results of the sensitivity analysis show that the difference in initial ice sheet boundary conditions overwhelmingly
417 determined the difference in final ice volume between the LGM and PGM in the ensemble of simulations. We tested the impact
418 of starting from LGM and PGM ice sheet configurations in Glimmer instead of the 18.2 ka BP ice sheet and found that this
419 caused an even larger difference in ice volume between the two glacials. Comparing the simulations that use the same initial
420 ice topography in FAMOUS and Glimmer (first set of experiments), to those that use different topographies (second set of
421 experiments), whilst keeping the ice cover consistent, reveals that the relative contribution from the initial ice sheet boundary
422 conditions, compared to the climate conditions, to the simulated differences between the LGM and PGM ice sheets, remains
423 similar. This suggests that the dominant feedback responsible for this result is the ice-albedo feedback rather than the
424 temperature-elevation feedback. A similar conclusion was obtained by Abe-Ouchi et al., (2007) who studied the relative
425 contribution to climate over ice sheets from the ice sheet itself and the orbital parameters and CO₂ concentration. They found
426 the cooling caused by the ice sheet themselves was the dominant effect, mostly due to albedo feedbacks, which increase with
427 ice sheet area. Kageyama et al., (2004) also highlighted in their study the importance of the albedo feedback on the maximum
428 modelled North American ice volume. They show that changes in vegetation are needed to initiate glaciation over North
429 America which is then accelerated by the ice-albedo feedback. The North American ice sheet was larger at the LGM than at
430 the PGM. However, this sensitivity analysis reveals that the difference in orbital parameters, GHGs and SSTs (climate)

431 between the LGM and PGM encourages the growth of a larger North American ice sheet at the PGM (Fig. 9a). This effect
432 would likely be even stronger if we had used the orbit at 137 ka BP (the timing of the minimum in Northern Hemisphere
433 summer insolation; Fig. 11a-c) since the PGM would have received even lower insolation in spring and early summer. This
434 result highlights the importance of the evolution of these climate factors and the ice sheets during the preceding glacial cycles
435 in determining the glacial maxima configurations. For example, during the start of the Last Glacial Cycle (MIS 5; ~115-80 ka
436 BP), the variation in 65° N summer insolation was relatively large as a result of changes in orbital parameters (Fig 11a-c),
437 which resulted in multiple cycles of growth and recession of the North American Ice Sheets during this period, but total ice
438 volume remained low (Bonelli et al., 2009; Ganopolski et al., 2010; Dalton et al., 2022). Insolation then reaches a minimum
439 at ~70 ka BP (Fig 11c) which, combined with decreasing concentrations of CO₂ (~190 ppm at ~65 ka BP; Fig. 11f), led to a
440 significant increase in ice sheet volume to almost LGM extent (Fig. 11d) and a switch to more widespread glacial conditions
441 at the MIS 5/MIS 4 transition (Bonelli et al., 2009; Dalton et al., 2022). The size of the NAIS at this time was large enough to
442 induce positive feedbacks, such as the ice-albedo feedback, allowing its maintenance throughout MIS 4 and MIS 3 (~70-30 ka
443 BP) despite an increase in insolation from ~50-30 ka BP (Fig. 11c). This was also supported by a continued decrease in CO₂
444 (Fig. 11f). Growth of the ice sheet could then continue to its glacial maximum extent following a further insolation and CO₂
445 decrease during MIS 2 (~30-21 ka BP) (Fig. 11c-f). In contrast, prior to the PGM there were peaks in insolation at ~172 and
446 ~148 ka BP that reached higher levels than were reached prior to the LGM during MIS 4 and MIS 3 (Fig. 11c; Berger; 1978).
447 This may have inhibited an initial significant build-up of ice over North America, as during MIS 4, preventing the initiation of
448 an ice-albedo feedback strong enough to enable the continued growth towards a larger LGM configuration and/or maintain its
449 volume through the second insolation peak. In addition, there was more time between the LGM and the insolation maximum
450 at ~50-30 ka BP compared to the PGM and the maximum at ~147 ka BP. Therefore, the PGM NAIS may have not had enough
451 time to regrow before insolation started to increase again. Thus, investigation of the processes and interactions that took place
452 prior to the glacial maxima will be needed to fully understand why the LGM and PGM NAIS configuration differed.



453

454

455

456

457

458

Figure 11. Evolution of climate proxies over the last two glacial-interglacial cycles: (a) precession index (red) with eccentricity as an envelope (yellow); (b) obliquity (Berger, 1978); (c) July insolation at 65° N (Berger and Loutre, 1999); (d) reconstruction of global mean sea level and uncertainty estimate (dotted lines) (Waelbroeck et al., 2002); (e) benthic $\delta^{18}\text{O}$ global stack record (Lisiecki and Raymo, 2005), and (f) EPICA Dome C carbon dioxide ice core records (Luthi et al., 2008; Bereiter et al., 2015). The PGM and LGM are indicated by the dotted line.

459

460

461

462

463

Additional feedbacks that played a role in the development of glacials into either an LGM-like or PGM-like mode are also missing in these simulations due to computational constraints. For example, the low resolution of the atmospheric component of FAMOUS means that it is capable of performing ensembles and long palaeo runs while directly coupled to an ice sheet model. However, it also means that many small-scale atmospheric processes (e.g. stationary wave response) caused by and affecting the ice sheet topography are not represented well (Kageyama and Valdes, 2000; Liakka and Nilsson, 2010; Beghin

464 et al., 2014; 2015; Liakka et al., 2012; 2016). Additionally, the shallow ice approximation used in Glimmer means that the ice
465 sheet will not be able to simulate marine instabilities of advance and retreat (Pattyn et al., 2012). This effect will be minimal
466 for the NAIS, but a more advanced ice sheet model would be required to simulate a marine ice sheet like the EIS.

467 As a reminder, the vegetation was kept fixed at pre-industrial distributions, but the vegetation prior to and next to the ice cover
468 has been shown to be very important for determining ice sheet expansion in models through the vegetation-albedo feedback
469 (Kageyama et al., 2004; Colleoni et al., 2009b; Horton et al., 2010; Stone and Lunt, 2013). Therefore, implementing glacial
470 maxima distributions or dynamical vegetation may affect the results since the reduction in forest and expansion of
471 tundra/shrubs compared to present day would increase the albedo of the surface next to the ice and affect the climate (Meissner
472 et al., 2003). Similarly, the prescribed SSTs and sea ice concentrations used introduce an additional source of uncertainty. As
473 well as impacting the global mean temperature and precipitation patterns in the simulations, the SSTs and sea ice used can
474 have local climate impacts that affect the simulated ice sheets. This includes causing a warming or cooling over the more
475 coastal areas affecting the melt rate, and impacting evaporation rates, which affects the amount of snowfall the ice sheets
476 receive. The SSTs used in this study are cooler (as a global average) than the multi-proxy and data assimilation LGM SST
477 reconstructions of Tierney et al., (2020) and Paul et al., (2020) and the constrained statistical reconstruction of Gandy et al.,
478 (2023) and Astfalck et al., (2024). HadCM3 also tends to simulate cooler SSTs compared to other PMIP4 models, although
479 they are similar to CESM1.2 (Kageyama et al., 2021). Therefore, the use of colder SSTs in this study causes lower global mean
480 temperature overall, but also would have caused a cooling next to the ice sheets and reduced snowfall, which would have
481 impacted the ice sheet growth in different ways (Marsiat and Valdes, 2001; Hofer et al., 2012; Astfalk et al., 2024). The latter
482 impact was shown to be most dominant in the study by Astfalck et al., 2024, suggesting that our simulated ice sheet volumes
483 may have been larger had we used their warmer LGM SST reconstruction, due to increased evaporation. Prescribing the ocean
484 forcing also neglects any effects changes in ocean conditions and ice sheets have on each other (e.g. Timmerman et al., 2010;
485 Colleoni et al., 2011; Ullman et al., 2014; Sherriff-Tadano et al., 2018; 2021). Using a dynamical ocean would include the
486 effects of meltwater and changes in atmospheric circulation, arising from the ice sheets, on ocean circulation and temperature,
487 which would in turn affect the climate, feeding back onto the ice sheets themselves. Further work will be required to investigate
488 the feedbacks between ice sheets and sea surface at the PGM, but this is beyond the scope of this study. We recommend the
489 use of a fully coupled atmosphere-ocean-vegetation-ice sheet model to further investigate these feedbacks. The effect of dust
490 deposition and ice dammed lakes have also been shown to have a large influence on the build-up of ice (e.g. Krinner et al.,
491 2004; 2006; Naafs et al., 2012; Colleoni et al., 2009a) however further model developments would be needed to investigate
492 these effects.

493 Finally, the Eurasian ice sheet also displayed important differences between the LGM and PGM and had a large influence on
494 the climate. It is likely that some of the differences in the configurations of the NAIS and EIS between the two glacial maxima
495 resulted from their interactions with each other (Beghin et al., 2014; 2015; Liakka et al., 2016). To investigate the EIS at the
496 PGM, we recommend the use of an efficient marine ice sheet model such as BISICLES that uses Adaptive Mesh Refinement

497 to refine the processes occurring at marine margins that are more important for the marine based Eurasian ice sheet (Cornford
498 et al., 2013; Gandy et al., 2019).

499 **5 Conclusions**

500 We have performed and compared ensemble simulations of the LGM and PGM using a coupled atmosphere-ice sheet model
501 (FAMOUS-ice) with prescribed surface ocean conditions and interactive North American and Greenland Ice Sheets. We tested
502 the relative importance of the initial ice sheet configuration versus the climate boundary conditions on the resulting ice sheet
503 volumes through sensitivity tests and factor decomposition analysis. The main conclusions of this study are as follows:

- 504 1. Successful simulations of the LGM and PGM North American and Greenland ice sheets are produced using a coupled
505 climate-ice sheet model. We find that uncertain model parameters tuned to produce a plausible LGM North American
506 Ice Sheet also perform well for the PGM.
- 507 2. The initial ice extents used as boundary conditions in coupled climate-ice sheet simulations have a much larger impact
508 on the modelled NAIS than the climate boundary conditions, causing a ~30% decrease in ice volume at the PGM
509 compared to the LGM. This is due to the ice-albedo feedback.
- 510 3. The climate of the PGM causes an increase in NAIS ice volume of ~6% compared to the LGM due to the orbital
511 configuration causing the Northern Hemisphere to receive less insolation in spring and early summer. Since the LGM
512 ice sheet was larger than the PGM, this suggests that the climate and ice sheet evolution prior to the glacial maxima
513 contributes to the differences seen between the LGM and PGM ice sheets.

514 **Appendix A: Eccentricity equation correction**

515 The equation for the role of eccentricity on solar insolation used in the simulations in this paper was:

516

$$517 S(t) = S_o \left(\left(1 + \frac{e^2}{2} \right) (1 + e \cos v) / (1 - e^2) \right)^2 \quad (4)$$

518

519 However, this is incorrect and has now been corrected in the model to:

520


$$521 S(t) = S_o ((1 + e \cos v) / (1 - e^2))^2 \quad (5)$$

522

523 where; S(t) is the incoming solar insolation, S_o is the solar constant, e is the eccentricity of the earth's orbit and v is
524 the true anomaly (the angle of earth's current position on its orbit).


525 The PGM experiment 'xpkv0' was re-run with the correct equation and shows that on average the SMB was slightly lower in
526 our simulations than it should have been (decreased by 16% at the end of the simulations), leading to slightly smaller ice sheets

527 (Fig. A1). However, the impact is small (and would be even smaller for the LGM given the lower eccentricity) and does not
528 affect our overall conclusions.




529
530 **Figure A1. (a) Difference between the SMB at the end of the experiments between the original simulation and the simulation using**
531 **the corrected eccentricity equation and (b) the evolution of ice sheet volume for both experiments.**

532 Appendix B: Sea surface temperatures



533
534 **Figure B1. Mean annual SSTs used in this study from HadCM3 for (a) LGM and (b) PGM and (c) the difference between the LGM**
535 **SST reconstruction used in Gandy et al., (2023) and the HadCM3 LGM SSTs.**



537

538 **Figure C1. Difference in the final ice thickness between the simulations with matching initial conditions in FAMOUS and Glimmer**
 539 **and the NROYa ensemble member for (a) the LGM and (b) the PGM.**

540 **Appendix D: Wave 2 methodology**

541 The ensemble design in this study was based on the ‘Not Ruled Out Yet’ (NROY) parameter combinations from a second
 542 wave of ensemble members that followed on from the 280 member ensemble performed in Gandy et al., (2023). From the first
 543 wave of simulations, only 18 out of these 280 members produced a large enough LGM North American Ice sheet to meet the
 544 volume and extent criteria they imposed (see details in reference). Further work was thus performed to augment the ensemble
 545 of simulations that met the NROY criteria. We used statistical emulation to identify plausible regions in the parameter space.
 546 As there was limited information to constrain the domain of plausibility in the parameter space, we instead implemented an
 547 early-stopping criteria that allowed us to prevent the full execution of model runs that were not expected to produce good ice
 548 sheets. To do this we first modelled, from Wave 1, the predicted equilibrium area of the ice sheet from the value of the initial
 549 surface mass balance. Mathematically, we specified;

550

$$551 \quad A = f(b) + \epsilon, \quad (6)$$

552

553 where A is the ‘equilibrium’ ice sheet area after 10,000 ice sheet years, b is the 20 year averaged SMB value over the
 554 ice sheet and $f(\cdot)$ may be any function. We considered f to be either linear or sampled from a Gaussian Process (GP) and found
 555 the linear model gave more conservative uncertainty estimates which was desired since the Wave 2 runs needed to bound the

556 NROY space. The predictive interval for the model is $P(b)=[f(b) + 3\sqrt{\text{var}(\epsilon)}, f(b) - 3\sqrt{\text{var}(\epsilon)}]$ and we targeted equilibrium ice
557 sheet areas in the interval $T = [1.5 \times 10^7 \text{ km}^2, 2 \times 10^7 \text{ km}^2]$. The interval T is analogous to the target interval defined using
558 Pukelsheim's 3-sigma rule in standard history matching (Pukelsheim, 1994). Plausible values of b satisfy the condition that
559 $P(b) \cap T$ is non-zero, that is, for b to be plausible, the predictive bound $P(b)$ and the plausible equilibrium ice sheet area T
560 must intersect. It was found that the 20-year averaged SMB had to be at least positive to produce a plausible ice sheet.

561 To further improve efficiency, we used statistical emulation to produce plausible values of b (and hence equilibrium ice sheet
562 areas); iterating the training data of the emulator with each wave of simulator runs. Define by \mathbf{x} the multivariate vector of
563 parameters that they build the emulator over: here \mathbf{x} comprised of the 4 most influential parameters *Fsnow*, *AV_GR*, *Daice*,
564 and *Flow Factor*. We model b with a random error process, $b \sim GP(\mathbf{x}) + \eta$, where the effects of the parameters not explicitly
565 represented in \mathbf{x} are handled by the stochasticity of the process represented by η . Values of b were sampled using a stratified
566 k-extended Latin Hypercube design (Williamson, 2015) and three sub-waves were executed, from which, a candidate set for
567 the Wave 2 ensemble was extracted.

568 The first sub-wave (Wave 1.1) samples 200 ensemble members, which are predicted from the emulator to have non-negligible
569 probability of positive SMB. This results in around 50% of simulations in this sub-wave having a positive SMB, an increase
570 from 15% in the original wave (Fig. D1, Wave 1.1). We attempt to refine the predictive bounds on the GP model twice more
571 (Fig. D1, Wave 1.2 and 1.3), with no improvement. This is likely due to the inherent stochasticity of the climate model and
572 cumulative effects of the parameters that they absorb into the predictive error term. At the end of this process of iterative short
573 waves, the candidate set contains over 1000 20-year long simulations that have a positive SMB over the North American ice
574 sheet. From this candidate set, and again using stratified k-extended Latin Hypercubes, we select an optimal (with respect to
575 space-filling and accounting for the previous Wave 1 runs) design of 200 ensemble members to continue for a full 10,000
576 years to an equilibrium North American Ice Sheet. These 200 simulations make up the Wave 2. For context, this workflow of
577 GP model sub-waves saved around 230,000 core hours (or about two months of real time) compared to running a full second
578 ensemble wave.


579 Out of these 200 Wave 2 simulations, 176 members were identified to be NROY based on the original volume and extent
580 thresholds. It is based on these results that we sub-sampled 62 parameter combinations for our simulations. This number of
581 simulations was selected to enable us to run long equilibrium LGM and PGM simulations over a full ensemble within
582 reasonable computational requirements. From the 176 NROY parameter combinations we randomly generated 10^7 candidate
583 designs of size 62 from which we selected an approximate maximin design. This is obtained by: first linearly transforming
584 each parameter onto the same range of $[0, 1]$ to aid comparability; before computing the minimum distance between a
585 parameter vector and its nearest neighbour; and then selecting the candidate design that maximised this distance. The resulting
586 design possesses parameter vectors which are well-spaced and thus adequately cover the NROY space.

587 Our simulations use slightly different orbital parameter values and sea surface conditions to that of Gandy et al., (2023) (see
588 Sect. 2.3). Thus, we do not expect the sample of 62 parameter combinations to provide full coverage of the NROY space but,

589 as seen in Sect. S2 of the supplementary information in Gandy et al., (2023), the output trends are sufficiently similar that we
 590 expect this to be close enough to an optimal sample. Whilst we may have also sampled some parameter combinations outside
 591 of the NROY space, we feel these will still provide valuable information about uncertainty in outputs at the LGM and PGM.
 592 Our detailed comparison to empirical evidence and other model data (see Sect. 2.4 and 3.1) identified six parameter
 593 combinations that match our criteria for LGM and PGM ice extent and volume, thus demonstrating the success of this approach.
 594 Further exploration of the parameter space may produce NROY simulations in a different part of the parameter space but
 595 would not change the conclusion of this paper.

596 Upon analysing the results, we found a technical error in the original Wave 2 ensemble which resulted in the values of the
 597 parameter *Daice* being shifted from its intended range of $-0.4-0\text{ K}^{-1}$ to $0-0.4\text{ K}^{-1}$, this means that the albedo of the bare ice was
 598 increasing with melting, which is likely not the case. This produced larger values of surface albedo and thus larger ice sheets
 599 in these Wave 2 simulations (not shown here). In the ensemble of simulations presented here, we corrected the *Daice* values
 600 to match the intended parameter range. In some simulations, the switch of *Daice* value from a large positive number to a large
 601 negative number would have resulted in a decrease in surface albedo and resulting ice sheet volume. This effect is negligible
 602 for values of *Daice* closer to zero.


603



604


605 **Figure D1. Ice volumes simulated in the successive ensemble sub-waves of simulations sampled to have a positive initial surface mass**
 606 **balance using the Gaussian Process emulator**

608



609

610 **Figure E1.** Southern area versus each of the 13 parameters varied for the LGM ensemble. The green shaded region shows the
 611 southern area constraint applied with the dotted line showing the exact area of the reconstruction and the solid line the solid line the
 612 minimum bound applied. The colour scale represents ice volume and the dots outlined in red are the six NROY LGM simulations
 613 with the red line on the colour bar showing the volume constraint.




614

615

Figure E2. Southern area versus each of the 13 parameters varied for the PGM ensemble. The colour scale represents ice volume and the dots outlined in red are the corresponding six NROY PGM simulations.

616



617

618

Figure E3. Difference in southern area versus each of the 13 parameters varied between the LGM and PGM ensemble members.

619

The colour scale represents difference in ice volume and the dots outlined in red are the six NROY simulations.

620 **Data availability**

621 The boundary conditions used in this study as well as the full ensemble ice sheet model output and volume and extent metrics,
622 climate timeseries for the NROY simulations and final ice volume data from the sensitivity tests are available at
623 <https://catalogue.ceda.ac.uk/uuid/5e48b31e413b480792e4156191b654f4>. All other model output data are available on request.

624 **Author contributions**

625 VLP lead the project and performed the majority of the work. VLP, LJG, RFI and NG designed the simulations and VLP and
626 NG prepared the initial and boundary conditions with support from OGP. VLP ran the simulations and analysed the results.
627 LCA and NG designed and performed the Wave 2 simulations the ensembles were sampled from, and JO did the sampling.
628 RSS provided technical and scientific support and updates for FAMOUS-Glimmer. PJV provided the PGM HadCM3 sea
629 surface temperature and sea ice dataset. VLP wrote the manuscript with comments and contributions from all co-authors. LJG,
630 RFI and NG supervised the project and LJG acquired the funding.

631 **Competing interests**

632 The authors declare they have no conflict of interest.

633 **Acknowledgements**

634 This research is primarily funded by the ‘SMB-Gen’ UKRI Future Leaders Fellowship MR/S016961/1, with LJG, JO, NG and
635 LCA supported by the award and VLP’s PhD studentship funded by the University of Leeds. RFI and RSS’s contributions
636 were supported by the RISICMAP19 NERC standard grant NE/T007443/1. LCA is supported by the ARC ITRH for
637 Transforming energy Infrastructure through Digital Engineering (TIDE), Grant No. IH200100009. OGP is funded by the
638 European Union’s Horizon 2020 research and innovation programme RiSeR project (grant agreement no. 802281). The
639 simulations were run on the high-performance research computing facilities of the University of Leeds and technical support
640 was provided by Richard Rigby from the Centre for Environmental Modelling and Computation (CEMAC). The authors would
641 like to thank Michel Crucifix, Peter Hopcroft, Pam Vervoort and Paul Valdes for discovering the eccentricity equation error
642 and providing the corrected equation.

643 **References**

- 644 Abe-Ouchi, A., Segawa, T., and Saito, F.: Climatic Conditions for modelling the Northern Hemisphere ice sheets throughout
645 the ice age cycle, *Clim. Past*, 3, 423–438, <https://doi.org/10.5194/cp-3-423-2007>, 2007.
- 646 Abe-Ouchi, A., Saito, F., Kawamura, K., Raymo, M. E., Okuno, J., Takahashi, K., and Blatter, H.: Insolation-driven 100,000-
647 year glacial cycles and hysteresis of ice-sheet volume, *Nature*, 500, 190-193, <https://doi.org/10.1038/nature12374>, 2013.
- 648 Allen, J. R. M., Forrest, M., Hickler, T., Singarayer, J. S., Valdes, P. J., and Huntley, B.: Global vegetation patterns of the past
649 140,000 years, *J. Biogeogr.*, 47, 2073-2090. <https://doi.org/10.1111/jbi.13930>, 2020.
- 650 Astfalck, L., Williamson, D., Gandy, N., Gregoire, L., and Ivanovic, R.: Coexchangeable Process Modeling for Uncertainty
651 Quantification in Joint Climate Reconstruction. *J. Am. Stat. Assoc.* 1–14. <https://doi.org/10.1080/01621459.2024.2325705>,
652 2024.
- 653 Batchelor, C. L., Margold, M., Krapp, M., Murton, D. K., Dalton, A. S., Gibbard, P. L., Stokes, C. R., Murton, J. B., and
654 Manica, A.: The configuration of Northern Hemisphere ice sheets through the Quaternary, *Nat. Commun.*, 10, 3713,
655 <https://doi.org/10.1038/s41467-019-11601-2>, 2019.
- 656 Beghin, P., Charbit, S., Dumas, C., Kageyama, M., Roche, D. M., and Ritz, C.: Interdependence of the growth of the Northern
657 Hemisphere ice sheets during the last glaciation: the role of atmospheric circulation, *Clim. Past*, 10, 345–358,
658 <https://doi.org/10.5194/cp-10-345-2014>, 2014.
- 659 Beghin, P., Charbit, S., Dumas, C., Kageyama, M., and Ritz, C.: How might the North American ice sheet influence the
660 northwestern Eurasian climate?, *Clim. Past*, 11, 1467–1490, <https://doi.org/10.5194/CP-11-1467-2015>, 2015.
- 661 Bereiter, B., Eggleston, S., Schmitt, J., Nehrbass-Ahles, C., Stocker, T. F., Fischer, H., Kipfstuhl, S., and Chappellaz, J.:
662 Revision of the EPICA Dome C CO₂ record from 800 to 600-kyr before present, *Geophys. Res. Lett.*, 42, 542–549,
663 <https://doi.org/10.1002/2014GL061957>, 2015.
- 664 Berger, A. L.: Long-Term Variations of Daily Insolation and Quaternary Climatic Changes, *J. Atmos. Sci.*, 35, 2362-2367,
665 [https://doi.org/10.1175/1520-0469\(1978\)035<2362:LTVODI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<2362:LTVODI>2.0.CO;2), 1978.
- 666 Berger, A., and Loutre, M. F.: Insolation values for the climate of the last 10 million years, *Quaternary Sci. Rev.*, 10, 297–317,
667 [https://doi.org/10.1016/0277-3791\(91\)90033-Q](https://doi.org/10.1016/0277-3791(91)90033-Q), 1991.
- 668 Bonelli, S., Charbit, S., Kageyama, M., Woillez, M.-N., Ramstein, G., Dumas, C., and Quiquet, A.: Investigating the evolution
669 of major Northern Hemisphere ice sheets during the last glacial-interglacial cycle, *Clim. Past*, 5, 329–345,
670 <https://doi.org/10.5194/cp-5-329-2009>, 2009.
- 671 Briggs, R. D., Pollard, D., and Tarasov, L.: A data-constrained large ensemble analysis of Antarctic evolution since the Eemian,
672 *Quaternary Sci. Rev.*, 103, 91-115, <https://doi.org/10.1016/j.quascirev.2014.09.003>, 2014.
- 673 Colleoni, F., Krinner, G., Jakobsson, M., Peyaud, V., and Ritz, C.: Influence of regional parameters on the surface mass balance
674 of the Eurasian ice sheet during the peak Saalian (140 kya), *Glob. Planet. Change*, 68, 132-148,
675 <https://doi.org/10.1016/j.gloplacha.2009.03.021>, 2009a.

676 Colleoni, F., Krinner, G., and Jakobsson, M.: Sensitivity of the Late Saalian (140 kyrs BP) and LGM (21 kyrs BP) Eurasian
677 ice sheet surface mass balance to vegetation feedbacks, *Geophys. Res. Lett.*, 36, L08704, doi:10.1029/2009GL037200, 2009b.

678 Colleoni, F., Liakka, J., Krinner, G., Jakobsson, M., Masina, S., and Peyaud, V.: The sensitivity of the Late Saalian (140 ka)
679 and LGM (21 ka) Eurasian ice sheets to sea surface conditions, *Clim. Dyn.*, 37, 531-553, [https://doi.org/10.1007/s00382-010-](https://doi.org/10.1007/s00382-010-0870-7)
680 [0870-7](https://doi.org/10.1007/s00382-010-0870-7), 2011.

681 Colleoni, F., Wekerle, C., Brandefelt, J., and Masina, S.: Constraint on the penultimate glacial maximum Northern Hemisphere
682 ice topography (~140 kyrs BP), *Quaternary Sci. Rev.*, 137, 97-112, <https://doi.org/10.1016/j.quascirev.2016.01.024>, 2016.

683 Cornford, S. L., Martin, D. F., Graves, D. T., Ranken, D. F., le Brocq, A. M., Gladstone, R. M., Payne, A. J., Ng, E. G., and
684 Lipscomb, W. H.: Adaptive mesh, finite volume modeling of marine ice sheets, *J. Comput. Phys.*, 232, 529–549,
685 <https://doi.org/10.1016/J.JCP.2012.08.037>, 2013.

686 Crossley, J. F., and Roberts, D. L.: Thermodynamic/dynamic Sea-ice model, Meteorological office, 1995.

687 Dalton, A. S., Margold, M., Stokes, C. R., Tarasov, L., Dyke, A. S., Adams, R. S., Allard, S., Arends, H. E., Atkinson, N.,
688 Attig, J. W., Barnett, P. J., Barnett, R. L., Batterson, M., Bernatchez, P., Borns Jr, H. W., Breckenridge, A., Briner, J. P.,
689 Brouard, E., Campbell, J. E., Carlson, A. E., ... Wright Jr, H. E.: An updated radiocarbon-based ice margin chronology for the
690 last deglaciation of the North American Ice Sheet Complex, *Quaternary Sci. Rev.*, 234, 106223,
691 <https://doi.org/10.1016/j.quascirev.2020.106223>, 2020.

692 Dalton, A. S., Stokes, C. R., and Batchelor, C. L.: Evolution of the Laurentide and Innuitian ice sheets prior to the Last Glacial
693 Maximum (115 ka to 25 ka), *Earth-Sci. Rev.*, 224, 103875, <https://doi.org/10.1016/j.earscirev.2021.103875>, 2022.

694 Davies-Barnard, T., Ridgwell, A., Singarayer, J., and Valdes, P.: Quantifying the influence of the terrestrial biosphere on
695 glacial–interglacial climate dynamics, *Clim. Past*, 13, 1381–1401, <https://doi.org/10.5194/cp-13-1381-2017>, 2017.

696 de Boer, B., van de Wal, R.S.W., Lourens, L.J., Bintanja, R., and Reerink, T. J.: A continuous simulation of global ice volume
697 over the past 1 million years with 3-D ice-sheet models, *Clim. Dyn.*, 41, 1365–1384. [https://doi.org/10.1007/s00382-012-1562-](https://doi.org/10.1007/s00382-012-1562-2)
698 [2](https://doi.org/10.1007/s00382-012-1562-2), 2013.

699 Dentith, J. E., Ivanovic, R. F., Gregoire, L. J., Tindall, J. C., and Smith, R. S.: Ocean circulation drifts in multi-millennial
700 climate simulations: the role of salinity corrections and climate feedbacks, *Clim. Dyn.*, 52, 1761–1781,
701 <https://doi.org/10.1007/S00382-018-4243-Y/FIGURES/15>, 2019.

702 Dyer, B., Austermann, J., D’Andrea, W. J., Creel, R. C., Sandstrom, M. R., Cashman, M., Rovere, A., and Raymo, M. E.: Sea-
703 level trends across the Bahamas constrain peak last interglacial ice melt, *Proc. Natl. Acad. Sci. U.S.A.*, 118, 33,
704 <https://doi.org/10.1073/pnas.2026839118>, 2021.

705 Dyke, A.S., Andrews, J. T. Clark, P. U., England, J. H., Miller, G. H., Shaw, J., and Veillette, J. J.: The Laurentide and Innuitian
706 ice sheets during the Last Glacial Maximum, *Quaternary Sci. Rev.*, 21, 9-31, [https://doi.org/10.1016/S0277-3791\(01\)00095-](https://doi.org/10.1016/S0277-3791(01)00095-6)
707 [6](https://doi.org/10.1016/S0277-3791(01)00095-6), 2002.

708 Ehlers, J., Gibbard, P.L. and Hughes, P.D.: Chapter 4 – Quaternary Glaciations and Chronology, in: Past Glacial Environments,
709 Second Edition, edited by: Menzies, J., and van der Meer, J.J.M., Elsevier, 77-101, [https://doi.org/10.1016/B978-0-08-100524-](https://doi.org/10.1016/B978-0-08-100524-8.00003-8)
710 [8.00003-8](https://doi.org/10.1016/B978-0-08-100524-8.00003-8), 2018.

711 Essery, R., Best, M., Betts, R., Cox, P. and Taylor, C.: Explicit Representation of Subgrid Heterogeneity in a GCM Land
712 Surface Scheme, *J. Hydrometeorol.*, 4, 530-543, [https://doi.org/10.1175/1525-7541\(2003\)004<0530:EROSHI>2.0.CO;2](https://doi.org/10.1175/1525-7541(2003)004<0530:EROSHI>2.0.CO;2),
713 2003.

714 Fyke, J. G., Weaver, A. J., Pollard, D., Eby, M., Carter, L., and Mackintosh, A.: A new coupled ice sheet/climate model:
715 description and sensitivity to model physics under Eemian, Last Glacial Maximum, late Holocene and modern climate
716 conditions, *Geosci. Model Dev.*, 4, 117–136, <https://doi.org/10.5194/gmd-4-117-2011>, 2011.

717 Gandy, N., Gregoire, L. J., Ely, J. C., Cornford, S. L., Clark, C. D., and Hodgson, D. M.: Exploring the ingredients required
718 to successfully model the placement, generation, and evolution of ice streams in the British-Irish Ice Sheet, *Quaternary Sci.*
719 *Rev.*, 223, 105915, <https://doi.org/10.1016/j.quascirev.2019.105915>, 2019.

720 Gandy, N., Astfalck, L. C., Gregoire, L. J., Ivanovic, R. F., Patterson, V. L., Sherriff-Tadano, S., Smith, R. S., Williamson, D.,
721 and Rigby, R.: De-tuning a coupled Climate Ice Sheet Model to simulate the North American Ice Sheet at the Last Glacial
722 Maximum, *J. Geophys. Res. Earth Surf.*, DOI: 10.1002/essoar.10512201.1, 2023.

723 Ganopolski, A. and Brovkin, V.: Simulation of climate, ice sheets and CO2 evolution during the last four glacial cycles with
724 an Earth system model of intermediate complexity, *Clim. Past*, 13, 1695–1716, <https://doi.org/10.5194/cp-13-1695-2017>,
725 2017.

726 Ganopolski, A., Calov, R., and Claussen, M.: Simulation of the last glacial cycle with a coupled climate ice-sheet model of
727 intermediate complexity, *Clim. Past*, 6, 229–244, <https://doi.org/10.5194/cp-6-229-2010>, 2010.

728 Gowan, E.J., Zhang, X., Khosravi, S., Rovere, A., Stocchi, P., Hughes, A. L. C., Gyllencreutz, R., Mangerud, J., Svendsen, J-
729 I., and Lohmann, G.: A new global ice sheet reconstruction for the past 80 000 years, *Nat. Commun.*, 12, 1199,
730 <https://doi.org/10.1038/s41467-021-21469-w>, 2021.

731 Gregoire, L., J. Modelling the Northern Hemisphere Climate and Ice Sheets during the Last Deglaciation, Ph.D. thesis, School
732 of Geographical Sciences, University of Bristol, UK, 2010.

733 Gregoire, L., Payne, A. and Valdes, P.: Deglacial rapid sea level rises caused by ice-sheet saddle collapses, *Nature*, 487, 219–
734 222, <https://doi.org/10.1038/nature11257>, 2012.

735 Gregoire, L. J., Valdes, P. J., and Payne, A. J.: The relative contribution of orbital forcing and greenhouse gases to the North
736 American deglaciation, *Geophys. Res. Lett.*, 42, 9970-9979, <https://doi.org/10.1002/2015GL066005>, 2015.

737 Gregoire, L. J., Otto-Bliesner, B., Valdes, P. J., and Ivanovic, R.: Abrupt Bølling warming and ice saddle collapse contributions
738 to the Meltwater Pulse 1a rapid sea level rise, *Geophys. Res. Lett.*, 43, 9130–9137, <https://doi.org/10.1002/2016gl070356>,
739 2016.

740 Gregoire, L. J., Ivanovic, R. F., Maycock, A. C., Valdes, P. J., and Stevenson, S.: Holocene lowering of the Laurentide ice
741 sheet affects North Atlantic gyre circulation and climate, *Clim. Dyn.*, 51, 3797-3813, [https://doi.org/10.1007/s00382-018-](https://doi.org/10.1007/s00382-018-4111-9)
742 [4111-9](https://doi.org/10.1007/s00382-018-4111-9), 2018.

743 Gregory, J. M., Browne, O. J. H., Payne, A. J., Ridley, J. K., and Rutt, I. C.: Modelling large-scale ice-sheet–climate
744 interactions following glacial inception, *Clim. Past*, 8, 1565–1580, <https://doi.org/10.5194/cp-8-1565-2012>, 2012.

745 Gregory, J. M., George, S. E., and Smith, R. S.: Large and irreversible future decline of the Greenland ice sheet, *Cryosphere*,
746 14, 4299–4322, <https://doi.org/10.5194/TC-14-4299-2020>, 2020.

747 Heinemann, M., Timmermann, A., Elison Timm, O., Saito, F., and Abe-Ouchi, A.: Deglacial ice sheet meltdown: orbital
748 pacemaking and CO₂ effects, *Clim. Past*, 10, 1567–1579, <https://doi.org/10.5194/cp-10-1567-2014>, 2014.

749 Hemming, S. R.: Heinrich events: Massive late Pleistocene detritus layers of the North Atlantic and their global climate
750 imprint, *Rev. Geophys.*, 42, RG1005, <https://doi.org/10.1029/2003RG000128>, 2004.

751 Hofer, D., Raible, C. C., Dehnert, A., and Kuhle, J.: The impact of different glacial boundary conditions on atmospheric
752 dynamics and precipitation in the North Atlantic region, *Clim. Past*, 8, 935–949, <https://doi.org/10.5194/cp-8-935-2012>, 2012.

753 Horton, D., Poulsen, C. and Pollard, D.: Influence of high-latitude vegetation feedbacks on late Palaeozoic glacial cycles, *Nat.*
754 *Geosci.*, 3, 572–577, <https://doi.org/10.1038/ngeo922>, 2010.

755 Huybers, P.: Early Pleistocene Glacial Cycles and the Integrated Summer Insolation Forcing, *Science*, 313, 508-511,
756 DOI:10.1126/science.1125249, 2006.

757 Izumi, K., Valdes, P., Ivanovic, R., and Gregoire, L.: Impacts of the PMIP4 ice sheets on Northern Hemisphere climate during
758 the last glacial period, *Clim. Dyn.*, 60, 2481-2499, <https://doi.org/10.1007/s00382-022-06456-1>, 2023.

759 Kageyama, M., and Valdes, P. J.: Impact of the North American ice-sheet orography on the Last Glacial Maximum eddies and
760 snowfall, *Geophys. Res. Lett.*, 27, 1515-1518, <https://doi.org/10.1029/1999GL011274>, 2000.

761 Kageyama, M., Charbit, S., Ritz, C., Khodri, M., and Ramstein, G.: Quantifying ice-sheet feedbacks during the last glacial
762 inception, *Geophys. Res. Lett.*, 31, 24, doi:10.1029/2004GL021339, 2004.

763 Kageyama, M., Albani, S., Braconnot, P., Harrison, S. P., Hopcroft, P. O., Ivanovic, R. F., Lambert, F., Marti, O., Peltier, W.
764 R., Peterschmitt, J.-Y., Roche, D. M., Tarasov, L., Zhang, X., Brady, E. C., Haywood, A. M., Legrande, A. N., Lunt, D. J.,
765 Mahowald, N. M., Mikolajewicz, U., ... Zheng, W.: The PMIP4 contribution to CMIP6-Part 4: Scientific objectives and
766 experimental design of the PMIP4-CMIP6 Last Glacial Maximum experiments and PMIP4 sensitivity experiments, *Geosci.*
767 *Model Dev.*, 10, 4035–4055, <https://doi.org/10.5194/gmd-10-4035-2017>, 2017.

768 Kageyama, M., Harrison, S. P., Kapsch, M.-L., Lofverstrom, M., Lora, J. M., Mikolajewicz, U., Sherriff-Tadano, S., Vadsaria,
769 T., Abe-Ouchi, A., Bouttes, N., Chandan, D., Gregoire, L. J., Ivanovic, R. F., Izumi, K., LeGrande, A. N., Lhardy, F., Lohmann,
770 G., Morozova, P. A., Ohgaito, R., Paul, A., Peltier, W. R., Poulsen, C. J., Quiquet, A., Roche, D. M., Shi, X., Tierney, J. E.,
771 Valdes, P. J., Volodin, E., and Zhu, J.: The PMIP4 Last Glacial Maximum experiments: preliminary results and comparison
772 with the PMIP3 simulations, *Clim. Past*, 17, 1065–1089, <https://doi.org/10.5194/cp-17-1065-2021>, 2021.

773 Krinner, G., Mangerud, J., Jakobsson, M., Crucifix, M., Ritz, C., and Svendsen, J-I.: Enhanced ice sheet growth in Eurasia
774 owing to adjacent ice-dammed lakes, *Nature*, 427, 429–432, <https://doi.org/10.1038/nature02233>, 2004.

775 Krinner, G., Boucher, O., and Balkanski, Y.: Ice-free glacial northern Asia due to dust deposition on snow, *Clim. Dyn.*, 27,
776 613–625, DOI:10.1007/s00382-006-0159-z, 2006.

777 Krinner, G., Diekmann, B., Colleoni, F., and Stauch, G.: Global, regional and local scale factors determining glaciation extent
778 in Eastern Siberia over the last 140,000 years, *Quaternary Sci. Rev.*, 30, 821-831,
779 <https://doi.org/10.1016/j.quascirev.2011.01.001>, 2011.

780 Lambeck, K., Rouby, H., Purcell, A., Sun, Y., and Sambridge, M.: Sea level and global ice volumes from the Last Glacial
781 Maximum to the Holocene, *Proc. Natl. Acad. Sci. U.S.A.*, 111, 15296–15303, <https://doi.org/10.1073/pnas.1411762111>, 2014.

782 Liakka, J., and Nilsson, J.: The impact of topographically forced stationary waves on local ice-sheet climate, *J. Glaciol.*, 56,
783 534–544, <https://doi.org/10.3189/002214310792447824>, 2010.

784 Liakka, J., Nilsson, J. and Löfverström, M.: Interactions between stationary waves and ice sheets: linear versus nonlinear
785 atmospheric response, *Clim. Dyn.*, 38, 1249–1262, <https://doi.org/10.1007/s00382-011-1004-6>, 2012.

786 Liakka, J., Löfverström, M., and Colleoni, F.: The impact of the North American glacial topography on the evolution of the
787 Eurasian ice sheet over the last glacial cycle, *Clim. Past*, 12, 1225–1241, <https://doi.org/10.5194/CP-12-1225-2016>, 2016.

788 Lisiecki, L. E., and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}O$ records,
789 *Paleoceanography*, 20, 1–17, <https://doi.org/10.1029/2004PA001071>, 2005.

790 Loulergue, L., Schilt, A., Spahni, R., Masson-Delmotte, V., Blunier, T., Lemieux, B., Barnola, J. M., Raynaud, D., Stocker,
791 T. F., and Chappellaz, J.: Orbital and millennial-scale features of atmospheric CH_4 over the past 800,000 years, *Nature*, 453,
792 383–386, <https://doi.org/10.1038/nature06950>, 2008.

793 Lunt, D. J., Haywood, A. M., Schmidt, G. A., Salzmann, U., Valdes, P. J., Dowsett, H. J., and Loptson, C. A.: On the causes
794 of mid-Pliocene warmth and polar amplification. *Earth Planet. Sci. Lett.*, 321-322, 128-138,
795 <https://doi.org/10.1016/j.epsl.2011.12.042>, 2012.

796 Lüthi, D., le Floch, M., Bereiter, B., Blunier, T., Barnola, J. M., Siegenthaler, U., Raynaud, D., Jouzel, J., Fischer, H.,
797 Kawamura, K., and Stocker, T. F.: High-resolution carbon dioxide concentration record 650,000–800,000 years before present,
798 *Nature*, 453, 379–382. <https://doi.org/10.1038/nature06949>, 2008.

799 Margari, V., Skinner, L. C., Hodell, D. A., Martrat, B., Toucanne, S., Grimalt, J. O., Gibbard, P. L., Lunkka, J. P., and Tzedakis,
800 P. C.: Land-ocean changes on orbital and millennial time scales and the penultimate glaciation, *Geology*, 4, 183–186,
801 <https://doi.org/10.1130/G35070.1>, 2014.

802 Marshall, S. J., James, T. S., and Clarke, G. K. C.: North American Ice Sheet reconstructions at the Last Glacial Maximum,
803 *Quaternary Sci. Rev.*, 21, 175-192, [https://doi.org/10.1016/S0277-3791\(01\)00089-0](https://doi.org/10.1016/S0277-3791(01)00089-0), 2002.

804 Marsiat, I., and Valdes, P.: Sensitivity of the Northern Hemisphere climate of the Last Glacial Maximum to sea surface
805 temperatures. *Clim. Dyn.* 17, 233–248, <https://doi.org/10.1007/s003820000108>, 2001.

806 Masson-Delmotte, V., Stenni, B., Pol, K., Braconnot, P., Cattani, O., Falourd, S., Kageyama, M., Jouzel, J., Landais, A.,
807 Minster, B., Barnola, J. M., Chappellaz, J., Krinner, G., Johnsen, S., Röthlisberger, R., Hansen, J., Mikolajewicz, U., and Otto-
808 Bliesner, B.: EPICA Dome C record of glacial and interglacial intensities, *Quaternary Sci. Rev.*, 29, 113-128,
809 <https://doi.org/10.1016/j.quascirev.2009.09.030>, 2010.

810 Meissner, K. J., Weaver, A. J., Matthews, H. D., and Cox, P. M.: The role of land surface dynamics in glacial inception: a
811 study with the UVic Earth System Model, *Clim. Dyn.*, 21, 515-537, <https://doi.org/10.1007/s00382-003-0352-2>, 2003.

812 Menviel, L., Capron, E., Govin, A., Dutton, A., Tarasov, L., Abe-Ouchi, A., Drysdale, R. N., Gibbard, P. L., Gregoire, L., He,
813 F., Ivanovic, R. F., Kageyama, M., Kawamura, K., Landais, A., Otto-Bliesner, B. L., Oyabu, I., Tzedakis, P. C., Wolff, E., and
814 Zhang, X.: The penultimate deglaciation: protocol for Paleoclimate Modelling Intercomparison Project (PMIP) phase 4
815 transient numerical simulations between 140 and 127 ka, version 1.0, *Geosci. Model Dev.*, 12, 3649–3685,
816 <https://doi.org/10.5194/gmd-12-3649-2019>, 2019.

817 Naafs, B. D. A., Hefter, J., Acton, G., Haug, G. H., Martinez-Garcia, A., Pancost, R., and Stein, R.: Strengthening of North
818 American dust sources during the late Pliocene (2.7 Ma), *Earth Planet. Sci. Lett.*, 317-318, 8-19,
819 <https://doi.org/10.1016/j.epsl.2011.11.026>, 2012.

820 Naafs, B. D. A., Hefter, J., and Stein, R.: Millennial-scale ice rafting events and Hudson Strait Heinrich(-like) Events during
821 the late Pliocene and Pleistocene: a review, *Quaternary Sci. Rev.*, 80, 1-28, <https://doi.org/10.1016/j.quascirev.2013.08.014>,
822 2013.

823 Niu, L., Lohmann, G., Hinck, S., Gowan, E., and Krebs-Kanzow, U.: The sensitivity of Northern Hemisphere ice sheets to
824 atmospheric forcing during the last glacial cycle using PMIP3 models, *J. Glaciol.*, 65, 645-661, doi:10.1017/jog.2019.42, 2019.

825 Niu, L., Lohmann, G., Gierz, P., Gowan, E. J., and Knorr, G.: Coupled climate-ice sheet modelling of MIS-13 reveals a
826 sensitive Cordilleran Ice Sheet. *Glob. Planet. Change*, 200, 103474, <https://doi.org/10.1016/j.gloplacha.2021.103474>, 2021.

827 Niu, L., Knorr, G., Krebs-Kanzow, U., Gierz, P., and Lohmann, G.: Rapid Laurentide Ice Sheet growth preceding the Last
828 Glacial Maximum due to summer snowfall. *Nat. Geosci.* 17, 440–449, <https://doi.org/10.1038/s41561-024-01419-z>, 2024.

829 Obrochta, S. P., Crowley, T. J., Channell, J. E. T., Hodell, D. A., Baker, P. A., Seki, A., and Yokoyama, Y.: Climate variability
830 and ice-sheet dynamics during the last three glaciations, *Earth Planet. Sci. Lett.*, 406, 198–212,
831 <https://doi.org/10.1016/J.EPSL.2014.09.004>, 2014.

832 Parker, R. L., Foster, G. L., Gutjahr, M., Wilson, P. A., Littler, K. L., Cooper, M. J., Michalik, A., Milton, J. A., Crocket, K.
833 C., and Bailey, I.: Laurentide Ice Sheet extent over the last 130 thousand years traced by the Pb isotope signature of weathering
834 inputs to the Labrador Sea, *Quaternary Sci. Rev.*, 287, 107564, <https://doi.org/10.1016/j.quascirev.2022.107564>, 2022.

835 Pattyn, F., Schoof, C., Perichon, L., Hindmarsh, R. C. A., Bueler, E., de Fleurian, B., Durand, G., Gagliardini, O., Gladstone,
836 R., Goldberg, D., Gudmundsson, G. H., Huybrechts, P., Lee, V., Nick, F. M., Payne, A. J., Pollard, D., Rybak, O., Saito, F.,
837 and Vieli, A.: Results of the Marine Ice Sheet Model Intercomparison Project, MISMIP, *Cryosphere*, 6, 573–588,
838 <https://doi.org/10.5194/tc-6-573-2012>, 2012.

839 Paul, A., Mulitza, S., Stein, R., and Werner, M.: Glacial Ocean Map (GLOMAP), PANGAEA [dataset],
840 <https://doi.org/10.1594/PANGAEA.923262>, 2020.

841 Peltier, W. R., Argus, D. F., and Drummond, R.: Space geodesy constrains ice age terminal deglaciation: The global ICE-
842 6G_C (VM5a) model, *J. Geophys. Res. Solid Earth*, 120, 450–487, doi:10.1002/2014JB011176, 2015.

843 Pollard, O. G., Barlow, N. L. M., Gregoire, L. J., Gomez, N., Cartelle, V., Ely, J. C., and Astfalck, L. C.: Quantifying the
844 uncertainty in the Eurasian ice-sheet geometry at the Penultimate Glacial Maximum (Marine Isotope Stage 6), *Cryosphere*, 17,
845 4751-4777, <https://doi.org/10.5194/tc-17-4751-2023>, 2023.

846 Pöppelmeier, F., Joos, F., and Stocker, T.F.: The Coupled Ice Sheet–Earth System Model Bern3D v3.0. *J. Climate*, 36, 7563–
847 7582, <https://doi.org/10.1175/JCLI-D-23-0104.1>, 2023.

848 Pukelsheim, F.: The Three Sigma Rule, *Am. Stat.*, 48, 88-91, <https://doi.org/10.2307/2684253>, 1994.

849 Quiquet, A., Roche, D. M., Dumas, C., Bouttes, N., and Lhardy, F.: Climate and ice sheet evolutions from the last glacial
850 maximum to the pre-industrial period with an ice-sheet–climate coupled model, *Clim. Past*, 17, 2179–2199,
851 <https://doi.org/10.5194/cp-17-2179-2021>, 2021.

852 Rabineau, M., Berné, S., Olivet, J.-L., Aslanian, D., Guillocheau, F., and Joseph, P.: Paleo sea levels reconsidered from direct
853 observation of paleoshoreline position during Glacial Maxima (for the last 500,000 yr), *Earth Planet. Sci. Lett.*, 252, 119–137,
854 <https://doi.org/10.1016/j.epsl.2006.09.033>, 2006.

855 Roberts, W. H. G., Valdes, P. J., and Payne, A. J.: Topography's crucial role in Heinrich Events, *Proc. Natl. Acad. Sci. U.S.A.*,
856 111, 16688-16693, <https://doi.org/10.1073/pnas.1414882111>, 2014.

857 Robinson, A., Calov, R., and Ganopolski, A.: Greenland ice sheet model parameters constrained using simulations of the
858 Eemian Interglacial, *Clim. Past*, 7, 381–396, <https://doi.org/10.5194/cp-7-381-2011>, 2011.

859 Rohling, E. J., Hibbert, F. D., Williams, F. H., Grant, K. M., Marino, G., Foster, G. L., Hennekam, R., de Lange, G. J., Roberts,
860 A. P., Yu, J., Webster, J. M., and Yokoyama, Y.: Differences between the last two glacial maxima and implications for ice-
861 sheet, $\delta^{18}\text{O}$, and sea-level reconstructions, *Quaternary Sci. Rev.*, 176, 1–28, <https://doi.org/10.1016/j.quascirev.2017.09.009>,
862 2017.

863 Rutt, I. C., Hagdorn, M., Hulton, N. R. J., and Payne, A. J.: The Glimmer community ice sheet model, *J. Geophys. Res. Earth*
864 *Surf.*, 114, 2004, <https://doi.org/10.1029/2008JF001015>, 2009.

865 Sellevold, R., van Kampenhout, L., Lenaerts, J. T. M., Noël, B., Lipscomb, W. H., and Vizcaino, M.: Surface mass balance
866 downscaling through elevation classes in an Earth system model: application to the Greenland ice sheet, *Cryosphere*, 13, 3193–
867 3208, <https://doi.org/10.5194/tc-13-3193-2019>, 2019.

868 Sherriff-Tadano, S., Abe-Ouchi, A., Yoshimori, M., Oka, A., and Chan W-L.: Influence of glacial ice sheets on the Atlantic
869 meridional overturning circulation through surface wind change, *Clim. Dyn.*, 50, 2881–2903, [https://doi.org/10.1007/s00382-
870 017-3780-0](https://doi.org/10.1007/s00382-017-3780-0), 2018.

871 Sherriff-Tadano, S., Abe-Ouchi, A., and Oka, A.: Impact of mid-glacial ice sheets on deep ocean circulation and global climate,
872 *Clim. Past*, 17, 95–110, <https://doi.org/10.5194/cp-17-95-2021>, 2021.

873 Sherriff-Tadano, S., Ivanovic, R., Gregoire, L., Lang, C., Gandy, N., Gregory, J., Edwards, T. L., Pollard, O., and Smith, R.
874 S.: Large ensemble simulations of the North American and Greenland ice sheets at the Last Glacial Maximum with a coupled
875 atmospheric general circulation-ice sheet model, *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2023-2082>, 2023.

876 Smith, R. N. B.: A scheme for predicting layer clouds and their water content in a general circulation model, *Q. J. R. Meteorol.*
877 *Soc.*, 116, 435-460, <https://doi.org/10.1002/qj.49711649210>, 1990.

878 Smith, R.S., and Gregory, J.: The last glacial cycle: transient simulations with an AOGCM, *Clim. Dyn.*, 38, 1545–1559,
879 <https://doi.org/10.1007/s00382-011-1283-y>, 2012.

880 Smith, R. S., Gregory, J. M., and Osprey, A.: A description of the FAMOUS (version XDBUA) climate model and control
881 run, *Geosci. Model Dev.*, 1, 53–68, <https://doi.org/10.5194/gmd-1-53-2008>, 2008.

882 Smith, R. S., George, S., and Gregory, J. M.: FAMOUS version xotzt (FAMOUS-ice): A general circulation model (GCM)
883 capable of energy- And water-conserving coupling to an ice sheet model, *Geosci. Model Dev.*, 14, 5769–5787,
884 <https://doi.org/10.5194/GMD-14-5769-2021>, 2021.

885 Snoll, B., Ivanovic, R. F., Valdes, P. J., Maycock, A. C., and Gregoire, L., J.: Effect of orographic gravity wave drag on
886 Northern Hemisphere climate in transient simulations of the last deglaciation, *Clim. Dyn.*, 59, 2067-2079,
887 <https://doi.org/10.1007/s00382-022-06196-2>, 2022.

888 Stokes, C. R., Tarasov, L., and Dyke, A. S.: Dynamics of the North American Ice Sheet Complex during its inception and
889 build-up to the Last Glacial Maximum, *Quaternary Sci. Rev.*, 50, 86-104, <https://doi.org/10.1016/j.quascirev.2012.07.009>,
890 2012.

891 Stone, E.J., and Lunt, D.J.: The role of vegetation feedbacks on Greenland glaciation, *Clim. Dyn.*, 40, 2671–2686,
892 <https://doi.org/10.1007/s00382-012-1390-4>, 2013.

893 Stone, E. J., Lunt, D. J., Annan, J. D., and Hargreaves, J. C.: Quantification of the Greenland ice sheet contribution to Last
894 Interglacial sea level rise, *Clim. Past*, 9, 621–639, <https://doi.org/10.5194/cp-9-621-2013>, 2013.

895 Svendsen, J. I., Alexanderson, H., Astakhov, V. I., Demidov, I., Dowdeswell, J. A., Funder, S., Gataullin, V., Henriksen, M.,
896 Hjort, C., Houmark-Nielsen, M., Hubberten, H. W., Ingólfsson, Ó., Jakobsson, M., Kjær, K. H., Larsen, E., Lokrantz, H.,
897 Lunkka, J. P., Lyså, A., Mangerud, J., ... Stein, R.: Late Quaternary ice sheet history of northern Eurasia, *Quaternary Sci.*
898 *Rev.*, 23, 1229–1271, <https://doi.org/https://doi.org/10.1016/j.quascirev.2003.12.008>, 2004.

899 Tarasov, L., and Peltier, W. R.: Greenland glacial history and local geodynamic consequences, *Geophys. J. Int.*, 150, 198-229,
900 <https://doi.org/10.1046/j.1365-246X.2002.01702.x>, 2002.

901 Tarasov, L., and Peltier, W. R.: A geophysically constrained large ensemble analysis of the deglacial history of the North
902 American ice-sheet complex, *Quaternary Sci. Rev.*, 23, 359-388, <https://doi.org/10.1016/j.quascirev.2003.08.004>, 2004.

903 Tarasov, L., Dyke, A. S., Neal, R. M., and Peltier, W. R.: A data-calibrated distribution of deglacial chronologies for the North
904 American ice complex from glaciological modelling, *Earth Planet. Sci. Lett.*, 315–316, 30–40,
905 <https://doi.org/10.1016/j.epsl.2011.09.010>, 2012.

906 Tierney, J. E., Zhu, J., King, J., Malevich, S. B., Hakim, G., and Poulsen, C.: Last Glacial Maximum SST proxy collection and
907 data assimilation, PANGAEA [dataset], <https://doi.org/10.1594/PANGAEA.920596>, 2020.

908 Timmermann, A., Knies, J., Timm, O. E., Abe-Ouchi, A., and Friedrich, T.: Promotion of glacial ice sheet buildup 60–115 kyr
909 B.P. by precessionally paced Northern Hemispheric meltwater pulses, *Paleoceanography*, 25, PA4208,
910 doi:10.1029/2010PA001933, 2010.

911 Ullman, D. J., Legrande, A. N., Carlson, A. E., Anslow, F. S., and Licciardi, J. M.: Assessing the impact of Laurentide ice
912 sheet topography on glacial climate, *Clim. Past*, 10, 487–507, <https://doi.org/10.5194/CP-10-487-2014>, 2014

913 Valdes, P. J., Armstrong, E., Badger, M. P. S., Bradshaw, C. D., Bragg, F., Crucifix, M., Davies-Barnard, T., Day, J. J.,
914 Farnsworth, A., Gordon, C., Hopcroft, P. O., Kennedy, A. T., Lord, N. S., Lunt, D. J., Marzocchi, A., Parry, L. M., Pope, V.,
915 Roberts, W. H. G., Stone, E. J., Tourte, G. J. L., and Williams, J. H. T.: The BRIDGE HadCM3 family of climate models:
916 HadCM3@Bristol v1.0, *Geosci. Model Dev.*, 10, 3715–3743, <https://doi.org/10.5194/gmd-10-3715-2017>, 2017.

917 Vizcaíno, M., Lipscomb, W. H., Sacks, W. J., van Angelen, J. H., Wouters, B., and van den Broeke, M. R.: Greenland Surface
918 Mass Balance as Simulated by the Community Earth System Model. Part I: Model Evaluation and 1850–2005 Results, *J.*
919 *Clim.*, 26, 7793–7812, <https://doi.org/10.1175/JCLI-D-12-00615.1>, 2013.

920 Waelbroeck, C., Labeyrie, L., Michel, E., Duplessy, J. C., McManus, J. F., Lambeck, K., Balbon, E., and Labracherie, M.:
921 Sea-level and deep water temperature changes derived from benthic foraminifera isotopic records, *Quaternary Sci. Rev.*, 21,
922 295–305, [https://doi.org/10.1016/S0277-3791\(01\)00101-9](https://doi.org/10.1016/S0277-3791(01)00101-9), 2002.

923 Willeit, M. and Ganopolski, A.: The importance of snow albedo for ice sheet evolution over the last glacial cycle, *Clim. Past*,
924 14, 697–707, <https://doi.org/10.5194/cp-14-697-2018>, 2018.

925 Willeit, M., Calov, R., Talento, S., Greve, R., Bernales, J., Klemann, V., Bagge, M., and Ganopolski, A.: Glacial inception
926 through rapid ice area increase driven by albedo and vegetation feedbacks, *Clim. Past*, 20, 597–623, <https://doi.org/10.5194/cp-20-597-2024>, 2024.

928 Williams, J. H. T., Smith, R. S., Valdes, P. J., Booth, B. B. B., and Osprey, A.: Optimising the FAMOUS climate model:
929 inclusion of global carbon cycling, *Geosci. Model Dev.*, 6, 141–160, <https://doi.org/10.5194/gmd-6-141-2013>, 2013.

930 Williamson, D.: Exploratory ensemble designs for environmental models using k-extended Latin Hypercubes, *Environmetrics*,
931 26, 268–283, <https://doi.org/10.1002/env.2335>, 2015.

932 Ziemen, F. A., Rodehacke, C. B., and Mikolajewicz, U.: Coupled ice sheet–climate modeling under glacial and pre-industrial
933 boundary conditions, *Clim. Past*, 10, 1817–1836, <https://doi.org/10.5194/cp-10-1817-2014>, 2014.

934 Zweck, C., and Huybrechts, P.: Modeling of the northern hemisphere ice sheets during the last glacial cycle and glaciological
935 sensitivity, *J. Geophys. Res.*, 110, D07103, doi:10.1029/2004JD005489, 2005.