

Impact of the Late Miocene Cooling on the loss of coral reefs in the Central Indo-Pacific

Benjamin Petrick¹, Lars Reuning¹, Miriam Pfeiffer¹, Gerald Auer², Lorenz Schwark¹

5 ¹ Institute of Geosciences, Kiel University, Kiel, Germany

² University of Graz, Graz, Austria

Correspondence to: Benjamin Petrick (benjamin.petrick@ifg.uni-kiel.de)

Abstract. The Late Miocene Cooling (LMC) has been recognized as a global event in the climate record and posited as the start of modern ecosystems. Whereas [climate](#) shifts in modern tropical terrestrial ecosystems around 7.0 - 5.4 Ma are known, little is known about the impact of the [LMC](#) on coral reefs, where few good proxy records exist. During the Pliocene, a stratigraphic interval is present in the Central Indo-Pacific, where reefs that were present at the start of the Messinian disappeared by the Early Pliocene. This “Pliocene [Reef Gap](#)” has often been ascribed to non-climatic factors. However, a lack of proxy data prevents an understanding of climatic changes during this time. Here, we present a TEX₈₆^H-based sea surface temperature (SST) record for the Coral Sea, suggesting that the LMC is present across the Central Indo-Pacific. During the LMC, SST at ODP Site 811 declined by about 2°C, and cooling lasted from 7.0 Ma to 5.4 Ma. This cooling has also been seen in other parts of the Central Indo-Pacific. The LMC caused many changes in the Central Indo-Pacific, including a southwest shift of the monsoon belt, changes in terrestrial inputs, and the strength of ocean currents. All these factors can be stressors affecting coral reef growth. This suggests the overall impact of the LMC was to increase the stress on reef systems, which could have provided a driver for the collapse of individual reefs and, therefore, a potential cause for the “Pliocene [Reef Gap](#).” The change in SST and other stressors associated with the cooling caused coral reef systems to collapse across the Central Indo-Pacific.

1. Introduction

1.1 LMC background

25 Global Sea Surface Temperature (SST) records have identified the Late Miocene Cooling (LMC) as a worldwide event occurring between 7.0- 5.4 Ma when SST globally decreased by about 6 °C (Herbert et al., 2016; Holbourn et al., 2018; Martinot et al., 2022; Tanner et al., 2020; Wen et al., 2023). However, the LMC does not occur in benthic δ¹⁸O stacks or splices (Westerhold et al., 2020; Zachos et al., 1994). It is a major climatic cooling that does not seem to be associated with any changes in ice volume or deep water temperatures (Herbert et al., 2016; Martinot et al., 2022; Tanner et al., 2020). Therefore, it has often been overlooked as a low-latitude climatic factor impacting tropical marine ecosystems.

30

Deleted: cooling

Deleted: reef

Deleted: gap

Deleted: Coral

35 The LMC has been linked to aridification in Asia and Africa due to changes in the monsoonal system (Dupont et al.,
2013; Feakins, 2013; Wen et al., 2023). This resulted in significant shifts from C₃ to C₄ plants in tropical zones, indicating the
expansion of tropical grasslands (Huang et al., 2007; Steinthorsdottir et al., 2021; Strömberg and Strömberg, 2011). Due to
grassland expansion in East Africa, it is thought that some of the earliest hominids began to evolve around 7 Ma to adapt to
40 the changed climatic conditions (Brunet, 2020). In the ocean, a biogenic bloom, marked by an increase in the $\delta^{13}\text{C}$ of benthic
foraminifera occurred just before the onset of the LMC at around 8 Ma (Diester-Haass et al., 2004; Drury et al., 2018; Lübbers
et al., 2019).

1.2 Causes of the LMC

45 The causes of the LMC are not well understood. The paradox of a major cooling without a concurrent increase in
glaciation has still not been fully explained. There are a series of short-term glacials, but these only occur after 6 Ma, almost
1 Mya after the onset of the cooling, and at the start of the warming following the LMC (Jöhneck et al., 2020). Therefore,
whatever triggered the LMC must have either had little impact on the size of glaciers or deep-water temperatures. There are
two primary explanations for the cooling associated with the LMC. These are gateway changes causing shifts in ocean
circulation and changes in atmospheric CO₂.

50 Ongoing re-organization in the Indonesian Throughflow (Hall, 2002, 2009; Gallagher et al., 2024) and the Isthmus
of Panama (Collins et al., 1996; Haug et al., 2001) may offer a tectonic mechanism for the LCM due to the re-organization of
the global thermohaline circulation. However, almost all known major thresholds in these systems date to the Pliocene or later
(Auer et al., 2019; Haug et al., 2001; De Vleeschouwer et al., 2018, 2019), making these tectonic restrictions of oceanic
gateways an unlikely cause. Furthermore, available SST data shows that changes observed during the LMC eventually reversed
55 after 5.4 Ma. This temporary and reversible pattern further emphasizes that long-term and permanent tectonic, oceanic gateway
closure could not have been the primary driver of the LMC.

An alternative hypothesis is that there was a reduction of atmospheric $p\text{CO}_2$ during the LMC. Lower $p\text{CO}_2$ would
explain the difference between the increased tropical presence of C₄ grasses, which were better adapted to lower atmospheric
CO₂ concentrations (Herbert et al., 2016; Wen et al., 2023). Model estimates, including atmospheric CO₂ reductions, also fit
60 well with temperature reconstructions, showing a cooling of about 2-3°C in the central Indo-Pacific (Martinot et al., 2022).
The cause of the Late Miocene CO₂ reduction is not well understood. One suggestion is that it was related to higher
oceanographic productivity, which would likely draw down atmospheric CO₂ (Holbourn et al., 2018). This could result from
the Biogenic Bloom at 8-7 Ma (Diester-Haass et al., 2004; Grant and Dickens, 2002). Another theory is that changes in plate
motion during the LMC led to a reduction of CO₂ input to the atmosphere (Herbert et al., 2022). Finally, it has been suggested
65 that the tectonic uplift of Papua New Guinea led to a long-term decrease in atmospheric CO₂ (Martin et al., 2023; Clift et al.,
2024).

Deleted: “

Deleted: ”

Deleted: lived

Deleted: ,

Deleted: and

Deleted: that

Deleted: which would draw down atmospheric

Deleted: ,

Deleted: be a consequence of

Deleted: “b

Deleted: b

Deleted: ”

Deleted: between

Deleted: ,

1.3 Miocene reef history

85 There was an extensive coral reef system on the Queensland Plateau during the Early to Middle Miocene (Feary et al., 1991; Betzler and Chaproniere, 1993). After 11 Ma, however, the coral reefs appear to have retreated (Isern et al., 1993, 1996). Benthic foraminiferal reconstructions of sea level show a gradual increase in relative water depth due to increasing subsidence of the Queensland Plateau between 13-8 Ma (Katz and Miller 1993). This roughly corresponds to the timing of the so-called Carbonate Crash, a period of low carbonate accumulation in the deep sea of all major tropical ocean basins (Lübbbers et al., 2019). It was hypothesized that this episode also reduced carbonate accumulation rates in reefs in the South China Sea (Mathew et al., 2020). A similar mechanism could have made the reefs on the Queensland Plateau more susceptible to subsidence-related drowning and later collapse due to climatic changes. Other reefs in the Coral Sea and surrounding areas show evidence of a later collapse, many occurring between 7.0-5.4 Ma during the LMC. For instance, in the southern Coral Sea, coral reefs on Marion Plateau disappeared between 7.5-5.7 Ma (Bashah et al., 2024; Eberli et al., 2010; Ehrenberg et al., 2006). On the other side of Australia, the NW Shelf barrier reef experienced a major drowning event between 7.2-5.9 Ma (Belde et al., 2017; Rosleff-Soerensen et al., 2012, 2016). Finally, studies of Early Pliocene sediments on the Queensland Plateau show hardgrounds on the shallow carbonate platforms, suggesting a complete absence of coral reefs during the 'Pliocene Reef Gap' (Droxler et al., 1993).

100 Globally, similar reef trends are seen. The highest abundance of coral reefs occurs in the Middle Miocene, followed by a slight decline towards the Late Miocene (Harrison et al., 2023; Perrin and Kiessling, 2012). However, by the Early Pliocene, there appears to have been a loss of coral reefs. This suggests that the major reef loss occurred between the beginning of the Messinian and the beginning of the Early Pliocene (7.25-5.33 Ma). Harrison et al. (2023) attribute reef loss in the central Indo-Pacific to several factors impacting individual reefs differently (Fig. 2). These include tectonic processes, sea level changes, and increases in terrestrial input. The authors reject climate change as a causal factor, arguing that SSTs during the Late Miocene were similar to modern ones, and that there is no evidence of major warming across this time (Harrison et al., 2023). However, there is evidence that the tropical water belt cooled and contracted during the LMC (Martinot et al., 2022; Liu et al., 2022). Moreover, there were changes in the global distribution of reefs during this time, including a decrease in their latitudinal range (Perrin and Kiessling, 2012).

1.4 Tropical SST change during the LMC

110 As discussed above, the climatic impact of the LMC has not been investigated as a driver of the "Pliocene Reef Gap" because many previous studies of benthic $\delta^{18}\text{O}$ isotope records did not show an increase of glaciation during the LMC (Harrison et al., 2023; Perrin and Kiessling, 2012). Furthermore, even after the LMC was identified, many early tropical records showed less than 1 °C cooling during the LMC (Herbert et al., 2016). As a result, it was concluded that the temperature impact of the LMC in the tropics was small. However, this conclusion has been challenged as most LMC records are based on U^{37} derived SST reconstructions (Herbert et al., 2016). These alkenone-based SST records have a saturation limit of 28 to 29 °C (Müller et al., 1998) when the proportion of the $\text{C}_{37:3}$ isomer used for the temperature calculation approaches zero (Grimalt et al., 2001).

Deleted: -

Deleted: Large b

Deleted: between

Deleted: t

Deleted: ance

Deleted: led to

Deleted: potentially could have made the reefs on the Queensland Plateau more susceptible to subsidence

Deleted: There is evidence of a reduction in carbonate platform area reduction due to stress caused by the decrease in carbonate production due to the Carbonate Crash from 13-8 Ma (Lübbbers et al., 2019; Mathew et al., 2020). This could have made the reefs more vulnerable to later collapse due to climatic changes....

Deleted: much

Deleted: reef

Deleted: gap'

Deleted:

Deleted: Mid

Deleted: -

Deleted: a

Deleted:

Deleted: by the Early Lower Early Pliocene, there appears to have been...

Deleted: to have caused

Deleted: major

Deleted: end

Deleted: -

Deleted: are

Deleted: ,

Deleted: proxy

Especially in carbonate rocks with low organic content and poor organic matter preservation, $U^{K_{37}}$ SSTs are considered unreliable for temperatures exceeding 26-27 °C (Pelejero and Calvo, 2003; Grimalt et al., 2001). Models show that many parts of the tropical and sub-tropical Miocene Ocean were too warm for applying the $U^{K_{37}}$ proxy (Burls et al., 2021). In fact, except ODP Site 722, all the sites where the $U^{K_{37}}$ proxy was used to reconstruct SSTs are located in the cold tongue originating from the East Pacific Equatorial Upwelling Zone (Herbert et al., 2016). ODP Site 722 is located in the Arabian Sea upwelling cell and is cooled by coastal upwelling driven by monsoonal winds (Bialik et al., 2020).

Mg/Ca and TEX_{86} records show a stronger cooling during the LMC. For instance, the West Pacific Warm Pool (WPWP) stack of TEX_{86} SSTs shows a cooling of 2 °C during the LMC (Liu et al., 2022). Furthermore, between 7-5 Ma, all the sites included in the WPWP stack show a cooling of 2-3 °C, suggesting a cooling of the entire WPWP (Liu et al., 2022; Zhang et al., 2014). This cooling is also confirmed by a Mg/Ca SST record from ODP Site 1146, one of the sites included in the WPWP TEX_{86} SST stack. In agreement with the TEX_{86} SSTs, the Mg/Ca SST shows a cooling of 2 °C (Holbourn et al., 2018). In the Indian Ocean, an Mg/Ca record from IODP Site 1443 shows a cooling of about 2 °C (Martinot et al., 2022a). However, these records mainly derive from the northern margins of the central Indo-Pacific warm water area. Therefore, while the magnitude of cooling and timing of the LMC in these records is similar, it is unclear whether this was a large-scale, tropical-wide event that could have affected coral reefs across the Central Indo-Pacific.

1.5 Objective


To better understand the link between tropical SSTs and the LMC, new records need to be produced from across the central Indo-Pacific. In this study, we present a TEX_{86}^H SST record covering the LMC from ODP Site 811, located on the Queensland Plateau in the Coral Sea (Fig. 1). This is an extension of a record previously published in Petrick et al. (2023) that spanned the period from 6.6-11.1 Ma. In this study, we added new TEX_{86}^H SST data between 2.5 and 6.6 Ma to investigate the LMC and the Pliocene. We chose this site because the Coral Sea has one of the highest coral reef densities in the world (Bridge et al., 2019) and is bordered by the Great Barrier Reef off the East Australian coast. The central Indo-Pacific is a hotspot of coral diversity during the Miocene and, therefore, a key reef area (Renema et al., 2008). There is abundant evidence of reef loss during the Late Miocene in the Coral Sea (Bashah et al., 2024; Betzler et al., 2024). This paper investigates whether climate change during the LMC could have led to the collapse of reef systems across the central Indo-Pacific and globally.

Deleted: above

Deleted: of

Deleted: higher than

Deleted: period



180 **Figure 1:** Map showing the central Indo-Pacific and the location of ODP Site 811 studied here (red star). All the sites included in the
 185 **TEX₈₆-derived West Pacific Warm Pool stack** (Liu et al., 2022) are indicated with black triangles (Fig 6). All Mg/Ca records are
 190 shown with black dots (Fig 5). The base map is from Ocean Data View (Schlitzer, 2021).

2 Methods

185 2.1 Biogeochemistry

We solvent-extracted 50 cc of sediment for this project, which resulted in between 50 and 60 g of sediment, suitable for
 190 obtaining sufficient lipids for TEX₈₆ determination. Dried and homogenized samples were Soxhlet extracted for 48 h using a
 solvent mixture of DCM: MeOH (9:1, v/v). The addition of activated copper turnings removed elemental sulfur. A Büchi
 solvent evaporator reduced excess solvent to a final volume of 2 ml. Samples were then transferred into a 4 ml vial, where the
 total extract (TE) was taken to dryness under a gentle stream of nitrogen. TEs were fractionated into aliphatic, aromatic, and
 polar fractions by silica gel-column chromatography (6 ml SPE column, 2.8 g Silica 60 mesh, 25–40 µm) using solvents with
 increasing polarity in an LC-TECH automated SPE system. NSO (polar) compounds were eluted with 14 ml DCM/ MeOH
 (1:1, v/v). The polar fraction was reconstituted in hexane/isopropanol (9:1, v/v) and re-chromatographed over aminopropyl-
 substituted silica gel (3 ml SPE column, 1.0 g aminopropyl-silica, 25–40 µm). The alcohol fraction containing the GDGTs was

Deleted: extracting

Deleted: organic material

eluted with 5 ml of hexane/isopropanol (9:1, v/v) and, after drying, was re-dissolved in hexane/isopropanol (99:1, v/v) to a final concentration of 6 mg/ml for injection into the HPLC/MS system.

GDGTs were measured on an AGILENT liquid chromatograph coupled to an AGILENT single quadrupole mass spectrometer following the analytical protocol of Hopmans et al. (2016). The HPLC instrument was equipped with an AGILENT HILIC silica column (2.1 x 150 mm; 1.5 μ m particle size) and a guard column maintained at 30°C. Detection of archaeological core lipids was achieved by single ion recording of their protonated molecular ions [M + H⁺], and compounds were quantified by integration of peak areas using AGILENT Masshunter© software. Calculation of TEX₈₆^H followed Kim et al. (2010). Reproducibility upon duplicate measurements showed a relative standard error of <2%.

3 Results


3.1 Site details

ODP Site 811 is located in the Coral Sea (16.516° S, 148.157 E) and has a modern water depth of 937.0 m (Fig. 1).

Benthic foraminifera data shows that the site was probably shallower during the start of the Late Miocene (<500 m) and that the site only reached its current depth around the Early Pliocene (Katz and Miller 1993). For this study, we focused on the record from Hole A between 113 and 27 m. The age model from this site was created using a new nannofossil stratigraphic study. This, including the data and error, was published by Petrick et al. (2023). The studied interval consists almost entirely of nannofossil to foraminiferal ooze. In the upper 70 mbsf, the pelagic components are mixed with fine, shallow water bank-derived particles. The sediment was interpreted as periplatform ooze (Fig. 2). One debris flow occurs close to the top of the studied interval (811A-4H-6), which was avoided during sampling (Fig. 2). The interval between 70 and 113 mbsf is characterized by purely pelagic sediments (Fig. 2). The carbonate content in the studied section always exceeds 90 wt. %. None of these lithologic boundaries match changes in the TEX₈₆ data or the timing of the LMC (Fig 2). The modern annual average SST at Site 811 is 26.1 °C (Locarnini et al., 2019). The average summer temperature is 28.32 °C, and the average winter temperature is 25.09 °C (Locarnini et al., 2019).

Deleted: Large b


Deleted: foraminifer



225 **Figure 2:** Lithologic column for ODP Hole 811A. The descriptions are from (Shipboard Scientific Party, 1991). Age model data from Petrick et al. (2023). Carbonate content data is from Davies et al. (1991). The LMC is marked on the figure.

3.2 TEX₈₆ tests

230 The new TEX₈₆^H record for 1.9-6.6 Ma complements previous data covering 6.6 to 11.1 Ma (Petrick et al., 2023). The data for this study and data from Petrick et al. (2023) are presented in supplemental data 1. Several tests were performed to evaluate the new data and ensure that it reflects SST changes and is not the result of nonthermal factors impacting GDGTs.



Deleted: a

Deleted: a

Deleted:

With only a few exceptions, the TEX₈₆^H data passed all applied tests. We will discuss the various tests, what they show, and why we decided to remove (or retain) data points.

The Methane Index (MI) excludes any data affected by gas-hydrate-related anaerobic [methane oxidation](#) (Zhang et al., 2011). The MI values are below the 0.5 value [recommended](#) for rejection (Fig. 3a). We also used the GDGT0% index to eliminate samples with GDGTs substantially originating from sedimentary archaeal methanogenesis (Weijers et al., 2006; Sinninghe Damsté et al., 2012). The values were well below the 67% cut-off for excessive methanogenesis (Fig. 3b). We used the ring index (RI) to evaluate whether the GDGTs deviate from modern values (Zhang et al., 2016). All our data fell within the acceptable error envelope of 0.3 (Fig. 3d). Finally, we used the 2/3 index to ensure that the GDGTs were being formed near the surface (Taylor et al., 2013; Hernández-Sánchez et al., 2014). This measures the relationship between the compounds GDGT 2 and GDGT 3, which is related to the depth of production of [archaeal GDGTs](#). While the appropriate cut-off for this test is still being debated, the data values are low (< 5) and indicate surface production (Rattanasriampaipong et al., 2022).

One GDGT index that did not fit recommendations for quality assurance was the BIT index (Fig. 3c). This index was developed to track the amount of terrigenous material that could interfere with the TEX₈₆ values via soil-sourced GDGTs (Schouten et al., 2013). The original cut-off point is 0.4, although there is a debate about whether that cut-off is too strict (Schouten et al., 2013) and how to evaluate variations in crenarchaeol shown to affect the BIT index (Fietz et al., 2011). Although within the GDGT suite analyzed here, 11 samples exceed a BIT value of 0.5 (Fig 3c), the data shows no covariance with either time, depth, or SST (Fig. S1). The highest BIT values neither match the highest nor lowest SST data. Finally, removing the high BIT samples does not affect the trends or conclusions of the paper. Therefore, we have decided to retain all data in the results based on the current debate regarding the reliability of the BIT index.

Deleted: oxidation of methane

Deleted: the

Deleted: proxy


Deleted: .

Deleted: T

Deleted: Supplementary

Deleted: ure

Deleted: major



265

Figure 3: Graphic illustration of the new ODP Site 811 data for various GDGT indices used for quality assurance for previously published data (Petrick et al., 2023). Quality criteria shown are a. Methane Index (MI, after Zhang et al., 2011), b. %GDGT-0-index (after Sinninghe Damsté et al., 2012), c. Branched vs. isoprenoid Index (BIT, proposed by Schouten et al., 2002), d. Ring Index (RI, proposed by Zhang et al., 2016). Quality assurance tests indicate that GDGT data are suitable for SST reconstruction and are not compromised by other environmental drivers.

270

3.3 SST trends

For this study, we follow the age model of Petrick et al. (2023), which updated the original shipboard (Davies et al., 1991) and a previously published age model (Isern et al., 1993, 1996) (Fig. 2). A part of the record presented in this study (6.6-11.1 Ma) was previously published in Petrick et al., (2023) and discussed in more detail in section 4.1 (Fig. 4). It is also important to note that the ODP Site 811 SST record has a lower resolution than other nearby records. This may mean that internal variability

275

- Deleted: r (
- Deleted:)
- Formatted: English (US), Not Highlight
- Deleted: (
- Deleted: ,
- Deleted:)
- Deleted: (
- Deleted:)
- Deleted: ¶

285 within the record could exaggerate the LMC cooling trend. Therefore, we used both a running average and a fixed time window
 for the LCM to quantify the amount of cooling in the record. However, the boundaries of the LMC are uncertain. We, therefore,
 estimated LMC cooling using two different definitions of the LMC (Table 1). First, we compared SSTs before the LMC (11.1-
 7.0 Ma) to the LMC, following Herbert et al. (2016), where the LMC is defined as 7.0-5.4 Ma. We then used the windows
 used in Martionot et al. (2022) for the pre-LMC (8.5-7.5 Ma) and maximum of the LMC (6.5-5.5 Ma). The data is summarized
 290 in Table 1. The average cooling at ODP Site 811 during the LMC is 2 +/- 0.2 °C across all windows. We will use 2 °C as our
 best estimate of LMC cooling at Site 811.

Deleted: height

Deleted: be using

Citation	Pre-LMC window	Average Pre-LMC SST at ODP Site 811	LMC window	LMC average SST C at ODP Site 811	Difference Pre-LMC minus LMC
Herbert et al., (2016)	11.1-7.0 Ma	30.0 °C	7.0-5.4 Ma	28.2 °C	1.8 °C
Martinot et al., (2022)	8.5-7.5 Ma	29.9 °C	6.5-5.5 Ma	27.9 °C	2.0°C

Table 1: different definitions of the LMC compared in this study, average SSTs, and LMC cooling at Site 811.

295 4 Discussion

4.1 Previous work at ODP 811

The Joides Resolution drilled ODP Site 811 as part of Expedition 133 in the Coral Sea (Davies et al., 1991) (Fig. 1).
 ODP Site 811 was located further south during the Late Miocene (likely ~ 19.4 to 17.5°S) (Van Hinsbergen et al., 2015). As
 part of the post-cruise work, a $\delta^{18}\text{O}$ record was produced using planktonic foraminifera, which showed SSTs were between
 300 18-24 °C during the Late Miocene (Isern et al., 1993, 1996). These cold temperatures were proposed to be the reason for the
 collapse of the coral reefs in the Coral Sea between 11-8 Ma (Isern et al., 1993, 1996). However, many similar foraminiferal
 $\delta^{18}\text{O}$ records showing cool tropical SSTs have been re-evaluated subsequently using biomarkers and well-preserved
 foraminifera. These studies show that post-depositional alteration of calcite is a larger source of error than originally thought
 (Nairn et al., 2021; Wilson et al., 2002). Petrick et al. (2023) used $\text{TEX}_{86}^{\text{H}}$ to reconstruct SSTs over the 8-12 Ma period when
 305 the reefs in the Coral Sea drowned (Isern et al., 1993) (Fig. 4). This record showed an average SST of 30 °C between 8-12 Ma.
 Petrick et al. (2023) concluded that a combination of low aragonite saturation and heat stress impaired coral growth, and that
relative sea-level rise combined with these stressors and potential changes in ocean circulation led to the drowning of the coral
 reefs (Petrick et al., 2023). Therefore, at ODP Site 811, the period of major reef loss occurred before the LMC (Isern et al.,
 1993, 1996). However, other major reef systems in the Coral Sea did drown between 7-5.4 Ma, including the Marion Plateau
 310 (Bashah et al., 2024). Therefore, while ODP Site 811 is not located in an area that experienced coral reef loss during the LMC,
 it can still be used to understand the climatic impact of the LMC on the Coral Sea and the wider central Indo-Pacific.

Deleted: a

Deleted: ,

Deleted: combined with changes in ocean circulation, and


Deleted: that

Deleted: regional

Deleted: sea

Deleted: ly

Deleted:



325 Figure 4: The $\text{TEX}_{86}^{\text{II}}$ -derived SST record for ODP Site 811 is shown in red, including data from 6.6 to 11.1 Ma taken from Petrick
 et al. (2023). The grey line represents an 8-point running average. The calibration error from Kim et al. (2010) is shown at the top.
 The sea level evolution is shown in blue after Miller et al. (2020). The relative abundance (%) of areas covered by reefs in the Central
 Indo-Pacific is indicated by black line and dots at the bottom, with data taken from Harrison et al. (2023). Note we have followed
 Harrison et al. (2023) figures by setting the Reef Abundance % for each faunal stage at the midpoint of the stage. The exception is
 330 the value of the Serravallian, which is shown at the end of the period (11.63 Ma) to fit it into the figure.

4.2 ODP Site 811 and the LMC

As shown above, the cooling at ODP Site 811 during the LMC is about 2 °C (Fig 4). This is consistent with other
 records from the Central Indo-Pacific, including ODP Site 1146 and IODP Site U1448 (Holbourn et al., 2018; Martinot et al.,
 335 2022) (Fig 5). This also matches the magnitude of cooling seen in TEX_{86} records from the WPWP (Liu et al., 2022b; Zhang et
 al., 2014b) (Fig 6). Therefore, the new ODP Site 811 data confirms previous findings (Holbourn et al., 2018; Liu et al., 2022;
 Martinot et al., 2022), showing that the cooling of the tropics was greater than the 0.5 °C during the LMC, as initially suggested

Deleted: shown

Deleted: ,

Deleted: putting

Deleted: originally

by Herbert et al., (2016). Furthermore, the data suggests that the cooling in the Central Indo-Pacific was uniform during the LMC.

345 Unfortunately, our records can only provide limited information regarding the causes of LMC. The ODP Site 811 temperature trends are consistent with the expected cooling inferred using climate modeling (Martinot et al., 2022), which estimates SSTs of around 28 °C and a cooling of about 2 °C at ODP Site 811 when atmospheric CO₂ is lowered to 280-ppm. Given that ODP Site 811 and IODP Site 1448 are both located in the Central Indo-Pacific warm pool but in different ocean basins and hemispheres, the fact that mean SSTs and LMC cooling are consistent with climate models could strengthen the argument that a change in atmospheric CO₂ caused the LMC. Therefore, our ODP Site 811 record suggests that changes in
350 $p\text{CO}_2$ may be responsible for the cooling seen in the record.

The major difference between the ODP Site 811 record and other records from the Central Indo-Pacific seems to be the post-cooling recovery of temperatures (Fig. 5,6). Not all sites return to pre-LMC temperatures. The timing of post-LMC recovery is also different between sites. ODP Site 811 remained cool after the LMC but fully recovered by the Early to Late Pliocene boundary (Fig. 4). Other records show a dramatic post-LMC recovery, such as IODP Site 1448 (Jöhnck et al., 2020)
355 between 5.5-5 Ma (Fig 5).

In contrast, the TEX₈₆ WPWP stack shows a long-term cooling trend and no clear recovery after the LMC, although there is warming around 4 Ma (Liu et al., 2022) (Fig. 6). Therefore, while the Central Indo-Pacific experienced the same 2-3 °C cooling during the LMC the recovery was heterogeneous (Figs 5,6). We conclude that if the “Pliocene Reef Gap” was triggered by climatic change, it was most likely the LMC cooling, not the post-LMC recovery.

360

Deleted: post

Deleted: therefore




Figure 5: ODP Site 811 $\text{TEX}_{86}^{\text{H}}$ -derived SSTs compared to the Mg/Ca-derived SST records. The gray bar marks the LMC as defined in Herbert et al., 2016. Sites are shown in Figure 1.

365


4.5 Potential stressors in the Central Indo-Pacific

As shown above, ODP Site 811 experienced a rapid and strong cooling associated with the LMC (Fig 2). Furthermore, SSTs did not recover until at least 5 Ma, well into the “Pliocene Reef Gap.” The cooling was widespread, as it was recorded at many sites in the Central Indo-Pacific. Therefore, it could have negatively impacted coral reefs and carbonate platform ecosystems. Studies show that globally, there was a reduction in the latitudinal extent of coral reefs during the Late Miocene, including the late LMC (Perrin and Kiessling, 2012). On the Marion Plateau, south of ODP Site 811, reefs disappeared around 7 Ma (Bashah et al., 2024; Ehrenberg et al., 2006; Isern et al., 2004). Today, an SST gradient of about 4 degrees exists between

370

Deleted: had negative impacts on

ODP Site 811 and the southern Marion Plateau (Locarnini et al., 2019). ODP Site 811 had average SSTs of about 28 °C and possibly as cold as 24 °C during the LMC. Therefore, SSTs could have been as cold as 20 °C on the southern Marion Plateau. Coral reefs have lower carbonate accumulation due to lower coral growth rates at these temperatures (Laugie et al., 2019; Lough, 2008; Lough and Barnes, 2000; Lough and Cantin, 2014). This can impair their ability to adapt to changing oceanographic conditions (Higuchi et al., 2015). Other key carbonate producers associated with reefs show a similar temperature dependence during the Miocene (Bassi et al., 2024). Porcelaneous larger benthic foraminifera (pLBF) diversity is at its lowest level during the Messinian (almost contemporaneous with the LMC), with only three species identified for the Central Indo-Pacific (Bassi et al., 2024). Since pLBF are a crucial contributor to carbonate production in Miocene coral reef environments, this decrease could have negatively impacted the accretion rate of the reef. So, the cold SSTs during the LMC could have negatively impacted many of these higher-latitude coral reefs and contributed to the latitudinal contraction of coral reefs (Perrin and Kiessling, 2012).



390 Figure 6: ODP Site 811 (red) temperature anomalies compared to the Tropical Anomaly (blue) from Herbert et al. (2016) and the WPWP stack (black) (Liu et al., 2022). SST data for this plot has been normalized to modern temperatures to better compare with the anomaly data presented by Herbert et al. (2016). This was done for ODP Site 811 and the WPWP stack by subtracting modern SSTs from the data. The gray bar delineates the LMC as defined by Herbert et al. (2016). Sites included in the WPWP stack are shown in Figure 1.

Deleted: also see experienced declines during the LMC

Deleted: LMC

Deleted: two

Deleted: modern

400 However, as discussed in Petrick et al. (2023), stressors other than SSTs can affect the coral reefs. A larger latitudinal
SST gradient during the LMC may have been associated with changes in the ocean circulation, as seen during other major
cooling events (Petrick et al., 2018; 2019). It has already been proposed that changes in the strength of ocean currents could
have caused greater erosion of carbonate platforms, possibly impacting the expansion of coral reefs (Betzler et al., 2024;
Betzler and Eberli, 2019). There is some evidence of stronger ocean currents during the LMC in the Central Indo-Pacific. On
405 the Marion Plateau, between 7.5 and 5.7 Ma, the intensification of the East Australian current caused an increase in drift
deposits (Bashah et al., 2024). On the NW Shelf, there is evidence of an intensification of NNE-SSW-oriented bottom currents
starting in the Late Miocene after 7.2 Ma (Thronberens et al., 2022), contributing to the demise of the regional reef system
(Belde et al., 2017). Therefore, a similar Central Indo-Pacific-wide strengthening of ocean currents could have occurred during
the LMC.

410 Changes in terrestrial input could also result from shifts in SSTs. There is evidence that SST cooling during the Late
Miocene led to southward shifts of the rain belts (Jöhnck et al., 2020). This may have increased rainfall and terrestrial input in
northern Australia and the Indonesian Archipelago (Jöhnck et al., 2020). Finally, in the Coral Sea, the weathering of Papua
New Guinea intensified, and there is evidence of increased terrestrial input to the northern Coral Sea from DSDP Sites 210
and 287 (Clift et al., 2024). During the Miocene, many coral reefs in the Indonesian Archipelago grew in turbid environments
415 (Santodomingo et al., 2016). However, increases in sediment input could have upset the delicate balance on which these corals
depended. Therefore, changing rainfall patterns during the LMC could have been an additional stressor for the coral reefs.

Therefore, the LMC was associated with major environmental changes that could have acted as stressors impacting
the coral reefs, causing widespread drowning. ~~Previous work in the southern Coral Sea has shown that similar changes in
420 temperature, terrestrial input, and ocean current led to changes in the benthic assemblage of carbonate producers (Hallock et
al., 2006). This modified the accretion potential of the carbonate platforms and contributed to their drowning in the Miocene
(Hallock et al., 2006). This is also similar to modern coral systems, where current temperature changes are accompanied by
multiple other environmental stressors, such as sea-level rise and increased sediment loads (Cornwall et al., 2021). While coral
reefs might be able to adapt to single stressors, such as higher sea levels, multiple stressors may add up synergistically and
trigger coral reef collapse (Darling and Côté 2013).~~

425 Finally, while the LMC cooling was unrelated to a sea level change, the global sea level increased after 5 Ma (Miller
et al., 2020) (Fig. 4). As a result, the carbonate platform tops were no longer in the photic zone when SSTs eventually returned
to the Middle Miocene levels during the Middle Pliocene. As a result, the corals could not grow back. It was noted that the re-
establishment of coral reefs in the Coral Sea was linked to a global sea level drop of ~2.9 Ma, which would have brought the
platforms back into the photic zone, allowing coral reefs to develop again (Droxler et al., 1993) (Fig 4). Therefore, it is likely
430 that after the drowning during the LMC, the reestablishment of the reefs was more related to changes in sea level than SST.

5 Conclusions

Deleted: late

Deleted: demonstrated

Deleted: the dominant carbonate producers due to many of the external forcings, including

Deleted: changes in

Deleted: ,

Deleted: of

Deleted: was a factor in the

Deleted: eventual

Deleted: of carbonate platforms

Deleted: (Hallock et al., 2006).

Deleted: Mid-

Deleted: -

The new TEX₈₆^H-derived SST data at ODP Site 811 shows that the LMC led to a relatively rapid drop in SSTs by about 2 °C in the southern part of the Central Indo-Pacific reef province. This is consistent with other records from the Central Indo-Pacific, which show a 2-3 °C drop in SSTs [c. 7-6 Ma ago](#). This decline in tropical SST preceded the “Pliocene Reef Gap” and could have been a stress factor contributing to the loss of coral reefs in the Central Indo-Pacific reef province. It is probably responsible for the loss of reefs on the Southern Marion Plateau. Additionally, there is evidence that the changes in global and regional SSTs caused by the LMC led to shifts in ocean currents, as well as rain belts. Together with the cooling of SST, these multiple stressors may explain some of the changes seen previously in individual [Upper Miocene reefs](#) and provide an overall explanation for the region-wide coral reef decline over such a short time interval. This likely explains the massive reduction in the extent of coral reefs in the “[Pliocene Reef Gap](#).” Therefore, this study emphasizes the detrimental impacts of climate change on coral reefs and the importance of limiting additional stressors, such as pollution, on reef ecosystems during global temperature change.

Data availability

The data for this paper is available both in supplements and at [the EU Open Research Repository Zenodo \(DOI: 10.5281/zenodo.10902264\)](#).

Author Contribution

All authors approved the manuscript and agreed to its submission. The corresponding author is B.P. All authors discussed the results and provided significant input to the final version of the manuscript. B.P. and L.R. designed the study. B.P. ran the project and processed the samples. L.S. performed the biomarker analysis in his lab and interpreted data with B.P. G.A., who provided a new-age model for the site. B.P., L.S., L.R., M.P., and G.A. provided vital feedback on the article.

Competing Interests

The authors declare they have no competing interests in this paper.

Acknowledgments

The authors would like to thank the participants of ODP Expedition 133 for recovering the core and providing the initial data and many of the studies used to reconstruct conditions in the Coral Sea. This project was funded with a grant from the Deutsche Forschungsgemeinschaft (DFG)—Project No. 447611930 [awarded to Benjamin Petrick, and contributes towards Austrian Science Fund \(grant no. P 36046-N\)P-](#). We thank the IODP core repository at Kochi for providing the samples. IODP has provided [critical](#) support for this project. We thank Erin McClymont, Andre Droxler, and Christian Betzler for their help and feedback on the paper. Finally, thanks to [Laurie Menviel and three](#) anonymous reviewers who helped improve the paper. All data needed to evaluate the conclusions in the paper are presented in the paper.

Deleted: between about

Deleted:

Deleted: late

Deleted: ‘

Deleted: R

Deleted: ’

Deleted: ary data one

Deleted: with a doi of

Deleted:

Deleted: for

Deleted: important

Deleted: two

References

- Auer, G., De Vleeschouwer, D., Smith, R. A., Bogus, K., Groeneveld, J., Grunert, P., Castañeda, I. S., Petrick, B., Christensen, B., Fulthorpe, C., Gallagher, S. J., and Henderiks, J.: Timing and Pacing of Indonesian Throughflow Restriction and Its Connection to Late Pliocene Climate Shifts, *Paleoceanogr Paleoclimatol*, 34, 635–657, 495 <https://doi.org/10.1029/2018PA003512>, 2019.
- Bashah, S., Eberli, G. P., and Anselmetti, F. S.: Archive for the East Australian current: carbonate contourite depositional system on the Marion Plateau, Northeast Australia, *Mar Geol*, 107224, <https://doi.org/https://doi.org/>, 2024.
- 500 [Bassi, D., Braga, J. C., Pignatti, J., Fujita, K., Nebelsick, J. H., Renema, W., and Iryu, Y.: Porcelaneous larger foraminiferal responses to Oligocene–Miocene global changes, *Palaeogeogr Palaeoclimatol Palaeoecol*, 634, 111916, <https://doi.org/10.1016/j.palaeo.2023.111916>, 2024.](https://doi.org/10.1016/j.palaeo.2023.111916)
- Belde, J., Back, S., Bourget, J., and Reuning, L.: Oligocene and Miocene Carbonate Platform Development In the Browse 505 Basin, Australian Northwest Shelf, *Journal of Sedimentary Research*, 87, 795–816, <https://doi.org/10.2110/jsr.2017.44>, 2017.
- Bellworthy, J. and Fine, M.: Warming resistant corals from the Gulf of Aqaba live close to their cold-water bleaching threshold, *PeerJ*, 9, e11100, <https://doi.org/10.7717/peerj.11100>, 2021.
- 510 Betzler, C. and Chaproniere, G. C. H.: Paleogene and Neogene Larger Foraminifers from the Queensland Plateau: Biostratigraphy and Environmental Significance, *Proceedings of the Ocean Drilling Program, 133 Scientific Results*, <https://doi.org/10.2973/ODP.PROC.SR.133.210.1993>, 1993.
- Betzler, C. and Eberli, G. P.: Miocene start of modern carbonate platforms, *Geology*, 47, 771–775, 515 <https://doi.org/10.1130/G45994.1>, 2019.
- Betzler, C., Brachert, T. C., and Kroon, D.: Role of climate in partial drowning of the Queensland Plateau carbonate platform (northeastern Australia), *Mar Geol*, 123, 11–32, [https://doi.org/10.1016/0025-3227\(95\)80002-S](https://doi.org/10.1016/0025-3227(95)80002-S), 1995.
- 520 Betzler, C., Hübscher, C., Lindhorst, S., Lüdmann, T., Hincke, C., Beaman, R. J., and Webster, J. M.: Seismic stratigraphic and sedimentary record of a partial carbonate platform drowning, Queensland Plateau, north-east Australia, *Mar Geol*, 470, 107255, <https://doi.org/https://doi.org/10.1016/j.margeo.2024.107255>, 2024.

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Black

- 525 Bialik, O. M., Auer, G., Ogawa, N. O., Kroon, D., Waldmann, N. D., and Ohkouchi, N.: Monsoons, Upwelling, and the
Deoxygenation of the Northwestern Indian Ocean in Response to Middle to Late Miocene Global Climatic Shifts,
Paleoceanogr Paleoclimatol, 35, <https://doi.org/10.1029/2019PA003762>, 2020.
- Brachert, T. C., Corrège, T., Reuter, M., Wrozyna, C., Londeix, L., Spreter, P., and Perrin, C.: An assessment of reef coral
calcification over the late Cenozoic, *Earth Sci Rev*, 204, 103154, <https://doi.org/10.1016/j.earscirev.2020.103154>, 2020.
- 530 Bridge, T. C. L., Beaman, R. J., Bongaerts, P., Muir, P. R., Ekins, M., and Sih, T.: The Great Barrier Reef and Coral Sea,
Springer, Cham, 351–367, https://doi.org/10.1007/978-3-319-92735-0_20, 2019.
- Brunet, M.: Sahelanthropus tchadensis dit « Toumaï » : le plus ancien membre connu de notre tribu, *Bull Acad Natl Med*, 204,
535 251–257, <https://doi.org/https://doi.org/10.1016/j.banm.2019.12.017>, 2020.
- Burls, N. J., Bradshaw, C. D., Boer, A. M. De, Herold, N., Huber, M., Pound, M., Donnadiou, Y., Farnsworth, A., Frigola, A.,
Gasson, E., Heydt, A. S. von der, Hutchinson, D. K., Knorr, G., Lawrence, K. T., Lear, C. H., Li, X., Lohmann, G., Lunt, D.
J., Marzocchi, A., Prange, M., Riihimaki, C. A., Sarr, A.-C., Siler, N., and Zhang, Z.: Simulating Miocene Warmth: Insights
540 From an Opportunistic Multi-Model Ensemble (MioMIP1), *Paleoceanogr Paleoclimatol*, 36, e2020PA004054,
<https://doi.org/10.1029/2020PA004054>, 2021.
- Clift, P. D., Du, Y., Mohtadi, M., Pahnke, K., Sutorius, M., and Böning, P.: The erosional and weathering response to arc-
continent collision in New Guinea, *J Geol Soc London*, 181, <https://doi.org/10.1144/jgs2023-207>, 2024.
- 545 Collins, L. S., Coates, A. G., Berggren, W. A., Aubry, M.-P. M. _P., and Zhang, J.: The late Miocene Panama isthmian strait,
Geology, 24, 687–690, 1996.
- Cornwall, C. E., Comeau, S., Kornder, N. A., Perry, C. T., van Hooidek, R., DeCarlo, T. M., Pratchett, M. S., Anderson, K.
550 D., Browne, N., Carpenter, R., Diaz-Pulido, G., D’Olivo, J. P., Doo, S. S., Figueiredo, J., Fortunato, S. A. V., Kennedy, E.,
Lantz, C. A., McCulloch, M. T., González-Rivero, M., Schoepf, V., Smithers, S. G., and Lowe, R. J.: Global declines in coral
reef calcium carbonate production under ocean acidification and warming, *Proc Natl Acad Sci U S A*, 118,
https://doi.org/10.1073/PNAS.2015265118/SUPPL_FILE/PNAS.2015265118.SD02.XLSX, 2021.
- 555 Crandall, E. D., Riginos, C., Bird, C. E., Liggins, L., Treml, E., Beger, M., Barber, P. H., Connolly, S. R., Cowman, P. F.,
DiBattista, J. D., Eble, J. A., Magnuson, S. F., Horne, J. B., Kochzius, M., Lessios, H. A., Liu, S. Y. V., Ludt, W. B., Madduppa,
H., Pandolfi, J. M., Toonen, R. J., Network, C. M. of the D. of the I.-P., and Gaither, M. R.: The molecular biogeography of

the Indo-Pacific: Testing hypotheses with multispecies genetic patterns, *Global Ecology and Biogeography*, 28, 943–960, <https://doi.org/https://doi.org/10.1111/geb.12905>, 2019.

560

Darling, E. S. and Côté, I. M.: 4.21 - Vulnerability of Coral Reefs, edited by: Pielke, R. A. B. T.-C. V., Academic Press, Oxford, 259–270, <https://doi.org/https://doi.org/10.1016/B978-0-12-384703-4.00427-5>, 2013.

Davies, P. J., Mackenzie, J. A., and Palmer-Julson, A.: Site 811-826 Northeast Australian Margin, in: Proc. ODP Init. Rep, 565 810, 1991.

DiCaprio, L., Gurnis, M., and Müller, R. D.: Long-wavelength tilting of the Australian continent since the Late Cretaceous, *Earth Planet Sci Lett*, 278, 175–185, <https://doi.org/10.1016/j.epsl.2008.11.030>, 2009.

570 Diester-Haass, L., Meyers, P. A., and Bickert, T.: Carbonate crash and biogenic bloom in the late Miocene: Evidence from ODP Sites 1085, 1086, and 1087 in the Cape Basin, southeast Atlantic Ocean, *Paleoceanography*, 19, 1007, <https://doi.org/10.1029/2003PA000933>, 2004.

575 Droxler, A. W., Haddad, G. A., Kroon, D., Gartner, S., Wuchang Wei, and McNeill, D.: Late Pliocene (2.9 Ma) partial recovery of shallow carbonate banks on the Queensland Plateau: signal of bank-top reentry into the photic zone during a lowering in sea level, Proc., scientific results, ODP, Leg 133, northeast Australian margin, 235–254, <https://doi.org/10.2973/ODP.PROC.SR.133.227.1993>, 1993.

580 Drury, A. J., Lee, G. P., Gray, W. R., Lyle, M., Westerhold, T., Shevenell, A. E., and John, C. M.: Deciphering the state of the late Miocene to early Pliocene equatorial Pacific, *Paleoceanogr Paleoclimatol*, <https://doi.org/10.1002/2017PA003245>, 2018.

Dupont, L. M., Rommerskirchen, F., Mollenhauer, G., and Schefuß, E.: Miocene to Pliocene changes in South African hydrology and vegetation in relation to the expansion of C4 plants, *Earth Planet Sci Lett*, 375, 408–417, <https://doi.org/https://doi.org/10.1016/j.epsl.2013.06.005>, 2013.

585

Eberli, G. P., Anselmetti, F. S., Isern, A. R., and Delius, H.: Timing of Changes in Sea-Level and Currents along Miocene Platforms on the Marion Plateau, Australia, in: *Cenozoic Carbonate Systems of Australasia*, SEPM (Society for Sedimentary Geology), 219–242, <https://doi.org/10.2110/sepm.095.219>, 2010.

- 590 Ehrenberg, S. N. N., McArthur, J. M. M., and Thirlwall, M. F. F.: Growth, Demise, and Dolomitization of Miocene Carbonate Platforms on the Marion Plateau, Offshore NE Australia, *Journal of Sedimentary Research*, 76, 91–116, <https://doi.org/10.2110/jsr.2006.06>, 2006.
- Feakins, S. J.: Pollen-corrected leaf wax D/H reconstructions of northeast African hydrological changes during the late
595 Miocene, *Palaeogeogr Palaeoclimatol Palaeoecol*, 374, 62–71, <https://doi.org/10.1016/j.palaeo.2013.01.004>, 2013.
- Feary, D. A., Davies, P. J., Pigram, C. J., and Symonds, P. A.: Climatic evolution and control on carbonate deposition in northeast Australia, *Palaeogeogr Palaeoclimatol Palaeoecol*, 89, 341–361, [https://doi.org/10.1016/0031-0182\(91\)90171-M](https://doi.org/10.1016/0031-0182(91)90171-M), 1991.
- 600 Fietz, S., Martínez-García, A., Hugué, C., Rueda, G., and Rosell-Melé, A.: Constraints in the application of the Branched and Isoprenoid Tetraether index as a terrestrial input proxy, *J Geophys Res*, 116, C10032, <https://doi.org/10.1029/2011JC007062>, 2011.
- 605 Gallagher, S. J., Auer, G., Brierley, C. M., Fulthorpe, C. S., and Hall, R.: Cenozoic History of the Indonesian Gateway, *Annu Rev Earth Planet Sci*, 52, 581–604, <https://doi.org/10.1146/annurev-earth-040722-111322>, 2024.
- Hall, R.: Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations, *J Asian Earth Sci*, 20, 353–431, [https://doi.org/10.1016/S1367-9120\(01\)00069-4](https://doi.org/10.1016/S1367-9120(01)00069-4), 2002.
- 610 Grant, K. M. and Dickens, G. R.: Coupled productivity and carbon isotope records in the southwest Pacific Ocean during the late Miocene–early Pliocene biogenic bloom, *Palaeogeogr Palaeoclimatol Palaeoecol*, 187, 61–82, [https://doi.org/10.1016/S0031-0182\(02\)00508-4](https://doi.org/10.1016/S0031-0182(02)00508-4), 2002.
- 615 Grimalt, J. O., Calvo, E., and Pelejero, C.: Sea surface paleotemperature errors in UK' 37 estimation due to alkenone measurements near the limit of detection, *Paleoceanography*, 16, 226–232, <https://doi.org/https://doi.org/10.1029/1999PA000440>, 2001.
- Groeneveld, J., Henderiks, J., Renema, W., McHugh, C. M., De Vleeschouwer, D., Christensen, B. A., Fulthorpe, C. S.,
620 Reuning, L., Gallagher, S. J., Bogus, K., Auer, G., Ishiwa, T., and Scientists, E. 356: Australian shelf sediments reveal shifts in Miocene Southern Hemisphere westerlies, *Sci Adv*, 3, e1602567, <https://doi.org/10.1126/sciadv.1602567>, 2017.
- Hall, R.: Southeast Asia's changing palaeogeography, *Blumea - Biodiversity, Evolution and Biogeography of Plants*, 54, 148–161, <https://doi.org/10.3767/000651909X475941>, 2009.

625 [Hallock, P., Sheps, K., Chapronière, G., and Howell, M., 2006. Larger benthic foraminifers of the Marion Plateau, northeastern Australia \(ODP Leg 194\): comparison of faunas from bryozoan \(Sites 1193 and 1194\) and red algal \(Sites 1196–1198\) dominated carbonate platforms. In Anselmetti, F.S., Isern, A.R., Blum, P., and Betzler, C. \(Eds.\), Proc. ODP, Sci. Results, 194, 1–31 \[Online\]. Available from World Wide Web: <\[http://www.odp.tamu.edu/publications/194_SR/VOLUME/CHAPTERS/009.PDF\]\(http://www.odp.tamu.edu/publications/194_SR/VOLUME/CHAPTERS/009.PDF\)>.](#)

630

Harrison, G. W. M., Santodomingo, N., Johnson, K. G., and Renema, W.: Is the Coral Triangle's future shown in a Pliocene reef gap?, *Coral Reefs*, <https://doi.org/10.1007/s00338-023-02412-5>, 2023.

635 Haug, G. H., Tiedemann, R., Zahn, R., and Ravelo, A. C.: Role of Panama uplift on oceanic freshwater balance, *Geology*, 29, 207–210, 2001.

Herbert, T. D., Lawrence, K. T., Tzanova, A., Peterson, L. C., Caballero-Gill, R., and Kelly, C. S.: Late Miocene global cooling and the rise of modern ecosystems, *Nat Geosci*, 9, 843–847, <https://doi.org/10.1038/ngeo2813>, 2016.

640 Herbert, T. D., Dalton, C. A., Liu, Z., Salazar, A., Si, W., and Wilson, D. S.: Tectonic degassing drove global temperature trends since 20 Ma, *Science*, 377, 116–119, 2022.

Deleted: ,

645 Hernández-Sánchez, M. T., Woodward, E. M. S., Taylor, K. W. R., Henderson, G. M., and Pancost, R. D.: Variations in GDGT distributions through the water column in the South East Atlantic Ocean, *Geochim Cosmochim Acta*, 132, 337–348, <https://doi.org/10.1016/j.gca.2014.02.009>, 2014.

Higuchi, T., Agostini, S., Casareto, B. E., Suzuki, Y., and Yuyama, I.: The northern limit of corals of the genus *Acropora* in temperate zones is determined by their resilience to cold bleaching, *Sci Rep*, 5, 18467, <https://doi.org/10.1038/srep18467>, 2015.

650

Van Hinsbergen, D. J. J., De Groot, L. V., Van Schaik, S. J., Spakman, W., Bijl, P. K., Sluijs, A., Langereis, C. G., and Brinkhuis, H.: A Paleolatitude Calculator for Paleoclimate Studies, *PLoS One*, 10, e0126946, <https://doi.org/10.1371/JOURNAL.PONE.0126946>, 2015.

655 Holbourn, A. E., Kuhnt, W., Clemens, S. C., Kochhann, K. G. D., Jöhnck, J., Lübbers, J., and Andersen, N.: Late Miocene climate cooling and intensification of southeast Asian winter monsoon, *Nat Commun*, 9, 1584, <https://doi.org/10.1038/s41467-018-03950-1>, 2018.

660 Hopmans, E. C., Schouten, S., and Sinninghe Damsté, J. S.: The effect of improved chromatography on GDGT-based
palaeoproxies, *Org Geochem*, 93, 1–6, <https://doi.org/10.1016/j.orggeochem.2015.12.006>, 2016.

Huang, Y., Clemens, S. C., Liu, W., Wang, Y., and Prell, W. L.: Large-scale hydrological change drove the late Miocene C4
plant expansion in the Himalayan foreland and Arabian Peninsula, *Geology*, 35, 531–534, <https://doi.org/10.1130/G23666A.1>,
665 2007.

Isern, A. R., McKenzie, J. A., and Müller, D. W.: Paleooceanographic Changes and Reef Growth off the Northeastern Australian
Margin: Stable Isotopic Data from ODP Leg 133 Sites 811 and 817 and DSDP Leg 21 Site 209, *Proceedings of the Ocean
Drilling Program, 133 Scientific Results*, <https://doi.org/10.2973/ODP.PROC.SR.133.230.1993>, 1993.

670

Isern, A. R., McKenzie, J. A., and Feary, D. A.: The role of sea-surface temperature as a control on carbonate platform
development in the western Coral Sea, *Palaeogeogr Palaeoclimatol Palaeoecol*, 124, 247–272, [https://doi.org/10.1016/0031-0182\(96\)80502-5](https://doi.org/10.1016/0031-0182(96)80502-5), 1996.

675 Isern, A. R., Anselmetti, F. S., and Blum, P.: A Neogene Carbonate Platform, Slope, and Shelf Edifice Shaped by Sea Level
and Ocean Currents, Marion Plateau (Northeast Australia), <https://doi.org/10.1306/M81928>, 1 January 2004.

John, C. M. and Mutti, M.: Relative Control of Paleooceanography, Climate, and Eustasy over Heterozoan Carbonates: A
Perspective from Slope Sediments of the Marion Plateau (ODP LEG 194), *Journal of Sedimentary Research*, 75, 216–230,
680 <https://doi.org/10.2110/jsr.2005.017>, 2005.

Jöhnc, J., Kuhnt, W., Holbourn, A., and Andersen, N.: Variability of the Indian Monsoon in the Andaman Sea Across the
Miocene-Pliocene Transition, *Paleoceanogr Paleoclimatol*, 35, <https://doi.org/10.1029/2020PA003923>, 2020.

685 Katz, M. E. and Miller, K. G.: Neogene Subsidence along the Northeastern Australian Margin: Benthic Foraminiferal
Evidence, *Proceedings of the Ocean Drilling Program, 133 Scientific Results*,
<https://doi.org/10.2973/ODP.PROC.SR.133.242.1993>, 1993a.

Katz, M. E. and Miller, K. G.: Neogene Subsidence along the Northeastern Australian Margin: Benthic Foraminiferal
690 Evidence, *Proceedings of the Ocean Drilling Program, 133 Scientific Results*,
<https://doi.org/10.2973/ODP.PROC.SR.133.242.1993>, 1993b.

- Kim, J.-H., van der Meer, J., Schouten, S., Helmke, P., Willmott, V., Sangiorgi, F., Koç, N., Hopmans, E. C., Damsté, J. S. S., and Damste, J. S. S.: New indices and calibrations derived from the distribution of crenarchaeal isoprenoid tetraether lipids: Implications for past sea surface temperature reconstructions, *Geochim Cosmochim Acta*, 74, 4639–4654, <https://doi.org/http://dx.doi.org/10.1016/j.gca.2010.05.027>, 2010a.
- 695
- Kim, J.-H., der Meer, J., Schouten, S., Helmke, P., Willmott, V., Sangiorgi, F., Koç, N., Hopmans, E. C., and Damsté, S.: New indices and calibrations derived from the distribution of crenarchaeal isoprenoid tetraetherlipids: Implications for past sea surface temperature reconstructions, *Geochimica et Cosmochimica Acta*, 74, 4639–4654, 2010b.
- 700
- Laugié, M., Michel, J., Pohl, A., Poli, E., and Borgomano, J.: Global distribution of modern shallow-water marine carbonate factories: a spatial model based on environmental parameters, *Sci Rep*, 9, 16432, <https://doi.org/10.1038/s41598-019-52821-2>, 2019.
- 705
- Liu, C., Fulthorpe, C. S., Austin, J. A., and Sanchez, C. M.: Geomorphologic indicators of sea level and lowstand paleo-shelf exposure on early–middle Miocene sequence boundaries, *Mar Geol*, 280, 182–194, <https://doi.org/10.1016/j.margeo.2010.12.010>, 2011.
- 710
- Liu, X., Huber, M., Foster, G. L., Dessler, A., and Zhang, Y. G.: Persistent high latitude amplification of the Pacific Ocean over the past 10 million years, *Nat Commun*, 13, 7310, <https://doi.org/10.1038/s41467-022-35011-z>, 2022b.
- 715
- Locarnini, R. A., Mishonov, A. V., Baranova, O. K., Boyer, T. P., Zweng, M. M., Garcia, H. E., Reagan, J. R., Seidov, D., Weathers, K. W., Paver, C. R., and Smolyar, I. V.: *World Ocean Atlas*, 1, 52, 2019.
- Lough, J. M.: Coral calcification from skeletal records revisited, *Mar Ecol Prog Ser*, 373, 257–264, <https://doi.org/10.3354/MEPS07398>, 2008.
- Lough, J. M. and Barnes, D. J.: Environmental controls on growth of the massive coral *Porites*, *J Exp Mar Biol Ecol*, 245, 225–243, [https://doi.org/https://doi.org/10.1016/S0022-0981\(99\)00168-9](https://doi.org/https://doi.org/10.1016/S0022-0981(99)00168-9), 2000.
- 720
- Lough, J. M. and Cantin, N. E.: Perspectives on massive coral growth rates in a changing ocean, *Biological Bulletin*, 226, 187–202, <https://doi.org/10.1086/BBLV226N3P187/ASSET/IMAGES/LARGE/Z1N0031435890005.JPEG>, 2014a.

725 Lübbers, J., Kuhnt, W., Holbourn, A. E., Bolton, C. T., Gray, E., Usui, Y., Kochhann, K. G. D., Beil, S., and Andersen, N.:
The Middle to Late Miocene “Carbonate Crash” in the Equatorial Indian Ocean, *Paleoceanogr Paleoclimatol*, 34, 813–832,
<https://doi.org/10.1029/2018PA003482>, 2019.

Martin, P. E., Macdonald, F. A., McQuarrie, N., Flowers, R. M., and Maffre, P. J. Y.: The rise of New Guinea and the fall of
730 Neogene global temperatures, *Proceedings of the National Academy of Sciences*, 120,
<https://doi.org/10.1073/pnas.2306492120>, 2023.

Martinot, C., Bolton, C. T., Sarr, A.-C., Donnadiou, Y., Garcia, M., Gray, E., and Tachikawa, K.: Drivers of Late Miocene
Tropical Sea Surface Cooling: A New Perspective From the Equatorial Indian Ocean, *Paleoceanogr Paleoclimatol*, 37,
735 e2021PA004407, <https://doi.org/https://doi.org/10.1029/2021PA004407>, 2022.

[Mathew, M., Makhankova, A., Menier, D., Sautter, B., Betzler, C., and Pierson, B.: The emergence of Miocene reefs in South
China Sea and its resilient adaptability under varying eustatic, climatic and oceanographic conditions, *Sci Rep*, 10, 7141,
<https://doi.org/10.1038/s41598-020-64119-9>, 2020.](https://doi.org/10.1038/s41598-020-64119-9)

740

Miller, K. G., Browning, J. V., John Schmelz, W., Kopp, R. E., Mountain, G. S., and Wright, J. D.: Cenozoic sea-level and
cryospheric evolution from deep-sea geochemical and continental margin records, *Sci Adv*, 6,
<https://doi.org/10.1126/SCIADV.AAZ1346>, 2020.

745

Müller, P. J., Kirst, G., Ruhland, G., von Storch, I., and Rosell-Mele, A.: Calibration of the alkenone paleotemperature index
U-37(K ') based on core-tops from the eastern South Atlantic and the global ocean (60 degrees N-60 degrees S), *Geochim
Cosmochim Acta*, 62, 1757–1772, 1998.

750 Nairn, M. G., Lear, C. H., Sosdian, S. M., Bailey, T. R., and Beavington-Penney, S.: Tropical Sea Surface Temperatures
Following the Middle Miocene Climate Transition From Laser-Ablation ICP-MS Analysis of Glassy Foraminifera,
Paleoceanogr Paleoclimatol, 36, e2020PA004165, <https://doi.org/10.1029/2020PA004165>, 2021.

Pelejero, C. and Calvo, E.: The upper end of the UK′37 temperature calibration revisited, *Geochemistry Geophysics
755 Geosystems*, 4, 1014, <https://doi.org/10.1029/2002gc000431>, 2003.

Perrin, C. and Kiessling, W.: Latitudinal Trends in Cenozoic Reef Patterns and their Relationship to Climate, in: Carbonate Systems during the Oligocene–Miocene Climatic Transition, Wiley, 17–33, <https://doi.org/10.1002/9781118398364.ch2>, 2012.

760

Petrick, B., McClymont, E. L., Littler, K., Rosell-Melé, A., Clarkson, M. O., Maslin, M., Röhl, U., Shevenell, A. E., and Pancost, R. D.: Oceanographic and climatic evolution of the southeastern subtropical Atlantic over the last 3.5 Ma, *Earth Planet Sci Lett*, 492, 12–21, <https://doi.org/10.1016/j.epsl.2018.03.054>, 2018.

765 Petrick, B., Martínez-García, A., Auer, G., Reuning, L., Auderset, A., Deik, H., Takayanagi, H., De Vleeschouwer, D., Iryu, Y., and Haug, G. H.: Glacial Indonesian Throughflow weakening across the Mid-Pleistocene Climatic Transition, *Sci Rep*, 9, 16995, <https://doi.org/10.1038/s41598-019-53382-0>, 2019.

Petrick, B., Reuning, L., Auer, G., Zhang, Y., Pfeiffer, M., and Schwark, L.: Warm, not cold temperatures contributed to a
770 Late Miocene reef decline in the Coral Sea, *Sci Rep*, 13, 4015, <https://doi.org/10.1038/s41598-023-31034-8>, 2023.

Renema, W., Bellwood, D. R., Braga, J. C., Bromfield, K., Hall, R., Johnson, K. G., Lunt, P., Meyer, C. P., McMonagle, L. B., Morley, R. J., O’Dea, A., Todd, J. A., Wesselingh, F. P., Wilson, M. E. J., and Pandolfi, J. M.: Hopping hotspots: global shifts in marine biodiversity., *Science*, 321, 654–7, <https://doi.org/10.1126/science.1155674>, 2008.

775

Rattanasriampaipong, R., Zhang, Y. G., Pearson, A., Hedlund, B. P., and Zhang, S.: Archaeal lipids trace ecology and evolution of marine ammonia-oxidizing archaea, *Proceedings of the National Academy of Sciences*, 119, e2123193119, <https://doi.org/10.1073/PNAS.2123193119>, 2022.

780 Rich, W. A., Carvalho, S., and Berumen, M. L.: Coral bleaching due to cold stress on a central Red Sea reef flat, *Ecol Evol*, 12, e9450, <https://doi.org/10.1002/ece3.9450>, 2022.

Rosleff-Soerensen, B., Reuning, L., Back, S., and Kukla, P.: Seismic geomorphology and growth architecture of a Miocene barrier reef, Browse Basin, NW-Australia, *Mar Pet Geol*, 29, 233–254, <https://doi.org/10.1016/j.marpetgeo.2010.11.001>, 2012.

785

Rosleff-Soerensen, B., Reuning, L., Back, S., and Kukla, P. A.: The response of a basin-scale Miocene barrier reef system to long-term, strong subsidence on a passive continental margin, Barcoo Sub-basin, Australian North West Shelf, *Basin Research*, 28, 103–123, <https://doi.org/10.1111/bre.12100>, 2016.

790 Santodomingo, N., Wallace, C. C., and Johnson, K. G.: Fossils reveal a high diversity of the staghorn coral genera *Acropora*
and *Isopora* (Scleractinia: Acroporidae) in the Neogene of Indonesia, *Zool J Linn Soc*, 175, 677–763,
<https://doi.org/10.1111/zoj.12295>, 2015.

Santodomingo, N., Renema, W., and Johnson, K. G.: Understanding the murky history of the Coral Triangle: Miocene corals
795 and reef habitats in East Kalimantan (Indonesia), *Coral Reefs* 2016 35:3, 35, 765–781, <https://doi.org/10.1007/S00338-016-1427-Y>, 2016.

Schlitzer, Reiner, Ocean Data View, odv.awi.de, 2021

800 Schouten, S., Hopmans, E. C., Schefuss, E., and Sinninghe Damsté, J. S.: Distributional variations in marine crenarchaeotal
membrane lipids: a new tool for reconstructing ancient sea water temperatures?, *Earth Planet Sci Lett*, 204, 265–274, 2002.

Schouten, S., Hopmans, E. C., Rosell-Melé, A., Pearson, A., Adam, P., Bauersachs, T., Bard, E., Bernasconi, S. M., Bianchi,
T. S., Brocks, J. J., Carlson, L. T., Castañeda, I. S., Derenne, S., Selver, A. D., Dutta, K., Eglinton, T., Fosse, C., Galy, V.,
805 Grice, K., Hinrichs, K.-U., Huang, Y., Huguet, A., Huguet, C., Hurley, S., Ingalls, A., Jia, G., Keely, B., Knappy, C., Kondo,
M., Krishnan, S., Lincoln, S., Lipp, J., Mangelsdorf, K., Martínez-García, A., Ménot, G., Mets, A., Mollenhauer, G., Ohkouchi,
N., Ossebaar, J., Pagani, M., Pancost, R. D., Pearson, E. J., Peterse, F., Reichart, G.-J., Schaeffer, P., Schmitt, G., Schwark,
L., Shah, S. R., Smith, R. W., Smittenberg, R. H., Summons, R. E., Takano, Y., Talbot, H. M., Taylor, K. W. R., Tarozo, R.,
Uchida, M., van Dongen, B. E., Van Mooy, B. A. S., Wang, J., Warren, C., Weijers, J. W. H., Werne, J. P., Woltering, M.,
810 Xie, S., Yamamoto, M., Yang, H., Zhang, C. L., Zhang, Y., Zhao, M., and Damsté, J. S. S.: An interlaboratory study of TEX
86 and BIT analysis of sediments, extracts, and standard mixtures, *Geochemistry, Geophysics, Geosystems*, 14, 5263–5285,
<https://doi.org/10.1002/2013GC004904>, 2013.

Shipboard Scientific Party: Principal Results and Summary, in: Proceedings of the Ocean Drilling Program, 133 Initial Reports,
815 vol. 133, edited by: Davies, P. J., McKenzie, J. A., and Palmer-Julson, A., Ocean Drilling Program, 59–72,
<https://doi.org/10.2973/odp.proc.ir.133.103.1991>, 1991.

Sinninghe Damsté, J. S., Ossebaar, J., Schouten, S., and Verschuren, D.: Distribution of tetraether lipids in the 25-ka
820 sedimentary record of Lake Challa: extracting reliable TEX86 and MBT/CBT palaeotemperatures from an equatorial African
lake, *Quat Sci Rev*, 50, 43–54, <https://doi.org/10.1016/j.quascirev.2012.07.001>, 2012.

- Spalding, M. D., Fox, H. E., Allen, G. R., Davidson, N., Ferdaña, Z. A., Finlayson, M., Halpern, B. S., Jorge, M. A., Lombana, A., Lourie, S. A., Martin, K. D., McManus, E., Molnar, J., Recchia, C. A., and Robertson, J.: Marine Ecoregions of the World: A Bioregionalization of Coastal and Shelf Areas, *Bioscience*, 57, 573–583, <https://doi.org/10.1641/B570707>, 2007.
- 825
- Steinthorsdottir, M., Coxall, H. K., de Boer, A. M., Huber, M., Barbolini, N., Bradshaw, C. D., Burls, N. J., Feakins, S. J., Gasson, E., Henderiks, J., Holbourn, A. E., Kiel, S., Kohn, M. J., Knorr, G., Kürschner, W. M., Lear, C. H., Liebrand, D., Lunt, D. J., Mörs, T., Pearson, P. N., Pound, M. J., Stoll, H., and Strömberg, C. A. E.: The Miocene: The Future of the Past, *Paleoceanogr Paleoclimatol*, 36, <https://doi.org/10.1029/2020PA004037>, 2021.
- 830
- Strömberg, C. A. E. and Strömberg, S.: Evolution of Grasses and Grassland Ecosystems, <https://doi.org/10.1146/annurev-earth-040809-152402>, 39, 517–544, <https://doi.org/10.1146/ANNUREV-EARTH-040809-152402>, 2011.
- 835
- Tagliaro, G., Fulthorpe, C. S., Gallagher, S. J., McHugh, C. M., Kominz, M., and Lavier, L. L.: Neogene siliciclastic deposition and climate variability on a carbonate margin: Australian Northwest Shelf, *Mar Geol*, 403, 285–300, <https://doi.org/10.1016/J.MARGEO.2018.06.007>, 2018.
- Tanner, T., Hernández-Almeida, I., Drury, A. J., Guitián, J., and Stoll, H.: Decreasing Atmospheric CO₂ During the Late Miocene Cooling, *Paleoceanogr Paleoclimatol*, 35, <https://doi.org/10.1029/2020PA003925>, 2020.
- 840
- Taylor, K. W. R., Huber, M., Hollis, C. J., Hernandez-Sanchez, M. T., and Pancost, R. D.: Re-evaluating modern and Palaeogene GDGT distributions: Implications for SST reconstructions, *Glob Planet Change*, 108, 158–174, <https://doi.org/10.1016/j.gloplacha.2013.06.011>, 2013.
- 845
- Thronberens, S., Back, S., Bourget, J., Allan, T., and Reuning, L.: 3-D seismic chronostratigraphy of reefs and drifts in the Browse Basin, NW Australia, *GSA Bulletin*, 134, 3155–3175, <https://doi.org/10.1130/B36286.1>, 2022.
- Vernon, J. E. N., Devantier, L. M., Turak, E., Green, A. L., Kininmonth, S., Stafford-Smith, M., and Petetson, N.: Delineating the Coral Triangle, *Galaxea, Journal of Coral Reef Studies*, 11, 91–100, <https://doi.org/10.3755/galaxea.11.91>, 2009.
- 850
- De Vleeschouwer, D., Auer, G., Smith, R., Bogus, K., Christensen, B., Groeneveld, J., Petrick, B., Henderiks, J., Castañeda, I. S., O'Brien, E., Ellinghausen, M., Gallagher, S. J., Fulthorpe, C. S., and Pälke, H.: The amplifying effect of Indonesian Throughflow heat transport on Late Pliocene Southern Hemisphere climate cooling, *Earth Planet Sci Lett*, 500, 15–27, <https://doi.org/10.1016/j.epsl.2018.07.035>, 2018.
- 855

- De Vleeschouwer, D., Petrick, B. F., and Martínez-García, A.: Stepwise Weakening of the Pliocene Leeuwin Current, *Geophys Res Lett*, 46, 8310–8319, <https://doi.org/10.1029/2019GL083670>, 2019.
- 860 Weijers, J. W. H., Schouten, S., Hopmans, E. C., Geenevasen, J. A. J., David, O. R. P., Coleman, J. M., Pancost, R. D., and Sinninghe Damsté, J. S.: Membrane lipids of mesophilic anaerobic bacteria thriving in peats have typical archaeal traits, *Environ Microbiol*, 8, 648–657, <https://doi.org/10.1111/j.1462-2920.2005.00941.x>, 2006.
- Wen, Y., Zhang, L., Holbourn, A. E., Zhu, C., Huntington, K. W., Jin, T., Li, Y., and Wang, C.: CO₂-forced Late Miocene cooling and ecosystem re-organizations in East Asia, *Proceedings of the National Academy of Sciences*, 120, e2214655120, <https://doi.org/10.1073/pnas.2214655120>, 2023.
- 875 Westerhold, T., Marwan, N., Drury, A. J., Liebrand, D., Agnini, C., Anagnostou, E., Barnet, J. S. K., Bohaty, S. M., De Vleeschouwer, D., Florindo, F., Frederichs, T., Hodell, D. A., Holbourn, A. E., Kroon, D., Lauretano, V., Littler, K., Lourens, L. J., Lyle, M., Pälike, H., Röhl, U., Tian, J., Wilkens, R. H., Wilson, P. A., and Zachos, J. C.: An astronomically dated record of Earth's climate and its predictability over the last 66 million years, *Science* (1979), 369, 1383–1387, <https://doi.org/10.1126/science.aba6853>, 2020.
- Wilson, P. A., Norris, R. D., and Cooper, M. J.: Testing the Cretaceous greenhouse hypothesis using glassy foraminiferal calcite from the core of the Turonian tropics on Demerara Rise, *Geology*, 30, 607–610, [https://doi.org/10.1130/0091-7613\(2002\)030<0607:TTCGHU>2.0.CO;2](https://doi.org/10.1130/0091-7613(2002)030<0607:TTCGHU>2.0.CO;2), 2002.
- Zachos, J. C., Stott, L. D., and Lohmann, K. C.: Evolution of early Cenozoic marine temperatures, *Paleoceanography*, 9, 353–387, <https://doi.org/10.1029/93PA03266>, 1994.
- 880 Zhang, Y. G., Zhang, C. L., Liu, X.-L., Li, L., Hinrichs, K.-U., and Noakes, J. E.: Methane Index: A tetraether archaeal lipid biomarker indicator for detecting the instability of marine gas hydrates, *Earth Planet Sci Lett*, 307, 525–534, <https://doi.org/10.1016/J.EPSL.2011.05.031>, 2011.
- 885 Zhang, Y. G., Pagani, M., and Liu, Z.: A 12-Million-Year Temperature History of the Tropical Pacific Ocean, *Science* (1979), 344, 84–87, <https://doi.org/10.1126/science.1246172>, 2014.
- Zhang, Y. G., Pagani, M., and Wang, Z.: Ring Index: A new strategy to evaluate the integrity of TEX 86 paleothermometry, *Paleoceanography*, 31, 220–232, <https://doi.org/10.1002/2015PA002848>, 2016.