



Variations in the Biological Pump through the Miocene: Evidence from organic carbon burial in Pacific Ocean sediments.

Mitchell Lyle

5 Annette Olivarez Lyle

College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, CEOAS Admin Bldg 104, Corvallis Oregon 97333, USA

Correspondence to: Mitchell Lyle (Lylem@oregonstate.edu)

10

Abstract. The biological pump, defined as the marine biological production and sedimentation of particulate organic carbon (C_{org}), is a fundamental process to fix atmospheric carbon dioxide in the oceans, transfer carbon away from the atmosphere to the deep ocean, and maintain the CO_2 level of the atmosphere. The level of carbon sequestration by the biological pump has varied throughout the last 50 million years, from particularly weak in the warm Eocene to much stronger in the Holocene. However, persistently warm climates in the more recent past, e.g., the Miocene Climate Optimum (MCO; 17 million years ago [Ma] to 13.8 Ma) also have affected the biological sequestration of carbon. A series of scientific ocean drill sites from the equatorial Pacific contain very low sedimentary C_{org} % in the period prior to 14 Ma but higher and much more variable C_{org} % afterward. Although lower absolute productivity may have contributed to the lower C_{org} burial at the MCO, higher relative C_{org} degradation also occurred. Ratios of C_{org} to other productivity indicators indicate higher relative loss of C_{org} . Temperature records imply that the higher C_{org} degradation occurred in the upper water column, and global cooling strengthened the biological pump but led to more variability in burial. Similar records of low C_{org} at the MCO can be found in the North Pacific, which suggest this was a global—rather than regional—change. A weakened biological pump during warm climate intervals helps to sustain periods of global warmth.

15

20

25

1 Introduction

The sequestration of organic carbon in pelagic sediments sums all the biological, physical, and chemical processes; from carbon fixation by photosynthesis, water-column transport and degradation, to ultimate sea floor deposition and burial. Heterotrophic consumption as well as inorganic oxidation greatly reduces the total particulate organic carbon (C_{org}) mass falling from the surface and its reactivity. C_{org} that is eventually buried in pelagic sediments is protected from further attack because the remainder is relatively recalcitrant after all the water column degradation and because C_{org} binds to sediments and is less accessible for further degradation (Hedges and Keil, 1995; Mayer, 1995). Buried organic matter represents a small fraction, less than 1%, of the original primary productivity in the pelagic realm (Suess, 1980, Muller and Suess, 1979; Martin et al., 1987). In the Pleistocene pelagic equatorial Pacific, C_{org} content found in surface sediments is typically low, ~ 0.2% (Murray and Leinen, 1993) while ocean margin surface sediments contain much higher C_{org} (between 1-2%). The difference results from lower primary productivity in the equatorial Pacific, deep waters (>4 km for the equatorial Pacific vs <1 km for the ocean margins), and much slower sedimentation rates that allow the C_{org} to be exposed to oxygenated sea water for thousands of years.

30

35

There is evidence that during eras of global warmth C_{org} degradation tends to be larger and occurs higher up in the water column (John et al, 2014; Boscolo-Golazzo et al, 2021). Upper water column C_{org} degradation results in poorer sequestration of



40 atmospheric CO₂, higher atmospheric CO₂ in dynamic equilibrium with the oceans, and higher nutrient levels in upwelled
waters. Unlike the last century, where there is intense global warming in the surface above a cold ocean, there is not necessarily a
strong oxygen minimum during stable warm periods. Long periods of warming deepen the permanent thermocline and weaken
the pycnocline, allowing more extensive mixing of oxygen into the ocean in contrast to the modern condition, where surface
warming strengthens the temperature and density contrast between the mixed layer and colder deeper waters. The resulting
45 strong pycnocline in the modern ocean restricts mixing and oxygen transport downward. Higher dissolved oxygen and higher
water temperatures found in long warm intervals thus intensifies C_{org} degradation in surface waters and reduces burial rates.
Nutrients and CO₂ resulting from C_{org} degradation cycle back to the surface faster under warm earth conditions. Surface water
that is subducted to shallow depths often reappears at the surface in decades rather than centuries for waters that are cycled to
abyssal depths. Therefore, the biological pump weakens under longer intervals of global warming (Boyd, 2015). Indeed, a recent
50 paper by Li et al (2023) estimated global C_{org} burial in the Neogene and found much less C_{org} burial than expected during the
Miocene Climate Optimum (MCO) interval.

The hypothesis that C_{org} degradation occurred much shallower in the water column during warm ocean conditions is supported
by John et al. (2014). They measured ocean depth gradients in carbon isotopes measured on planktonic foraminifera compared to
55 earth system model representations for the Paleocene-Eocene Thermal Maximum (PETM) and during the greenhouse Eocene to
conclude that the Eocene had shallower C_{org} degradation and recycling. Similarly, Boscolo-Galazzo et al (2021) examined the
change in the abundance of planktonic foraminifera living at different depths in the ocean in the period from 15 to 0 Ma and used
stable carbon isotope values of the foraminiferal tests, combined with earth system models, to monitor changes in carbon flux
through the water column. When combined with modeling of temperature dependent C_{org} degradation, their results were
60 consistent with cooling having caused more particulate C_{org} rain to penetrate deeper into the water column to affect a stronger
biological pump.

The biological pump can also be studied by examining the C_{org} buried in the sediments. The C_{org} mass accumulation rate (MAR;
burial flux) depends upon all the processes in the water column above the sediments and is the obverse of measurements that
65 could be made in the water column because C_{org} does not sprout from the sediments below. It is a measure of the integrated
effectiveness of the biological pump. While high primary productivity increases the flow of particulate C_{org} to the sea floor,
increases in heterotrophic consumption by zooplankton and microbes, reduce the resulting downward flux. Because of the
fundamental temperature dependence of metabolic activity (e.g., see Brown et al, 2004), higher water temperatures increases the
rate of C_{org} degradation by zooplankton and microbes, and decreases C_{org} burial (e.g., Boscolo-Galazo et al., 2018).

70 Preservation of C_{org} in sediments can also be studied by comparing it to other paleoproductivity measures that are better
preserved. For example, Biogenic barium (bio-Ba) deposition is strongly correlated to C_{org} rain from the euphotic zone (Dymond
and Collier, 1996) and is better and more consistently preserved at the sea floor under typical oxygenated conditions (Dehairet
et al, 1980; Dymond et al., 1992; Ganeshram et al., 2003). Lyle and Baldauf (2015) studied relative CaCO₃ dissolution in this way,
75 using ratios of CaCO₃ to Ba. Under warm earth conditions associated with early Cenozoic greenhouse conditions (Olivarez Lyle
and Lyle, 2005, 2006, Lyle et al., 2005) buried C_{org} % was extremely low (Eocene average of 0.03%). Olivarez Lyle and Lyle
(2005, 2006) used bio-Ba burial to estimate an 'expected' C_{org} MAR assuming modern conditions. All C_{org} burial during the
Eocene was much lower than modern C_{org} burial relative to production, indicating much higher heterotrophic consumption of
C_{org} within the water column. The ratio of C_{org} to Ba is thus an indicator of relative C_{org} preservation.



80

We produced long-term, low-resolution C_{org} and $CaCO_3$ data between 2008 and 2013 for drill sites extending into the Miocene in the Pacific Ocean (Fig 1, Table I). In this paper we report discrete C_{org} measurements we have made on 3 scientific drill sites from the eastern and central equatorial Pacific (ODP Site 574, and IODP Sites U1337, and U1338) and combine them with data from the Ontong Java Plateau in the western equatorial Pacific (ODP Sites 806 and 807; Stax and Stein, 1993). We also report

85

briefly on two sites from the northwest Pacific (Sites 884 and 1208) to show that changes in C_{org} burial were not limited to the equatorial region. We use these data to study changes in the biological carbon pump from the MCO to the present by comparing the patterns of C_{org} preservation through time and, where we have data, by comparing how C_{org} survives relative to bio-Ba, a better preserved paleoproductivity indicator.

90


Site	location	N is +		water depth (m)	5 Ma		10 Ma		15 Ma		20 Ma		Notes
		Latitude	Longitude		Latitude	Longitude	Latitude	Longitude	Latitude	Longitude			
U1338	E. equatorial Pacific	2.508	-117.969	4205	1.43	-113.79	0.49	-110.00	-0.37	-106.31	eq crossing at 13 Ma
U1337	E. equatorial Pacific	3.833	-123.206	4466	2.55	-119.07	1.50	-115.31	0.53	-111.63	-0.36	-107.93	eq crossing at 18 Ma
574	E. equatorial Pacific	4.209	-133.33	4571	2.60	-129.18	1.35	-125.41	0.21	-121.73	-0.86	-118.02	eq crossing at 16 Ma
806	W. equatorial Pacific	0.319	159.361	2521	-1.99	163.62	-3.41	167.39	-4.63	171.04	-5.90	174.69	eq crossing at <1 Ma
807	W. equatorial Pacific	3.607	159.625	2804	1.33	160.94	-0.05	164.76	-1.24	168.45	-2.48	172.13	eq crossing at 10 Ma
1208	N. Pacific, Shatsky Rise	36.127	158.208	3346	33.83	163.05	32.42	167.37	Hiatus to Paleocene >12 Ma
884	N. Pacific, Detroit Seamount	51.45	168.337	3827	49.05	173.09	47.52	177.48	46.19	-178.30	44.82	-174.21	eq crossing at 10 Ma

Table I: Locations of drill sites discussed in this paper and paleo-locations of the sites through the Miocene

95

Li et al. (2023) used a global set of mass accumulation rates to show that there was a global low in C_{org} MAR during the MCO, with a primary objective of the paper to test the “Monterey hypothesis” that high $d^{13}C$ between 17 and 13 Ma resulted from higher C_{org} burial. They found that the MCO interval had globally low C_{org} burial and was not a cause for high $d^{13}C$ at that time. The objective in this paper is different. We chose to examine a focused region (the equatorial Pacific) and use additional sedimentary data to separate changes in production from changes in preservation in the ultimate C_{org} record. We examine 4 hypotheses for the low C_{org} burial at the MCO— (1) low primary productivity, (2) C_{org} degradation in warmer deep waters, (3) C_{org} degradation in warmer surface waters, and (4) a fundamental change in the proxy- C_{org} relationship. We find that the low C_{org} MAR in the equatorial Pacific results largely from better preservation of sedimentary C_{org} as the earth cooled through lower degradation in the surface ocean.

100





105 **Figure 1: Drill site locations discussed in this paper. Paleo-locations for each site are shown at 5 Myr intervals, using a fixed hotspot rotation pole for the Pacific tectonic plate. The equatorial ODP drill sites 806 and 807 were drilled on ODP leg 130 (Stax and Stein, 1993. Sites U1337 and U1338 were drilled on IODP expedition 321 (Lyle et al, 2019; Pälike et al, 2010). Site 574 was drilled on DSDP Leg 85 (Mayer et al, 1985). Two other sites are briefly discussed—Site 1208 on the Shatsky Rise (Bralower et al., 2002) and Site 884 near Detroit Seamount at the northern end of the Hawaii-Emperor chain (Rea et al., 1993). Yellow numbers indicate the age of each site's equator crossing, a local high in primary productivity and sediment deposition.**

110

2 Analytical Methods

We measured over 1,600 samples (1,215 presented here plus 388 previously published) for C_{org} and $CaCO_3$, from 5 Sites of the Deep Sea Drilling Project, Ocean Drilling Program, and Integrated Ocean Drilling Program (Sites 574, 884, 1208, U1337, and U1338, see Fig. 1). Freeze-dried sediment samples were analyzed using coulometry and furnace methodologies presented in
115 Lyle et al., (2000) which provide accurate C_{org} values at very low levels. First, Total carbon of the sediment sample was measured through combustion in a 1000°C furnace and analyzed using a coulometer. Organic carbon was determined by acidifying a second, larger aliquot of the sample before analysis to remove the $CaCO_3$ -based CO_2 , then analysing the remaining residue via combustion in pure oxygen at 1000°C via coulometry. From these two measurements, the $CaCO_3$ fraction is calculated based on the difference between the CO_2 measured for Total carbon and the organic carbon fraction. We found that
120 this method is accurate for estimating $CaCO_3$ down to the <1% level and for C_{org} at the 0.01% level (Olivarez Lyle and Lyle, 2005).

We typically do not use C_{org} data from DSDP or ODP that were analysed shipboard because the method is inaccurate below 0.3% C_{org} when in the presence of significant sedimentary $CaCO_3$ (Olivarez Lyle and Lyle, 2005). Using our method of analysis,
125 data for Site 574 sediments older than 12 Ma are reported in Piela et al (2012). C_{org} data spanning 12 to 2 Ma were analyzed later and are published here. Similarly, C_{org} data from Sites 884 and 1208 are also first published here. For Site 884, we used shipboard C_{org} when the $CaCO_3$ % was less than 15% because the error of the shipboard method at this level of $CaCO_3$ is relatively small (<1.8 wt.% carbon from the $CaCO_3$ fraction vs ~0.1 to 0.4% for the C_{org} fraction). Utilising these shipboard data at Site 884 helped us to complete a full record of deposition. For Sites U1337 and U1338, C_{org} data are reported in Wilson (2014)
130 but were not discussed in her thesis. All the data are reported here in the Supplemental Material; Tables S1 to S20.

We include data from Stax and Stein (1993) for Sites 806 and 807. They measured C_{org} by dissolving an unweighed split of the sample with HCl and measuring C_{org} after dissolving the $CaCO_3$. They then corrected the split C_{org} to that of the total sample by measuring total carbon on a separate split of the sample and calculating the relative proportion of C_{org} . Using this method, they
135 could dissolve a much larger split of sediment for C_{org} , so that they were accurate at low C_{org} contents while also being able to measure other C_{org} properties.

The net deposition and preservation of C_{org} at Sites U1337, U1338, and Site 574 is estimated from ratios to the barium content in the sediments. Extensive work has shown this element is precipitated in the water column as $BaSO_4$ via organic matter oxidation,
140 is well-preserved, and importantly, is a good proxy for primary productivity (Dymond et al, 1992, Dymond and Collier, 1996; Ganeshram et al., 2003). Although barium sulfate can form through other processes, in equatorial Pacific sediments, the Ba content is primarily biogenic in origin and not diluted by other Ba-containing components, for example, by barium in terrigenous clays (Piela et al., 2012; Lyle and Baldauf, 2015)). We calibrated Ba measurements from X-ray fluorescence (XRF) (normalized



145 median-scaled ‘NMS’, see Lyle et al, 2012) to discrete ICP-MS analyses for Sites U1337 and U1338 (Wilson, 2014). At Site 574 we used ICP-MS barium data from Piela et al. (2012) for sediments older than 12 Ma.

150 We use total XRF-estimated total BaSO₄ rather than bio-Ba to ratio to other biogenic elements in this paper to investigate C_{org} preservation. Total barium is justified, as stated, because the equatorial Pacific sediments at Sites U1337 and U1338 primarily consist of biogenic remains and contain little clays, so the majority of Ba is biogenic in origin. We further tested this assumption based on an estimate of terrigenous, authigenic oxide, and authigenic clay-based Ba at Site U1338 and found that biogenic BaSO₄ averaged 93% ±4% of total BaSO₄. The greatest error in our biogenic barium estimates occurs when this fraction is at its lowest in the presence of relatively abundant clays, and are located primarily at the tops of the sites. Such a bias towards shallow core depths consequently skews the C_{org}/Ba ratio to lower values in younger sediments where the C_{org}/Ba and estimated preservation is typically highest.

155

3 Age Models and Mass Accumulation Rates (MAR)

160 For all sites we calculated C_{org} mass accumulation rates (MAR), equivalent to C_{org} burial flux, as another indication of the rate of C_{org} deposition and burial. MAR has units of mass per unit area per unit time and typical reported as “grams/(cm² x 10³yr)”. For any component in a sedimentary mixture, calculating its mass accumulation rate eliminates artifacts in the resulting data profile over depth/time that are necessarily caused by variable deposition of the other sedimentary components. Converting weight % data to MAR is particularly important for evaluating changes in minor components such as C_{org} because change in deposition of the major components can dominate and distort actual changes of the component of interest. However, MAR calculations are subject to errors (for example, sedimentation rates) that are primarily caused by an imprecise age model. This issue is addressed below.

165

We developed age models for all sites to calculate MAR to further study rates of burial. See the Supplemental Material for details and for data from each of the sites. MARs are developed by making an age model (age vs. depth profile) that is differentiated to calculate a sedimentation rate (thickness of sediment deposited per unit time; cm per kyr) over the time interval of interest. Sedimentation rate, multiplied by the Dry Bulk Density results in a bulk MAR (g solid/cm²/kyr) for the sediment over a given age interval. Individual component MARs are calculated by multiplying the bulk MAR by the weight fraction (wt %/100) of the component in the dry sample.

170

3.1 Sites U1337, U1338

175 For Site U1337 (24 Ma crust) we used an astro-chronologically-tuned age model to 20 Ma by combining the Drury et al (2017, 2018) age model to 8.2 Ma and correlating the 20 Myr combined isotope record from Site U1337 (Holbourn et al., 2015; Tian et al., 2018) to the stable isotope stack from Westerhold et al (2020) to 20 Ma. For Site U1338 (18 Ma crust) we used the established age model to 8.2 Ma (Lyle et al., 2019). The U1338 ages greater than 8.2 Ma were derived by correlating the U1338 CaCO₃ profile to that from Site U1337, which is justified because CaCO₃ profiles are very similar across the central and eastern Pacific (Lyle et al, 2019; Mayer,1991).

180

3.2 Site 574

Site 574 (35 Ma crust) was drilled in 1985 during Leg 85 of the Deep Sea Drilling Project and lies about 1000 km west of Site U1337. It was one of the first sites to have hydraulically piston cored sediments. We XRF-scanned the upper 5 cores of Holes



185 574 and 574A and used these data and the GRA bulk density record in deeper sections to make a new splice to ~17.4 Ma (225 m
composite depth, the base of Holes 574 and 574A) using the *Code for Ocean Drilling Data* software (CODD; <https://www.codd-home.net/>, last access: 08 August 2023). The new splice was then correlated to Site U1337 by a CaCO₃ profile produced by
using the Site 574 GRA bulk density to estimate CaCO₃ content. In the eastern and central Pacific Ocean, the GRA bulk density
is highly correlated with changes in CaCO₃ content (Mayer, 1991). The noncarbonate fraction in the equatorial Pacific is
190 primarily low density, high porosity biogenic silica remains of diatoms. The GRA bulk density data is also much higher
resolution than the discrete carbonate analyses used for calibration. The spliced GRA density record from Site 574 is in the
supplemental tables with assigned ages (Supplemental Table S2), as well as the new splice developed for Site 574 (Supplemental
Table S1).

3.3 Sites 806 and 807

195 We used polynomial fits to ages updated to the biostratigraphic ages used for Sites U1337 and U1338 (Expedition 320/321
scientists, 2010) for the biostratigraphic datum levels for Sites 806 and 807 reported in the Initial Reports volume (Kroenke et
al., 1991). Such fits are less accurate than a direct correlation, but these sites do not have long stable isotope records, nor was it
possible to correlate the carbonate records to the eastern Pacific. While not as accurate as the other age models, ages should still
be good to ± 0.2 Ma.

200

3.4 Sites 1208 and 884

Age-depth profiles for Sites 1208 (Shatsky Rise, to 12.4 Ma) and 884 (Detroit Seamount, at the north end of the Hawaii-Emperor
seamount chain, to 19.7 Ma) were made using linear interpolations between magnetochrons. We used Evans (2006) for
magnetochrons from Site 1208, and shipboard paleomagnetic data for Site 884 (Rea et al., 1993). All magnetochrons have
205 updated ages based on Westerhold et al. (2020). Accuracy of ages should be better than 0.1 Ma. The profiles are listed in the
supplemental material. Site 1208 was located on the Shatsky Rise itself and had an extreme slowdown in sedimentation from the
middle Miocene through the late Cretaceous which may have affected the early part of the profile.

210 Bulk MAR for each site was calculated as the product of sedimentation rate (from the age/depth profiles) and the dry bulk
density to yield the total mass of sediment deposited over a given age interval (bulk MAR). Dry bulk density was estimated by
correlating discrete physical properties data from each drill site to the reported GRA wet bulk density data. We used the
correlation to develop a higher resolution dry bulk density profile from the GRA data.

4 Results

4.1 Organic Carbon Variations:

215 There is a striking difference between the the C_{org} records from the Pacific equatorial sites (U1338, U1337, 574, 806 and 807)
older than ~13 Ma and the younger sections, as illustrated by Figure 2. Low sedimentary C_{org} %, C_{org} MAR, and C_{org}/Ba within
the MCO changes to higher and more scattered values between 14 and 12 Ma, after the MCO. C_{org} wt.% was much lower in the
older record as compared to the Pliocene and Pleistocene (Fig 2A). Within the MCO, between 17 and 14 Ma, C_{org} wt.% in these 5
220 sites averaged 0.043 ± 0.014 %, in contrast to a nearly 3-fold increase (0.124 ± 0.058 %) in the Pleistocene and Pliocene (0-4
Ma) and we note the MCO values are more similar to the average measured for Eocene equatorial Pacific (C_{org} roughly 0.03
wt%, >34 Ma; Olivarez Lyle and Lyle, 2006). Both the concentration of sedimentary C_{org} and the variability were much lower in
the period prior to 14 Ma.



225 High dilution by other sedimentary components can lower C_{org} %, but dilution in the equatorial Pacific requires higher preservation and burial of either biogenic $CaCO_3$ or biogenic SiO_2 relative to modern conditions. Nonetheless, this possibility can be evaluated by calculating the C_{org} MAR which removes the dilution effect and is a measure of the actual burial flux. Fig 2B shows the C_{org} MAR data for the 5 equatorial Pacific sites. We note a relatively low and consistent C_{org} MAR between 18 and 14 Ma. Somewhat higher levels of C_{org} MAR occur prior to 19 Ma and higher but very scattered MAR after 14 Ma.

230

4.2 Variations in Biogenic Barium

Ba was measured at 3 of the sites, all in the high-productivity region of the eastern Pacific. The C_{org}/Ba ratio compares the burial of the more labile C_{org} (<1% C_{org} preserved) to that of the better-preserved biogenic Ba (~30% preservation, Dymond et al, 1992, Dymond and Lyle, 1994). Since Ba has a linear relationship with C_{org} in the particulate rain of the modern equatorial Pacific Ocean (Dymond and Collier, 1996), changes in the C_{org}/Ba ratio is interpreted to reflect the relative preservation of C_{org} . Fig 2C shows that there was lower C_{org} burial relative to Ba prior to 14 Ma than later in the records.

235 Ba was measured by ICPMS at Site 574 for the sediment column older than 12 Ma (Piela et al, 2012). Sites U1337 and U1338 have Ba measured by scanning XRF for the entire composite sediment column (Lyle et al., 2012; Wilson, 2014, Lyle and
240 Baldauf, 2015). All have low C_{org}/Ba ratios (poor C_{org} preservation) prior to 14 Ma in the MCO, and higher ratios with much greater variation since then.




Figure 2: C_{org} % (A) C_{org} MAR (B) and C_{org}/Ba (C) for the equatorial sites 574, 806, 807, U1337, and U1338 to show evidence of low C_{org} contents and poor C_{org} preservation over the Miocene Climate Optimum (MCO, 17-13.8 Ma).



245

Low C_{org} % during the MCO is not confined to the equatorial Pacific but also found in the north Pacific. We also analyzed two Sites from the North Pacific to produce organic carbon records which show increased C_{org} burial as the Miocene progressed. At Sites 1208 on the Shatsky Rise, and Site 884 on Detroit Seamount (Fig 1), C_{org} MAR has increased strongly post MCO (Supplemental tables S8 and S10) A straightforward interpretation of these records, however, is complicated by the fact that there is a major increase in sedimentation rate at Site 1208 after a Paleocene hiatus which also causes increased C_{org} MAR post 7.5 Ma. As such, it is unclear to what extent the local sedimentation regime played a role in the increase of C_{org} MAR relative to regional levels of preservation. At Site 884, sedimentation rates increase in the early Miocene, leading to higher C_{org} MAR after the MCO (see supplemental material Fig S5). There is no reason to suspect that the Site 884 record was strongly influenced by changes in the local sedimentary environment, however, there is overall much higher clay deposition here that might have affected C_{org} burial. In summary, these two Sites suggest a general increase in C_{org} MAR throughout the Pacific after the MCO, but more records need to be produced and evaluated to support the hypothesis. However, it is important to highlight the data from a recent study by Li et al (2023) who found surprisingly low C_{org} burial across the MCO worldwide, in support of the hypothesis that a global slowdown in C_{org} deposition occurred during the MCO.

250

255

260

5 Discussion and Implications

We expect variability in the deposition of biogenic particulate matter resulting from large scale tectonic-biogeographic processes and from global intervals of high productivity, as well as from regional variation in primary productivity. We note that C_{org} depositional variability from 12 to 0 Ma is high, between 0.03 and 0.40 wt.%. The variation is partly caused by the geographic position of the site relative to the high equatorial productivity zone, and partly from the presence of high productivity intervals since the MCO. Some may also result from protection of C_{org} by the deposition of other sedimentary components. Nevertheless, we note that more slowly accumulating sediments of the Pliocene and Pleistocene tend to have high C_{org} % (Figs 3 and 4) but without corresponding high biogenic MARs. In other words, factors other than high productivity have caused better preservation of C_{org} between 8 and 0 Ma since high C_{org} % is not necessarily found with other indicators of high paleoproductivity. All records in the equatorial Pacific show evidence for low C_{org} % and C_{org} MAR during the MCO (Fig 2), while sediments from the Pliocene and Holocene have relatively high C_{org} % even though many of the sites exhibit low C_{org} MAR.

265

270

We interpret the scatter to reflect the localized response to the timing of productivity drivers combined with better C_{org} preservation in more recent times. There are 2 major productivity drivers in the equatorial Pacific: First, when the site is carried across the equatorial high productivity region by tectonic motion of the Pacific plate (line of high productivity for the Neogene; Lyle, 2003, Moore et al., 2004), and second, during high productivity intervals which typically are limited in both space and time. One example is the late Miocene Biogenic Bloom, between 8 and 4.5 Ma (Dickens and Owen; 1999; Diester-Haas et al, 2002; Lyle and Baldauf, 2015; Karatsolis et al., 2022; Gastaldello et al., 2023), and an earlier high productivity interval between 13 and 10.5 Ma (Lyle and Baldauf, 2015) and extending to ~14 Ma (Holbourn et al., 2014).

275

280

5.1 Tectonic passage through the equatorial productivity zone and equatorial primary productivity

Modern studies of equatorial biological productivity, including direct measurements in the water column, MAR in surface sediments, and those based on interpretations of satellite color find that the particulate flux both at the surface and to sediments is highest at the equator and strongly decreases to the north and south (Wyrski, 1981; Chavez and Barber, 1987; Dugdale et al., 1992; Murray and Leinen, 1993; Honjo et al., 1995; Behrenfeld et al, 2005). This pattern of high equatorial biogenic flux has




285 been found throughout the Neogene (Moore et al., 2004, Berger, 1973). The records are strong evidence that the equatorial
divergence driven by the SE trade winds crossing the equator has been a persistent feature of the Cenozoic oceans that causes
high primary productivity at the equator. The magnitude of equatorial productivity has not remained constant through time,
however, but has waxed and waned along with global climate change. We expect to find a change to higher productivity and
deposition of biogenic sediments as the movement of the Pacific Plate brings a drill site into position at the equator, and a
290 decrease in biogenic MARs as the site is moved away from the equator.

For example, Piela et al (2012) found that high biogenic silica mass accumulation rate, bio-Si MAR, and barium mass
accumulation rate, Ba MAR, occurred during the Site 574 paleo-equator crossing at 16.25 Ma, despite low C_{org} MAR and low
 $CaCO_3$ MAR. They hypothesized that the dissolution of $CaCO_3$ exposed more C_{org} in surface sediments to potential degradation
and reduced C_{org} with respect to other productivity signals. At site U1338, high burial rates of biogenic components other than
 C_{org} coincide with the period that Site U1338 was within $\pm 0.5^\circ$ of the paleo-equator during the middle Miocene (= 55 km
distance, 16-10 Ma, Fig 3). C_{org} MAR, unlike the other biogenic components, was not as enhanced during the MCO, implying
that C_{org} burial was minimized during the MCO and immediately after. Also, at Site U1338, the high C_{org} MAR and opal MAR at
12 Ma are roughly equivalent to modern MARs in surface sediments as reported by Murray and Leinen (1993). $CaCO_3$ MAR is
300 not only affected by high productivity but also by changes in carbonate dissolution through time. Nevertheless, much of the
variation in $CaCO_3$ at Site U1338 is common with variations in both opal and $BaSO_4$ MARs, indicating a strong productivity
signal at the site.

The site U1337 oxygen isotope record is the 15-20 Ma part of the Westerhold et al. (2020) Cenozoic oxygen isotope splice and
thus, along with Site U1338, represents the MCO interval in the equatorial Pacific. The sediments from Site U1337 (Fig 4)
305 crossed the equator from the south-east to north-west at an earlier time than Site U1338 (from 21 to 15 Ma versus 16 to 10 Ma,
respectively) and had relatively low biogenic MARs in the older sediments as compared to later in the U1337 section. This is in
part caused by sediment focusing in the younger part of the record, especially in the intervals 6.2-5.4 Ma and 4.5-3 Ma (Lyle et
al., 2019, and its supplemental material). The sediment focusing is shown by anomalously high sedimentation rates for sediments
at this latitude in the Pleistocene and by erosional channels to the northeast of the site. Nevertheless, sediment focusing has not
310 affected the time interval spanning the MCO for which we find very low C_{org} % and C_{org} MAR. Such low values contrast with
the expected pattern of higher biogenic sedimentation beneath the high productivity region of the paleo-equator. We also note
that Bio-Barium MARs at Site 1337 are 3 times higher than at Site 1338 during the MCO, implying greater primary productivity
and organic carbon production, yet this signal was not preserved in the sediments.

315 Sites 806 and 807, in the western Pacific on Ontong Java Plateau, crossed the equatorial region at <1 Ma and 10 Ma,
respectively. In the modern ocean, the upwelling signal in the far western equatorial Pacific is not nearly as strong as in the
eastern equatorial Pacific (Behrenfeld et al., 2005; Rousseaux and Gregg, 2017). Site 806 has its highest C_{org} MAR in the
Pleistocene as expected by its recent equator crossing. However, Site 807 shows little sign of its equator crossing in C_{org} MAR at
320 10 Ma, although it and Site 806 have a high C_{org} MAR resulting from the Late Miocene Biogenic Bloom. However, both sites
have low C_{org} MAR during the warm MCO.




325 **Figure 3:** A) C_{org} wt %, B) C_{org} MAR, C) $BaSO_4$ MAR, D) Biogenic SiO_2 MAR, and E) $CaCO_3$ MAR time series for Site U1338, on ocean crust formed at 18 Ma.. High biogenic MARs are characteristic of later high productivity intervals. However, the MCO has low C_{org} contents and C_{org} MAR, despite Site U1338 being relatively near the equator at that time. Paleo-equator line marks the time when plate tectonic movement aligned the site with the equator.

5.2 Past high productivity intervals

330 Figure 3 has time series of all biogenic MAR components at Site U1338. Note the interval from roughly 14.5 Ma to 11.5 Ma



indicates high primary production despite having low C_{org} burial within the MCO. High productivity during this interval at Site U1338 was first noted by Holbourn et al. (2014), for the period near 14 Ma, suggested that enhanced equatorial Pacific biogenic silica production was driven by changes in insolation which helped to draw down high atmospheric CO_2 and cause cooling at the



335

Figure 4: A) C_{org} wt %, B) C_{org} MAR, C) $BaSO_4$ MAR, D) Biogenic SiO_2 MAR, and E) $CaCO_3$ MAR time series for Site U1337, to the west of Site U1338, on 24 Myr ocean crust. The biogenic MAR time series are more complex here because of sediment focusing in the younger part of the record. However, the MCO has low C_{org} contents and C_{org} MAR, despite Site U1337 being relatively near the equator at that time. Paleo-equator marks the time when plate tectonic movement aligned the site with the equator.

340



the end of the MCO. In addition, there is a well-documented global productivity interval between 8 and 4.5 Ma found globally as the “Late Miocene Biogenic Bloom”, (LMBB) (Dickens and Owen, 1999; Diester-Haas et al, 2002; Lyle and Baldauf, 2015; Drury et al, 2017; Lyle et al, 2019; Karatsolis et al, 2022.) Both episodes are clearly observed at Site U1338. At Site U1337, there is similar levels of biogenic Si MAR to Site U1338 during the older interval but there is another interval filled with laminated diatom mats between 10 and 12 Ma (200-250 m CCSF) that might represent accumulation near the subduction boundary between the South Equatorial Current and North equatorial countercurrent (Exp 320/321 Scientific Party, 2010b). So, at Site U1337, the two separate higher productivity intervals are joined by a third interval of higher deposition. Both Sites 806 and 807 show elevated C_{org} MAR associated with the LMBB indicating that these global high productivity intervals affect all sites in the equatorial Pacific.

350

5.3 Four Hypotheses to explain low C_{org} burial at the MCO

We propose four working hypotheses for the cause of low levels of C_{org} MAR along the Pacific equator during the MCO: (1) generally low primary productivity during the MCO, (2) warmer deep waters and increased C_{org} degradation in the lower water column or sediment surface, (3) increased degradation of C_{org} in surface waters; and (4) low apparent C_{org} as an artifact of changing proxy relationships, e.g., Ba fixation in relation to C_{org} fixation in particles. In only one hypothesis (1, low primary productivity) does the relationship between C_{org} production, production of productivity proxies and their ultimate burial remain the same; the others assume a change in these relationships occurred during the MCO.

355

5.3.1 Hypothesis (1): Low productivity is the primary factor for low C_{org} during the MCO

Under hypothesis (1), the observed low C_{org} % during the MCO reflects low average productivity during the MCO interval. However, the data do not support this hypothesis. Each of the sites were in a different position relative to the equatorial high productivity zone at the time of the MCO. Those near the equator had as low a C_{org} % and C_{org} MAR as those farther away. Furthermore, there is no evidence of global low productivity during the MCO. Site U1338 which was in the equatorial zone late in the MCO, has increases in C_{org} % and C_{org} MAR at the end of the MCO. Since other paleoproductivity indicators tended to be high during the MCO it is apparent that low C_{org} MAR is not a result from low MCO primary productivity.

365

In summary, while we find some evidence for somewhat lower productivity during the MCO, high primary productivity associated with drill site equator crossings are reflected in increased burial of other biogenic sediment components, but not for C_{org} . Therefore, we conclude that additional factors caused the lower C_{org} MAR during the MCO.

370


5.3.2 Hypothesis (2): warmer deep waters and more C_{org} degradation in the lower water column or sediment surface led to low C_{org} MAR during the MCO

Particulate matter falls through the oceans at a rate of ~100 m/day (Honjo et al, 1982) meaning that it spends about 10 days above 1000 m in the upper water column and about a month in the deep waters below before reaching bottom, assuming an average ocean 4 km water depth. In addition, particulate C_{org} spends decades to centuries at the sediment surface before final burial. Provided that particulate C_{org} has survived the passage through the upper water column, it is possible that temperatures or other processes in the lower water column have controlled burial of C_{org} by temperature dependent degradation of C_{org} . If the change in C_{org} degradation occurs primarily in the lower water column, one consequence is that the biological pump would still function, albeit at a somewhat lower rate.

380



There are two problems with the hypothesis for lower water column control of C_{org} burial—first, most C_{org} is remineralized in the upper water column in the modern oceans, so only the more recalcitrant C_{org} fractions or those protected in some way survive to abyssal depths. Second, the cooling of the abyssal Pacific occurs much later than the end of the MCO (Fig 5). Any temperature-linked degradation of C_{org} in deep waters should decrease only after the deep waters cool. As Fig 5 illustrates, Pacific deep waters were only slightly warmer within the MCO than immediately after that period and stayed at a relatively constant temperature of 7° C (Lear et al, 2015) until about 6 Ma. After 6 Ma deep waters cooled gradually to a modern temperature of about 2°C. Therefore, we expect a C_{org} signal caused primarily by deep water cooling to have a much different signal than that observed.



390 **Figure 5:** C_{org} wt % time series in the equatorial Pacific along with bottom water temperature (BWT, Lear et al, 2015, from Site 806) and a stacked temperate water sea surface temperature (SST, Herbert et al., 2022) expressed as anomaly from modern. We show the temperate SST anomaly record because the data are based on alkenone paleotemperatures, and older equatorial SST is higher than temperatures that can be measured by alkenones. BWT in the Pacific did not change much until after 6 Ma, too late to affect C_{org} degradation in the MCO.

395

The abyssal temperature change during the MCO is relatively small. Lear et al. (2015) data show that the MCO was 0.4°C warmer than the next 3 Myr of the temperature record at Site 806 (Fig 5). Kochann et al. (2017), in contrast, describe an abyssal temperature change of 1.7° at the end of the MCO at Site U1338 and 2.2°C at Site 1146 in the South China Sea. However, they did not observe consistently warm temperatures through the intervals in the MCO that they measured, in contrast to the consistently low C_{org} contents. Kochann et al. (2017) noted a transient 2.6°C warming of bottom water temperatures at the onset of peak MCO warmth (15.5 Ma) at sites U1337 and U1338. However, both Lear et al (2015) and Kochann et al. (2017) found intermittent warming during the MCO, thus inconsistent with abyssal warmth being a driver of the low C_{org} over the entire MCO.

400

A low percentage of particulate matter derived from total productivity reaches the sea floor, and sediment trap studies have shown that the particulate flux that reaches the sea floor is relatively rich in C_{org} . The deepest sediment traps in the Joint Global Ocean Flux Study (JGOFS) tropical Pacific Experiment caught a particulate flux that contains around 5 wt % C_{org} (Honjo et al., 1995; particulate rain from sediment traps between 2191 m and 3618 m depth within $\pm 5^\circ$ of the equator), which, subsequently, must degrade down to the typical content found in surface equatorial sediments, around 0.2 to 0.3% (Murray and Leinen, 1993; Pahl et al, 1989). This order of magnitude loss of organic carbon, from the deepest waters to surface sediment, appears to be a

405



410 consistent level of degradation in the pelagic equatorial Pacific, as is also shown by core top values of C_{org} in the drill site data. Interestingly, highest C_{org} found in surface sediments measured in the JGOFS transect by Murray and Leinen (1993) are in off-equatorial sites, where clay is much more abundant and sedimentation rates are much lower. The lower equatorial C_{org} is partly a dilution effect by additional $CaCO_3$ compensated by higher sedimentation rates near the equator (Murray and Leinen, 1993).

415 If high abyssal temperature were a major factor in the C_{org} degradation, one might expect a major change in temperature associated with the MCO. Instead, there is only a minor abyssal temperature change at the end of the MCO (Fig 5; Lear et al, 2015). The major temperature decline occurred between 6 and 3 Ma associated with the end of the Miocene and Pliocene. Between 14 and 6 Ma, the BWT stayed between 6 and 7 °C, relatively constant, before dropping to 3° at about 2.5 Ma.

420 5.3.3: Hypothesis (3): warmer surface waters and more C_{org} degradation in the upper water column during the MCO

Our preferred hypothesis is that higher degradation in surface waters during the MCO reduced C_{org} burial via diminished rates of C_{org} transferred to deep waters, while post-MCO cooling led to increased opportunities for greater C_{org} transfer to the abyss, partly dependent on the productivity regime over each drill site (Fig 5, Herbert et al, 2022). Note, the Herbert et al (2022) data is expressed as an anomaly, relative to modern SST, so that multiple sites that remained below alkenone saturation could be
425 combined into one profile.

Fig 5 illustrates the large drop, roughly 9°C, in Northern Hemisphere mid-latitude SST immediately following the MCO. Because the SST data from Herbert et al. (2022) are based on alkenones, and the equatorial SSTs prior to 12 Ma were near or above the maximum temperature that alkenones record (~29°C, Rouselle et al., 2013), the midlatitude Northern Hemisphere SST
430 anomaly change is displayed in Figure 5. The tropical temperature change is probably smaller than that of the NH midlatitudes and may be confined more to the surface. Matsui et al (2017), found a stronger oxygen isotope gradient between surface and subsurface foraminifera during the MCO and proposed that the increased gradient resulted from warmer surface waters in the equatorial Pacific relative to subsurface waters. Using typical oxygen isotope change with temperature, the tropical surface
435 waters during the MCO were about 3°C warmer than the subsurface, even if these subsurface waters also warmed because of a deeper thermocline. We expect that surface equatorial waters should have warmed less than those of higher latitudes because of the ease of heat loss in the tropics through evaporation from warm water to the air.

There is no direct evidence of higher C_{org} degradation in surface waters, but modeling and observations of plankton distribution point to a loss of C_{org} primarily within the surface ocean layers, above 1000 m. Supplemental material from Boscolo-Galazzo et al. (2021) showed that there were much steeper $d^{13}C$ depth gradients in older time intervals, which modeled to a much shallower
440 and sharper O_2 minimum than in the Holocene. Using their temperature dependent model, the Holocene flux of particulate C_{org} at equatorial Pacific Site U1338 was 3 to 4 times greater at 600 m relative to 15 Ma for the same level of primary productivity. Similarly, we observe a factor of 3 to 4 increase in C_{org} sediment content over this same time interval at Site U1338. If this continued through the water column, a much smaller C_{org} flux was sequestered in abyssal waters during the warm MCO climate
445 relative to modern conditions. Also, it supports a hypothesis that sedimentary C_{org} contents are roughly proportional to the C_{org} particulate rain escaping the surface ocean.

However, C_{org} content of particulate rain that arrives at the abyssal seafloor is significantly larger than the C_{org} buried in the surface sediments, as noted previously. In the modern ocean, the particulate rain captured in deep sediment traps within the



450 equatorial Pacific region ($\pm 5^\circ$ of the equator) contains about 5% C_{org} (Table 5 of Honjo et al., 1995) versus the surface sediments that have a C_{org} concentration between 0.23 and 0.33% (Murray and Leinen, 1993). Lower C_{org} MAR away from the equator reflects the lower particulate rain rates away from the equator, and not the lower C_{org} contents in the particulate rain. Differences in $CaCO_3$ rain versus bio- SiO_2 rain were observed but did not strongly affect the total C_{org} preserved. In the Holocene, productivity apparently affects the rate of deposition of the particulate rain, but not so much its composition. This also supports
455 the role that the upper water column plays to determine both the magnitude of the biological pump and the level of C_{org} content in sediments.

We note that in periods after the MCO, there is an increase in the ratio of C_{org} to Ba during periods already identified as high productivity intervals. At Site U1338, where we have the best record, high C_{org}/Ba reflects periods when productivity was
460 relatively high globally, e.g., the late Miocene Biogenic Bloom (Dickens and Owen; 1999; Diester-Haas et al, 2002; Lyle and Baldauf, 2015; Drury et al, 2017; Lyle et al, 2019; Karatsolis et al, 2022).

5.3.4 Hypothesis (4): A change in proxy relationships for productivity, changing estimates of paleoproductivity

Another hypothesis worth considering is that during the MCO there is a change in the response of the proxy that varies from the expected modern response. For example, diatom deposition or C_{org}/Ba ratio might behave differently with respect to C_{org}
465 production. Under these conditions there could be lower actual C_{org} export to the interior ocean than that indicated by the proxy, minimizing the deposition of C_{org} without indicating lower productivity. We believe that there is some likelihood that the relationship between export of particulate C_{org} from the euphotic zone to other biogenic components may be somewhat different under warm earth conditions but propose that these differences result primarily from relative changes in C_{org} consumption in the
470 upper water column.

Dymond and Collier (1996) described how C_{org} rained out of the modern equatorial Pacific relative to Ba. They found lower C_{org} particulate rain away from the equator, corresponded to much lower ratio of C_{org} to Ba (~ 30). In contrast, this ratio is ~ 150 near the equator where C_{org} rain was high. The data suggests relatively rapid formation of Ba in microenvironments within particulate
475 rain, followed by a loss of C_{org} versus Ba. There is more complete consumption of C_{org} where the C_{org} flux is lower. The sediment C_{org} to Ba ratios found in sediments (Fig 2) are much lower because of the high degradation of C_{org} at the sea floor relative to Ba ($>20x$ reduction for C_{org} vs $\sim 3x$ for Ba) before burial. However, the amount of Ba fixed in micro-environments depends on the Ba composition of seawater. In the modern ocean, Ba rain is significantly lower in the Atlantic than in the Pacific relative to C_{org} because of the lower dissolved Ba in the Atlantic (Dymond and Collier, 1996).

480 If the modern observations across the equatorial region are consistent with changes that might occur in a warm interval like the MCO, we expect lower C_{org} in the particulate rain relative to Ba. Conceivably then, the C_{org} rain might slow even though the Ba flux did not. This could happen if dissolved Ba is fixed into barite ($BaSO_4$) relatively early in the rain of particulates, so that later degradation of C_{org} only affects the C_{org}/Ba leaving surface waters (Fig 3). However, the C_{org} MAR resembles the opal MAR to a
485 certain extent after the end of the MCO, indicating that the C_{org} MAR has a profile like a different proxy for productivity when surface waters cool. Perhaps the presence of diatoms causes a more effective transport of particulate C_{org} to the sea floor.

In time series at Site U1338 we find significant non-random changes in the ratio of C_{org} to Ba, associated with apparent changes in productivity (Fig 3). Specifically: high C_{org}/Ba associated with the late Miocene Biogenic Bloom, and during a period around



490 12 Ma that shows high biogenic Si and CaCO₃ deposition as well. These time series show that the biogenic components have their individual processes that lead from creation to burial, so that care needs to be taken to quantitatively ascribe a certain level of primary productivity to the remains found in the sediments. Nevertheless, the lack of C_{org} response during the MCO likely results from upper water column processes.

495 6 Conclusions

In earlier work we have found that warm earth conditions in the Eocene are marked by very low levels of C_{org} burial (Olivarez Lyle and Lyle, 2005, 2006). In this study we also show that warm earth conditions during the Miocene Climate Optimum are also characterized by a low level of C_{org} burial compared to later in the sedimentary record. The low levels are represented in both the weight % C_{org} and C_{org} MAR's, and as low ratios of C_{org} relative to other better preserved biogenic components like BaSO₄, despite relatively high deposition of other paleoproductivity proxies like biogenic silica. We formed 4 hypotheses to explain the low C_{org} at the MCO: lower productivity, higher degradation in the lower water column, higher degradation in the upper water column, and a change in relationships between proxies, and rejected all except higher degradation the upper water column.

500 We have identified the upper water column as the region where C_{org} is preferentially removed from the particulate rain to the sea floor, an indication of a 'short circuit' in the biological pump under extreme global warmth once the ocean equilibrates. We observe that the average C_{org} content of equatorial Pacific MCO sediments was about 0.04 % organic carbon, 5 times lower than modern surface sediment of 0.2 to 0.3% C_{org} (Murray and Leinen, 1993). If preserved C_{org} in surface sediments is roughly proportional to the rain of C_{org} that reaches the sea floor, then about 5 times less C_{org} in the particulate rain reached the seafloor at the MCO, caused by higher metabolic degradation in surface waters.

505 While such a proportionality of C_{org} in particulate rain is an oversimplification of early diagenesis in pelagic environments, it is an example of how the pelagic sedimentary environment responds to warm earth conditions. Better diagenetic models under low sedimentation rates and oxic conditions might improve our ability to hindcast particulate rain in the past. Clearly, though, there is a sedimentary response to these processes in the water column.

515 Author Contributions

Mitchell Lyle helped to collect the cores from Sites U1337 and U1338. He organized the XRF analyses along the continuous sediment sections (splices) and found funding for the discrete calibration carbon samples. He had primary responsibility for writing the paper. Annette Olivarez Lyle trained students, supervised both the bio-SiO₂ and the carbon analyses (CaCO₃ and C_{org}) and was responsible for quality control of the results. She also helped write and edit the paper.

Competing Interests

The authors declare that they have no conflict of interest.

525

Acknowledgements

We thank the scientific party and crew of the *D/V JOIDES Resolution* on the 2009 Exp 320/321 IODP expedition for collection and initial description of Sites U1337 and U1338. We also thank the students and technical staff that ran the carbon analyses



530 (Chris Piela, Bianca Romero, Julia Wilson, Anna Stepanova). We also thank Joanna T. Lyle for reviewing the initial draft of this paper.

References

- Arrhenius, G. O. S.: Sediment cores from the east Pacific., in: Reports of the Swedish Deep Sea Expedition, 1947-1948, 5, 1-228, 1952.
- Behrenfeld, M. J., Boss, E. S., Siegel, D. A., and Shea, D. M.: Carbon-based ocean productivity and phytoplankton physiology from space, *Global Biogeochemical Cycles*, 19, GB1006, 10.1029/2004GB002299, 2005.
- 535 Berger, W. H.: Cenozoic sedimentation in the eastern tropical Pacific, *Geological Society of America Bulletin*, 84, 1941-1954, 1973.
- Boscolo-Galazzo, F., Crichton, K. A., Ridgwell, A., Mawbey, E. M., Wade, B. S., and Pearson, P. N.: Temperature controls carbon cycling and biological evolution in the ocean twilight zone, *Science*, 371, 1148-1152, 2021.
- Boyd, P. W.: Toward quantifying the response of the oceans biological pump to climate change, *Frontiers in Marine Science*, 2, Article 77, 10.3389/fmars.2015.00077, 2015.
- 540 Bralower, T.J., Premoli Silva, I., Malone, M.J., et al. *Proc. ODP, Init. Repts.*, 198: College Station, TX (Ocean Drilling Program), 2002. doi: 10.2973/odp.proc.ir.198.2002
- Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., and West, G. B.: Toward a metabolic theory of ecology, *Ecology*, 85, 1771-1789, 2004.
- 545 Chavez, F. P. and Barber, R. T.: An estimate of new production in the equatorial Pacific, *Deep-Sea Research*, 34, 1229-1243, 1987.
- Dehairs, F., Chesselet, R., and Jedwab, J.: Discrete suspended particles of barite and the barium cycle in the open ocean, *Earth and Planetary Science Letters*, 49, 528-550, 1980.
- Dickens, G. R. and Owen, R. M.: The latest Miocene-early Pliocene biogenic bloom: a revised Indian Ocean perspective, *Marine Geology*, 161, 75-91, 1999.
- 550 Diester-Haass, L., Billups, K., and Emeis, K.-C.: In search of the late Miocene-early Pliocene “biogenic bloom” in the Atlantic Ocean (Ocean Drilling Program Sites 982, 925, and 1088), *Paleoceanography*, 20, PA4001, 10.1029/2005PA001139, 2005.
- Drury, A. J., Westerhold, T., Frederichs, T., Tian, J., Wilkens, R., Channell, J. E. T., Evans, H., John, C. M., Lyle, M., and Röhl, U.: Late Miocene climate and time scale reconciliation: Accurate orbital calibration from a deep-sea perspective, *Earth and Planetary Science Letters*, 475, 254-266, 10.1016/j.epsl.2017.07.038, 2017.
- 555 Drury, A. J., Lee, G. P., Gray, W. R., Lyle, M., Westerhold, T., Shevenell, A. E., and John, C. M.: Deciphering the state of the late Miocene to early Pliocene equatorial Pacific, *Paleoceanography and Paleoclimatology*, 33, 246-263, 10.1002/2017PA003245, 2018.
- Dugdale, R. C., Wilkerson, F. P., Barber, R. T., and Chavez, F. P.: Estimating new production in the equatorial Pacific Ocean at 150°W, *Journal of Geophysical Research*, 97, 681-687, 1992.
- Dymond, J., Suess, E., and Lyle, M.: Barium in deep-sea sediment: A geochemical proxy for paleoproductivity, *Paleoceanography*, 7, 163-181, 1992.
- 560 Dymond, J. and Lyle, M.: Particle fluxes in the ocean and implications for sources and preservation of ocean sediments, in: *Material Fluxes on the Surface of the Earth*, edited by: Hay, W. W., Andrews, J. T., Baker, V. R., Dymond, J., Kump, L. R., Lerman, A., Martin, W. R., Meybeck, M., Milliman, J. D., Rea, D. K., and Sayles, F. L., National Academy Press, Washington DC, 125-143, 1994.
- Dymond, J. and Collier, R.: Particulate barium fluxes and their relationships to biological productivity, *Deep Sea Research II*, 43, 1283-1308, 1996.
- 565 Evans, H. F.: *Magnetic Stratigraphy and Environmental Magnetism of Oceanic Sediments*, Geology, PhD. University of Florida, 204 pp., 2006.



- Expedition 320/321 Scientists, 2010a. Methods. In Pälike, H., Lyle, M., Nishi, H., Raffi, I., Gamage, K., Klaus, A., and the Expedition 320/321 Scientists, *Proc. IODP*, 332/321: Tokyo (Integrated Ocean Drilling Program Management International, Inc.).
570 doi:10.2204/iodp.proc.320321.102.2010
- Expedition 320/321 Scientists, 2010b. Site U1337. In Pälike, H., Lyle, M., Nishi, H., Raffi, I., Gamage, K., Klaus, A., and the Expedition 320/321 Scientists, *Proc. IODP*, 332/321: Tokyo (Integrated Ocean Drilling Program Management International, Inc.).
doi:10.2204/iodp.proc.320321.109.2010
- Ganeshram, R. S., Francois, R., Commeau, J., and Brown-Leger, S. L.: An experimental investigation of barite formation in seawater,
575 *Geochimica et Cosmochimica Acta*, 67, 2599-2605, 2003.
- Gastaldello, M. E., Agnini, C., Westerhold, T., Drury, A. J., Sutherland, R., Drake, M. K., Lam, A. R., Dickens, G. R., Dallanave, E., Burns, S., and Alegret, L.: The Late Miocene-Early Pliocene Biogenic Bloom: An Integrated Study in the Tasman Sea, *Paleoceanography and Paleoclimatology*, 38, e2022PA004565, <https://doi.org/10.1029/2022PA004565>, 2023.
- Hedges, J. I. and Keil, R. G.: Sedimentary organic matter preservation: an assessment and speculative synthesis, *Marine Chemistry*, 49,
580 81-115, 1995.
- Herbert, T. D., Dalton, C. A., Liu, Z., Salazar, A., Si, W., and Wilson, D. S.: Tectonic degassing drove global temperature trends since 20 Ma, *Science*, 377, 116-119, [10.1126/science.abl4353](https://doi.org/10.1126/science.abl4353), 2022.
- Holbourn, A., Kuhnt, W., Lyle, M., Schneider, L., Romero, O., and Andersen, N.: Middle Miocene climate cooling linked to intensification of eastern equatorial Pacific upwelling, *Geology*, 42, 19-22, [10.1130/G34890.1](https://doi.org/10.1130/G34890.1), 2014.
- 585 Holbourn, A., Kuhnt, W., Kochhann, K. G. D., Andersen, N., and Meier, K. J. S.: Global perturbation of the carbon cycle at the onset of the Miocene Climate Optimum, *Geology*, 43, 123-126, [10.1130/G36317.1](https://doi.org/10.1130/G36317.1), 2015
- Honjo, S., Manganini, S. J., and Poppe, L. J.: Sedimentation of Lithogenic Particles in the Deep Ocean, *Marine Geology*, 50, 199-220, 1982.
- Honjo, S., Dymond, J., Collier, R., and Manganini, S. J.: Export production of particles to the interior of the equatorial Pacific Ocean during the 1992 EqPac experiment, *Deep-Sea Research*, 42, 831-870, 1995.
590
- John, E. H., Wilson, J. D., Pearson, P. N., and Ridgwell, A.: Temperature-dependent remineralization and carbon cycling in the warm Eocene oceans, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 413, 158-166, [10.1016/j.palaeo.2014.05.019](https://doi.org/10.1016/j.palaeo.2014.05.019), 2014.
- Karatsolis, B.-Th., Lougheed, B. C., De Vleeschouwer, D., and Henderiks, J.: Abrupt conclusion of the late Miocene-early Pliocene biogenic bloom at 4.6-4.4 Ma, *Nature Communications*, 13, 353, <https://doi.org/10.1038/s41467-021-27784-6>, 2022.
- 595 Kochann, K. G. D., Holbourn, A., Kuhnt, W., and Xu, J.: Eastern equatorial Pacific benthic foraminiferal distribution and deep water temperature changes during the early to middle Miocene, *Marine Micropaleontology*, 133, 28-39, DOI [10.1016/j.marmicro.2017.05.002](https://doi.org/10.1016/j.marmicro.2017.05.002), 2017.
- Kroenke, L.W., Berger, W.H., Janecek, T.R., et al., 1991. *Proc. ODP, Init. Repts.*, 130: College Station, TX (Ocean Drilling Program).
[doi:10.2973/odp.proc.ir.130.1991](https://doi.org/10.2973/odp.proc.ir.130.1991)
- 600 Li, Z., Zhang, Y. G., Torres, M., and Mills, B. J. W.: Neogene burial of organic carbon in the global ocean, *Nature*, 613, 90-95, <https://doi.org/10.1038/s41586-022-05413-6>, 2023.
- Lear, C. H., Coxall, H. K., Foster, G. L., Lunt, D. J., Mawbey, E. M., Rosenthal, Y., Sosdian, S. M., Thomas, E., and Wilson, P. A.: Neogene ice volume and ocean temperatures: Insights from infaunal foraminiferal Mg/Ca paleothermometry, *Paleoceanography*, 30, doi:10.1002/2015PA002833, 2015.
- 605 Lima, I. D., Lam, P. J., and Doney, S. C.: Dynamics of particulate organic carbon flux in a global ocean model, *Biogeosciences*, 11, 1177-1198, doi:10.5194/bg-11-1177-2014, 2014.



- Lyle, M., Mix, A., Ravelo, C., Andreasen, D., Heusser, L., and Olivarez, A.: Millennial-scale CaCO₃ and C-org events along the northern and central California margin: stratigraphy and origins, in: Proceedings of the Ocean Drilling Program, Scientific Results, edited by: Lyle, M., Koizumi, I., Moore, T. C., jr., and Richter, C., Ocean Drilling Program, College Station TX, 163-182, 2000.
- 610 Lyle, M.: Neogene carbonate burial in the Pacific Ocean, *Paleoceanography*, 18, DOI 10.1029/2002PA000777, 2003.
- Lyle, M. W., Olivarez Lyle, A., Backman, J., and Tripathi, A.: Biogenic sedimentation in the Eocene equatorial Pacific: the stuttering greenhouse and Eocene carbonate compensation depth, in: Proceedings of the Ocean Drilling Program, Scientific Results, Leg 199, edited by: Lyle, M., Wilson, P., Janecek, T. R., and Firth, J., Ocean Drilling Program, College Station TX, 10.2973/odp.proc.sr.199.219.2005, 2005.
- 615 Lyle, M., Olivarez Lyle, A., Gorgas, T., Holbourn, A., Westerhold, T., Hathorne, E. C., Kimoto, K., and Yamamoto, S.: Data report: raw and normalized elemental data along the U1338 splice from X-ray Fluorescence scanning, Proceedings of the Integrated Ocean Drilling Program, 320/321, 10.2204/iodp.proc.320321.2010, 2012.
- Lyle, M. and Baldauf, J.: Biogenic sediment regimes in the Neogene equatorial Pacific, IODP Site U1338: Burial, production, and diatom community, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 433, 106-128, 10.1016/j.palaeo.2015.04.001, 2015.
- 620 Lyle, M., Drury, A. J., Tian, J., Wilkens, R., and Westerhold, T.: Late Miocene to Holocene high-resolution eastern equatorial Pacific carbonate records: stratigraphy linked by dissolution and paleoproductivity, *Climate of the Past*, 15, 1715-1739, <https://doi.org/10.5194/cp-15-1715-2019>, 2019.
- Martin, J. H., Knauer, G. A., Karl, D. M., and Broenkow, W. W.: VERTEX: carbon cycling in the northeast Pacific, *Deep-Sea Research*, 34, 267-285, 1987.
- 625 Matsui, H., Nishi, H., Kuroyanagi, A., Hayashi, H., Ikehara, M., and Takashima, R.: Vertical thermal gradient history in the eastern equatorial Pacific during the early to middle Miocene: Implications for the equatorial thermocline development, *Paleoceanography*, 32, 729-743, doi:10.1002/2016PA003058, 2017.
- Mayer, L. A., Theyer, F., Barron, J., Dunn, D. A., Handyside, T., Hills, S., Jarvis, I., Nigrini, C. A., Pisias, N., Pujos, A., Saito, T., Stout, P., Thomas, E., Weinrich, N., and Wilkens, R. H.: Initial Reports of the Deep Sea Drilling Program, Leg 85, U.S. Gov't Printing Office, Washington, 1021 pp.1985.
- 630 Mayer, L. M.: Sedimentary organic matter preservation: an assessment and speculative synthesis-a comment, *Marine Chemistry*, 49, 123-126, 1995.
- Mayer, L. A.: Extraction of high-resolution carbonate data for palaeoclimate reconstruction, *Nature*, 352, 148-150, 1991.
- Moore, T. C. Jr., Backman, J., Raffi, I., Nigrini, C., Sanfilippo, A., Palike, H., and Lyle, M.: The Paleogene tropical Pacific: Clues to circulation, productivity, and plate motion, *Paleoceanography*, 19, doi:10.1029/2003PA000998, 2004.
- 635 Muller, P. J. and Suess, E.: Productivity, sedimentation rate, and sedimentary organic matter in the oceans -I. Organic carbon preservation, *Deep-Sea Research*, 26A, 1347-1362, 1979.
- Murray, R. W. and Leinen, M.: Chemical transport to the seafloor of the equatorial Pacific Ocean across a latitudinal transect at 135°W: tracking sedimentary major, trace, and rare earth element fluxes at the equator and the Intertropical Convergence Zone, *Geochimica et Cosmochimica Acta*, 57, 4141-4163, 1993.
- 640 Murray, R. W., Leinen, M., and Isern, A. R.: Biogenic flux of Al to sediment in the central equatorial Pacific Ocean: Evidence for increased productivity during glacial periods, *Paleoceanography*, 8, 651-671, 1993.
- Olivarez Lyle, A. and Lyle, M.: Organic carbon and barium in Eocene sediments: Possible controls on nutrient recycling in the Eocene equatorial Pacific Ocean, in: Proceedings of the Ocean Drilling Program, Scientific Reports Volume 199, edited by: Lyle, M., Wilson, P.A., Janecek, T.R., et al., Ocean Drilling Program, College Station TX, 1-33; http://www-odp.tamu.edu/publications/199_IR/chap_106/chap_106.htm, 2005.
- 645



- Olivarez Lyle, A. and Lyle, M.: Organic carbon and barium in Eocene sediments: Is metabolism the biological feedback that maintains end-member climates? *Paleoceanography*, 21, 1-13; doi:10.1029/2005PA001230, 2006.
- Pälike, H., Lyle, M., Nishi, H., Raffi, I., Gamage, K., Klaus, A., and the Expedition 320/321 Scientists, 2010. *Proc. IODP*, 320/321: Tokyo (Integrated Ocean Drilling Program Management International, Inc.). doi:10.2204/iodp.proc.320321.2010
- 650 Piela, C., Lyle, M., Marcantonio, F., Baldauf, J., and Olivarez Lyle, A.: Biogenic sedimentation in the equatorial Pacific: Carbon cycling and paleoproduction, 12-24 Ma, *Paleoceanography*, 27, PA2204, 2218 pp, 10.1029/2011PA002236, 2012.
- Prahl, F. G., Muehlhausen, L. A., and Lyle, M.: An organic geochemical assessment of oceanographic conditions at MANOP Site C over the past 26,000 years, *Paleoceanography*, 4, 495-510, 1989.
- 655 Rea, D.K., Basov, L.A., Janecek, T.R., Palmer-Julson, A., et al., 1993. *Proc. ODP, Init. Repts.*, 145: College Station, TX (Ocean Drilling Program).
- Rouselle, G., Beltran, C., Sicre, M.-A., Raffi, I., and De Rafelis, M.: Changes in sea-surface conditions in the Equatorial Pacific during the middle Miocene-Pliocene as inferred from coccolith geochemistry, *Earth and Planetary Science Letters*, 361, 412-421, 10.1016/j.epsl.2012.11.003, 2013.
- 660 Rousseaux, C. S. and Gregg, W. W.: Forecasting ocean chlorophyll in the equatorial Pacific, *Frontiers in Marine Science*, 4, 10.3389/fmars.2017.00236, 2017.
- Stax, R. and Stein, R.: 34. Long-Term Changes In The Accumulation Of Organic Carbon In Neogene Sediments, Ontong Java Plateau, *Proceedings of the Ocean Drilling Program, Scientific Results*, 130, 573-584, 1993.
- Suess, E.: Particulate organic carbon flux in the oceans - surface productivity and oxygen utilization, *Nature*, 288, 260-263, 1980.
- 665 Tian, J., Ma, X., Zhou, J., Jiang, X., Lyle, M., Shackford, J. K., and Wilkens, R.: Paleoceanography of the east equatorial Pacific over the past 16 Myr and Pacific-Atlantic comparison: High resolution benthic foraminiferal $\delta^{18}O$ and $\delta^{13}C$ records at IODP Site U1337, *Earth and Planetary Science Letters*, 499, 185-196, 10.1016/j.epsl.2018.07.025, 2018.
- van Andel, T. H., Heath, G. R., and Moore, T. C. jr.: *Cenozoic History and Paleoceanography of the central equatorial Pacific Ocean*, GSA Memoir 143, Geological Society of America, 133 pp.1975.
- 670 Westerhold, T., Marwan, N., Drury, A. J., Liebrand, D., Agnini, C., Anagnostou, E., Barnet, J. S. K., Bohaty, S. M., De Vleeschouwer, D., Florindo, F., Frederichs, T., Hodell, D. A., Holbourn, A. E., Kroon, D., Lauretano, V., Littler, K., Lourens, L. J., Lyle, M., Pälike, H., Röhl, U., Tian, J., Wilkens, R. H., Wilson, P. A., and Zachos, J. C.: An astronomically dated record of Earth's climate and its predictability over the last 66 million years, *Science*, 369, 1383-1387, 2020.
- Wilson, J. K.: Early Miocene carbonate dissolution in the eastern equatorial Pacific, PhD. Thesis, Oceanography, Texas A and M University, 155 pp., 2014.
- 675 Wyrтки, K.: An estimate of equatorial upwelling in the Pacific, *Journal of Physical Oceanography*, 11, 1205-1214, 1981.