

Strong volcanic-induced climatic shocks on historical Moselle wine production

Fredrik Charpentier Ljungqvist^{1,2,3}, Bo Christiansen⁴, Lea Schneider^{5,6}, and Peter Thejll⁴

¹Department of History, Stockholm University, 106 91 Stockholm, Sweden

²Bolin Centre for Climate Research, Stockholm University, 106 91 Stockholm, Sweden

³Swedish Collegium for Advanced Study, Linneanum, Villavägen 6c, 752 38 Uppsala, Sweden

⁴Danish Meteorological Institute, Sankt Kjelds Plads 11, 2100 Copenhagen Ø, Denmark

⁵Department of Geography, Justus-Liebig-University, 35390 Giessen, Germany

⁶Center for international Development and Environmental Research, Justus-Liebig-University, 35390 Giessen, Germany

Correspondence: Fredrik Charpentier Ljungqvist (fredrik.c.l@historia.su.se)

Abstract. In central and southern Europe, grapevine is a climate-sensitive agricultural product of great economic importance, both in historical times and today. We systematically investigated the climatic impact, focusing on volcanic-forced abrupt cooling, on two long annual records of wine production quantity (spanning 1444–1786) from the Moselle Valley in present-day Luxembourg, close to the northern limit of viticulture in Europe. We present a consistent picture of the impact of volcanic eruptions on wine production through climate. To this end, we applied superposed epoch analysis – an appropriate method for detecting episodic signals in non-stationary time-series – in combination with a bootstrap procedure to estimate the statistical significance. We also assessed the long-term relationship between different annual and seasonal climate parameters and wine production in the Moselle Valley. Robust and highly significant wine production declines occurred in the years immediately following major volcanic events. Warmer, and to a lesser extent drier, climate condition had a moderately strong, but persistent, positive effect on wine production. We also find a volcanic cooling signature in spring and summer in temperature reconstructions. However, the detected volcanic signature in the Moselle Valley wine production is considerably stronger than the one found for Central Europe in tree-ring-based reconstructions and is instead more akin to the strong volcanic signature present in Fennoscandian tree-ring series. On the basis of our findings, we encourage further compilation, publication, and analyses of additional wine production series containing unique biological and climatic information.

15 1 Introduction

The climatic response to volcanic forcing during the past few millennia has been studied intensively from local to global scales, typically using tree-ring data from strongly temperature-limited growth environments. The impacts on agricultural productivity, including on viticulture, in less extreme climatic settings remain far less well understood. The study of long time-series of agricultural productivity, like grapevine harvest quantities, offers insights into the effects of such abrupt cooling on biological records with properties possibly differing from those of tree-ring series. This, in turn, may shed new light on still unresolved

questions about the effects volcanic-induced cooling has on tree growth, in addition to gaining a better understanding of the role of major volcanic events for past agriculture.

25 Grapevine (henceforth ‘vine’) is a highly climate-sensitive crop whose phenology is comparatively well understood (Jones et al., 2005; Bock et al., 2013; de Cortazar Atauri et al., 2017; Drappier et al., 2019; Morales-Castilla et al., 2020; Droulia and Charalampopoulos, 2022). The biological requirements of both modern and historical European viticulture have been documented in detail (Lachiver, 1988; Mullins et al., 1992; Guerreau, 1995). An increasingly warmer climate, with longer growing seasons, benefits viticulture (vine growing) near its latitudinal and altitudinal limits (Malheiro et al., 2010; Guiot et al., 2023), whereas climatic warming at the same time can have adverse effects on viticulture in regions with a hot and dry Mediterranean climate (Cook and Wolkovich, 2016; Venios et al., 2020; Andrade et al., 2024; van Leeuwen et al., 2024). Grapes
30 were the most important cash crop, of an economic importance only second to grain, in much of Europe during the medieval and early modern periods, and vine is and was more climate-sensitive than grain (Pfister and Wanner, 2021). Relationships between different parameters associated with wine production and summer temperature in late medieval and early modern Europe have been previously described in the literature (Ladurie Le Roy and Baulant, 1980; Le Roy Ladurie et al., 2006; Labbé et al., 2019; Pfister et al., 2024). In Central Europe, the quantity of the grape harvest was positively correlated to June–July
35 temperature, the quality to July–September (and especially August) temperature, and the harvest dates to April–July/August temperature (Pfister, 1981; Pfister et al., 2024). Viticulture in Central Europe is, and was, especially sensitive to the effect of spring frost (Meier et al., 2018). The strong cooling during the climax of the Little Ice Age *c.* 1570–1710 (Wanner et al., 2022) made viticulture non-viable in parts of Central Europe – with considerable economic and social consequences (Landsteiner, 1999).

40 Large, but more short-lived, reductions in vine harvests – and thus wine production declines – could be triggered by sharp volcanic-induced cooling. Major volcanic eruptions have a significant impact on climate (Robock, 2000; Oppenheimer, 2011) through a negative radiative forcing related to an increased atmospheric loading of aerosols and dust, and can induce significant global to hemispheric scale cooling for one to several years (Stoffel et al., 2015; Sigl et al., 2015; Burke et al., 2023). The climate effect of an eruption does not only depend on its magnitude, but also on factors such as the geographical position of
45 the volcano and the season of the eruption (Guillet et al., 2017). Quantitative estimates of volcanic-induced cooling during pre-industrial times have mostly been derived from large-scale temperature reconstructions based on tree-ring data (Schneider et al., 2015; Stoffel et al., 2015; Wilson et al., 2016; Anchukaitis et al., 2017; Büntgen et al., 2021). In altitudinal and latitudinal tree-line ecotones, tree growth is particularly sensitive to summer temperature (Körner, 2021) and short-term climatic perturbations are archived in anomalies of annual ring width or wood density (Esper et al., 2016; Hartl-Meier et al., 2017). The precise annual
50 dating of tree-ring chronologies allows for a clear attribution of temperature deviations to volcanic eruptions (Anchukaitis et al., 2012; D’Arrigo et al., 2013; Esper et al., 2013a; Schneider et al., 2017).

The adverse impacts on agricultural productivity and, hence, on human society of volcanic-induced cooling in pre-industrial times have been well-documented in recent scholarship (D’Arrigo et al., 2020; Guillet et al., 2020; Huhtamaa et al., 2022; Stoffel et al., 2022; White et al., 2022; Martin et al., 2023). This is, in particular, the case at the northern edge of grain
55 agriculture in Europe (Ljungqvist et al., 2021, 2024). The effects of volcanic-induced cooling on viticulture in pre-industrial

Europe remains less well understood. However, it has been demonstrated that larger, especially tropical, volcanic eruptions resulted in significantly later grape harvest dates for one to two years after the volcanic event (Meier et al., 2007; Guillet et al., 2017; Pfister and Wanner, 2021). The effects of individual major volcanic forcing events on wine production have also been studied. For example, Brönnimann and Krämer (2016) studied the effects of the 1815 Tambora volcano eruption, followed by the cold summer of 1816 in central Europe. The extent to which the cooling was due to volcanic forcing was evaluated, as well as its biophysical effects (e.g., low harvest yields) and subsequent societal consequences.

Numerous records, not least from France and Switzerland, exist of grape harvest dates (Yiou et al., 2012). They have been proven to be strongly related to spring and summer temperatures, and have been utilised in numerous documentary-based temperature reconstructions (e.g., Chuine et al., 2004; Kiss et al., 2011; Možný et al., 2016a; Labbé et al., 2019), and also drought reconstructions (Možný et al., 2016b), despite possible biases in the long-term trends (Garcia de Cortazar-Atauri et al., 2010; Garnier et al., 2011). Grape harvest dates have been proven to be particularly skillful in capturing extremely cold years (Maurer et al., 2011), such as those frequently following volcanic forcing events (Sigl et al., 2015), although the skill of capturing extremely warm years are less good (Keenan, 2007). Likewise, a number of long wine price series exist from different portions of Europe (Allen and Unger, 2019), but the possible climate signal embedded in the wine price series remains relatively unexplored. In contrast to grape harvest dates and wine price series, long and continuous series of wine production (or grapevine harvest quantity or quality) are rare for medieval and early modern Europe (see, though, the recently published ones by Pfister et al., 2024). However, two long annual wine production quantity records, covering 1444–1786 with gaps, were already published by Yante (1985). They derive from Grevenmacher and Remich in the Moselle Valley, in present-day Luxembourg, close to the latitudinal limit of European viticulture (Fig. 1a–b).

In this article, we systematically investigate the impact from climate, with particular focus on the influence from volcanic forcing, on the two above-mentioned long wine production quantity series. We will first assess the effects of volcanic forced cooling on wine production using superposed epoch analysis (SEA) appropriate for studying episodic events in non-stationary time-series. Then, we will evaluate the effects of volcanic forcing on Moselle Valley pre-industrial climate and explore whether the wine production series contribute information useful for understanding volcanic-induced cooling in the region between the Alps and Scandinavia in comparison to temperature-sensitive tree-ring data. Finally, we will investigate both the short- and long-term relationship between different annual and seasonal climate parameters and wine production quantities in order to understand and constrain the effects of climate on wine production variability. Through utilising bootstrap and phase-scrambling-based techniques to estimate significance, we intend to pay particular attention to issues related to statistical significance.

We emphasise that this article is of a statistical nature. Our aim is to investigate the statistical relationships between volcanic forcing, climate and wine production in the Moselle Valley region. The rest of article is organised as follows: After presenting our data and methods (Section 2), we start with investigating the influence of volcanic eruptions on wine production (subsection 3.1), followed by the influence of volcanic eruptions on climate (subsection 3.2), and ending with the influence of climate extremes on the wine production (subsection 3.3). In Section 4 we discuss the interpretations and implications of the results and their associated uncertainties, starting with the volcanic and climatic signals embedded in the wine production series (sub-

90 section 4.1) followed by a comparative discussion about the volcanic forcing effects on wine production in relation to tree-ring growth (subsection 4.2). We conclude the article with a short conclusion and outlook (Section 5).

2 Materials and methods

In this section we start with describing the wine production data (Section 2.1), followed by the volcanic forcing dataset (Section 2.2), and then the (palaeo)climate series (Section 2.3). All the datasets are summarised in Table 2. We end the section with
95 describing the various statistical methods applied (Section 2.4).

2.1 The Moselle Valley wine production series

The two annual wine production quantity series from Grevenmacher and Remich in the Moselle Valley (49.5°N, 6.35°E) of present-day Luxembourg are based on official records of a tax of one-ninth levied on the wine production. They cover the 1444–1786 period, though with a gap between 1684–1741 (except for the years 1698–1701) in connection with the French
100 occupation of Luxembourg (Yante, 1985). It is unfortunate that much of the cold period coinciding with Maunder Minimum of solar activity (*c.* 1645–1715; Eddy, 1976; Steinhilber et al., 2009) is missing in the Moselle Valley wine production data. However, the last three decades of the 16th century were at least as cold as any part of the Maunder Minimum (Pfister and Wanner, 2021; Wanner et al., 2022). Thus, we still capture the full range of temperature variability.

While the tax was originally levied, when introduced in the 13th and 14th centuries, on every ninth basket of grapevines
105 harvested, it soon instead became levied on every ninth barrel of wine produced. However, the number of barrels of wine produced closely followed the number of baskets of grapevines harvest. Thus, the number of barrels of wine produced mimics the quantity of the vine harvest. It is uncertain to what extent changes over time in tax exemptions affected the wine quantity at inter-annual and longer time-scales. Furthermore, the tax appears in fact to, at times, have been both higher and lower than one-ninth of the wine production. The degree of fraud and resistance to taxation might also have varied over time (see also Pfister,
110 1984). Sufficient to say is that the long-term trend in the Grevenmacher and Remich series are not reliable. The source-critical problems associated with tithe records to estimate actual agricultural productivity variations have been extensively treated in the agrarian history literature (e.g., Kain, 1979; Le Roy Ladurie and Goy, 1982; Leijonhufvud, 2001; Skoglund, 2023). This research on tithes as a historical source for agricultural productivity has shown that long-term trends contain larger biases than short-term variability. Grevenmacher shows a long-term rather stable, or slightly decreasing, production trend, whereas
115 Remich shows a long-term production increase (Fig. 1a). Nevertheless, the correlation coefficients between the Grevenmacher and Remich series for linearly detrended and 11-year high-pass filtered data, respectively, are $r = 0.55$ and $r = 0.62$. Both correlations are significant at a $p = 0.01$ level (two-tailed test).

Modern observations show that temperature is the single most important climatic factor for viticulture in the Moselle region (Urhausen et al., 2011). The soil in Grevenmacher is limestone and in Remich is alluvial. Grain is the main crop but
120 terraces of vines are widespread too (Yante, 1985). The annual mean temperature during the twentieth century was around 10°C with around 650 mm annual precipitation and slightly over 1400 hours of sunshine. Years with less than 500 mm pre-




Figure 1. **A** Annual wine production in number of barrels for each series (*in black*) and volcanic sulfate flux derived from Greenland sulfate deposition in kg/km^2 (*in red*). **B** Location of the two sites with wine data from 1444–1786 (with gaps) in the Moselle Valley of Luxembourg.

125 precipitation tend to produce the best wine. Spring frost is a known problem for viticulture in the region (Meier et al., 2018). The Moselle Valley was during the early modern period a rather marginal viticulture area (about 90% of the wine was white) with often relatively low wine quality, with the wine of Remich considered superior to that of Grevenmacher. Consequently, the vine was grown to much larger extent in Remich than in Grevenmacher (Yante, 1985).

2.2 Volcanic forcing data

We employ the volcanic stratospheric sulfur injection data from Toohey and Sigl (2017) (lists provided by M. Sigl, priv. comm., 2022), an updated version of Sigl et al. (2015), that is utilised for the recommended forcing in state-of-the-art PMIP4 ‘past1000’ climate model simulations (Jungclaus et al., 2017). This dataset provides forcing year, not eruption year (which is sometimes unknown), and depending on the seasonality of the eruption the peak forcing year can be the same year as eruption year or the following year. Toohey and Sigl (2017) used composite ice-core data from Greenland and Antarctica, separately, to generate a robust list of dates of volcanic eruptions. We only use the dates for forcing events with aerosol loading in the Northern Hemisphere (NH) (flux is provided as the deposited sulfate found in the ice cores given in kg/km^2). We restrict our list to those eruptions, since it is likely that even strong Southern Hemisphere extra-tropical eruptions do not contribute much to an aerosol loading in the NH.

140 For use in the superposed epoch analysis, we identify key years with NH volcanic forcing of various strengths. In the Toohey and Sigl (2017) NH flux series, for the period we use in the analysis, there are 34 (27) eruptions listed with strengths above $0 \text{ kg}/\text{km}^2$ (the smallest eruption in the series is near $3.2 \text{ kg}/\text{km}^2$); there are 28 (22) eruptions stronger than $5 \text{ kg}/\text{km}^2$; 16 (12) stronger than $10 \text{ kg}/\text{km}^2$; 11 (7) stronger than $15 \text{ kg}/\text{km}^2$, and 9 (6) stronger than $20 \text{ kg}/\text{km}^2$. Here, the numbers in parentheses give the count of eruptions that also cover complete wine data, in the sense that we only include eruptions not less than 5

Table 1. Year and flux (in kg/km²) of the volcanic forcing events 1444–1786 based on Toohey and Sigl (2017). We include the volcanic events occurring 1684–1741 when we have a gap in the wine production series because the events are used for the SEA analysis based on climate extremes.

Year	NH flux [kg/km ²]	Year	NH flux [kg/km ²]
1453	24.80	1646	12.70
1458	35.30	1654	6.20
1463	3.20	1667	18.30
1470	7.30	1673	5.90
1477	27.00	1693	6.96
1480	8.70	1695	24.72
1502	5.30	1707	5.71
1510	12.10	1720	3.77
1528	5.70	1721	4.28
1537	4.50	1729	25.37
1554	5.20	1739	18.09
1567	13.20	1755	6.20
1585	23.30	1762	9.30
1595	11.80	1766	13.20
1600	38.10	1770	3.70
1637	6.80	1783	109.50
1640	41.10	1786	4.40

years from a wine data gap. In our analyses, we use all the eruptions, as recent results suggest that weak eruptions can have significant climate effects (Chim et al., 2023). However, we also tested the effects of using only eruptions larger than the thresholds mentioned above. The eruption years included are listed in Table 1. In addition, tested the sensitivity of the SEA by progressively excluding the strongest volcanic eruptions.

145 2.3 Climate data

Instrumental climate data have only a short period of overlap with the Grevenmacher and Remich wine production series. The closest instrumental temperature station to Luxembourg in the homogenised European Climate Assessment (ECA) temperature dataset (Squintu et al., 2019) is the monthly-averaged data for De Bilt in The Netherlands (van Engelen and Geurts, 1985) at a distance of about 300 km. Data from De Bilt overlaps with the wine production series from 1742 to 1786, covering only a
150 period of 45 years. The short overlap with the De Bilt instrumental record limits the reliability and confidence in results derived from it.

We therefore employ reconstructed series of temperature and precipitation, as summarised in Table 2. We use seasonally resolved gridded reconstructed temperature data from Luterbacher et al. (2004) and Xoplaki et al. (2005) (henceforth only cited as Luterbacher et al. (2004)) and seasonally resolved gridded reconstructed precipitation from Pauling et al. (2006) back to 1500 (Table 2). Both the Luterbacher et al. (2004) and Pauling et al. (2006) reconstructions include instrumental data during the 18th century. Data are extracted from the local grid-cell(s) covering Luxembourg. In addition, we use the seasonal resolved reconstructed temperature data from Dobrovolný et al. (2010), which is available as a Central European average for the same period. For the full period since 1444 we use the June–August temperature reconstruction by Luterbacher et al. (2016), as updated by Ljungqvist et al. (2019), based mainly on tree-ring data but also containing the documentary-based June–August season data from above-mentioned Dobrovolný et al. (2010) reconstruction. Furthermore, we employ for the full period the solely tree-ring-based NTREND gridded field reconstruction (Anchukaitis et al., 2017) of May–August temperature. For comparisons against volcanic-induced cooling in the Alpine region, we use the June–September temperature reconstruction from the Löttschental, Switzerland, which is derived from a strong temperature signal in maximum latewood density tree-ring series (Büntgen et al., 2006). June–August soil moisture (drought) conditions for the entire period are obtained from the tree-ring width based gridded Old World Drought Atlas (Cook et al., 2015) providing self-calibrated Palmer Drought Severity Index (scPDSI) values (van der Schrier et al., 2011).

2.4 Statistical methods

The episodic nature of the volcanic eruptions is an advantage when analysing their effects on variables with unknown, or even dubious, low-frequency variations and trends, such as wine production quantities. We simply compare the strength of the mean superposed standardised signal of the considered time-series (wine production or climate conditions) in the years immediately following the eruptions with the mean superposed standardised signal in the years before (five years). Using superposed epoch analysis (SEA; Chree, 1913, 1914; von Storch and Zwiers, 1999; Rao et al., 2019), we calculate the average of this difference across K eruptions, that are covered by the considered time-series – so the number of eruptions might vary slightly for different time-series – and stronger than the chosen threshold. We standardised the time-series to zero mean and unit variance before conducting the SEA. The significance is estimated by a bootstrap procedure, assuming only that the eruptions are independent. If we have K eruptions, we pick K random years in the whole period under consideration and calculate the superposed epochs from these years. Note, that the years with actual volcanic eruptions should be selected with the same probability as any other year. Otherwise, a bias would be introduced. We do this 3000 times to get a distribution of the SEA under the null-hypothesis that volcanic eruptions have no impact on the considered time-series (e.g., wine production). Comparing the original SEA with this distribution, we can calculate the p -value as the probability of having a lower SEA than the observed. We report significance in the SEA at both the $p = 0.05$ and $p = 0.01$ significance levels. Furthermore, our results are robust to changes in the details of the analysis, such as in the length of the epochs or whether the average is calculated over the whole epoch and not just the half preceding the eruptions. The SEA results are based on time-series standardised to unit variance over the whole period. However, similar results are obtained when the standardisation is performed individually over running-windows (corresponding to variance equalisation).

Table 2. Wine production data, (palaeo)climate and volcanic forcing data series used in this study with information of parameters, period covered, grid-cell (centre coordinates), season, data type, AR1 (autoregressive coefficient at lag 1), and data source.

Data-sets	Period	Grid-cell	Season	Data type(s)	AR1	Source/reference
<i>Wine production data</i>						
Grevenmacher	1444–1786	49.5°N, 6.35°E	Annual	Documentary	0.41	Yante (1985)
Remich	1444–1786	49.5°N, 6.35°E	Annual	Documentary	0.62	Yante (1985)
<i>Temperature data</i>						
Moselle temperature	1500–	49.5°N, 6.5°E	Seasonal	Multi-proxy/instr.	0.04 ^a	Luterbacher et al. (2004); Xoplaki et al. (2005)
Central Europe	1500–	Not gridded	Seasonal	Documentary	0.06 ^a	Dobrovolný et al. (2010)
EuroMed2k (updated) ^b	Full	47.5°N, 7.5°E	Jun–Aug	Tree-ring/doc.	0.67	Ljungqvist et al. (2019)
NTREND	Full	47.5°N, 7.5°E	May–Aug	Tree-ring	0.45	Anchukaitis et al. (2017)
Lötschental, Switzerland	Full	46.42°N, 7.83°E	Jun–Sept	Tree-ring	0.55	Büntgen et al. (2006)
<i>Hydroclimate data</i>						
Moselle precipitation	1500–	49.5°N, 6.5°E	Seasonal	Multi-proxy/instr.	0.07 ^a	Pauling et al. (2006)
scPDSI	Full	49.5°N, 6.5°E	JJA	Tree-ring	0.39	Cook et al. (2015)
<i>Volcanic forcing data</i>						
NH flux	Full	Not gridded	Annual	Ice-core	—	Toohey and Sigl (2017)

^a Mean AR1 of the four different seasonal windows (December–February, March–May, June–August, and September–November).

^b Updated by Ljungqvist et al. (2019) from the first EuroMed2k reconstruction version published by Luterbacher et al. (2016).

To further investigate the relationship between climate and wine production quantities, we calculated the Pearson correlation coefficient (r) between the wine production series and the various (palaeo)climate reconstructions. The significance of correlations were calculated using $p = 0.05$ with a phase-scrambling test, which is more conservative than the common Student’s t -test (for details, see Ljungqvist et al., 2022, 2023). To emphasise inter-annual variability, we employed 11-year high-pass filtering of the data. For studying low-frequency variability, we likewise employed an 11-year low-pass filter. This emphasises variability at decadal scales, while inter-annual fluctuations and noise are suppressed. The choice of 11 years as the filter cut-off length is chosen to match the time-scale of the SEA. Although we have used a simple 11-year smoothing filter, other more complicated filters (e.g., Gaussian filters) provided almost identical results. During an exploratory data analysis phase, we also compared the use of the parametric Pearson correlation coefficient with the non-parametric Kendall’s and Spearman correlation coefficients, finding that they resulted in similar correlation patterns.

3 Results

In this section we study the influence of volcanic eruptions on wine production (subsection 3.1), the influence of volcanic eruptions on climate (subsection 3.2), and the influence of climate extremes on wine production (subsection 3.3). We will mainly




Figure 2. Superposed epoch analysis (SEA) of volcanic forcing, after NH sulfate flux kg/km^2 size, on standardised wine production quantity for Grevenmacher, Remich. The grey bands indicate $p = 0.05$ (light grey) and $p = 0.01$ (dark grey) significance levels, calculated by a bootstrap method (randomly selected key years).

focus on the episodic connection using SEA analysis, while we will additionally use correlation analysis in subsection 3.3. Together, this will provide evidence for a volcanic and climatic influence on the wine production quantity, mainly through spring and summer temperatures.

3.1 Effects of volcanic forcing on Moselle Valley wine production

A main focus of this study is the effects of volcanic-forced abrupt cooling on the Moselle Valley wine production. Superposed epoch analysis (SEA) reveals a strong decrease in wine production the year following volcanic forcing events for both Grevenmacher and Remich. The SEA signal itself is shown in Fig. 2 with the blue curves, while the dark and light shaded bands indicate the $p = 0.01$ and $p = 0.05$ significance levels, respectively. This volcanic-related production decrease pattern is similar regardless of whether all volcanic events are included or only those exceeding certain thresholds for NH forcing flux (as demonstrated in the different panels in Fig. 2). In addition, progressively excluding the strongest volcanic eruptions one by one had only a minor impact on the results (not shown). Conclusions were unchanged.

In general, we find a strong, significant, and consistent decrease in wine production in the years following volcanic forcing events for both the Grevenmacher and Remich series. For all thresholds the signal (blue curve in Fig. 2) is about half a standard deviation and the p -value is below 0.05 and often below 0.01 (shaded regions in Fig. 2). Slightly smaller signals are found for Remich compared to Grevenmacher. Including all eruptions, we find the largest signal in the first year after the eruptions. The (taxed) wine production was even zero, or close to it, in the first year following certain major volcanic events: 1481, 1601,




Figure 3. For each climate series in Table 2, a SEA based on the years of the volcanic forcing event has been performed. The lowest p -value (two-tailed test) in the year of the volcanic forcing event and the following 5 years are shown. The x-axis is logarithmic, and p -values of 0.01 and 0.05 are shown with the red vertical lines. The p -values are calculated using the bootstrap method. The significant signals in the temperature/precipitation series all correspond to colder/wetter conditions. To the right is shown the lag where the lowest p -value is found. Volcanic forcing events of all strengths are included, the exact number is dependent on data availability and is also shown.

215 1674, 1767, and 1784. However, when increasing the threshold and only including stronger (and fewer) eruptions, we see a tendency for the strongest wine production decrease to be delayed. For the largest eruptions, the strongest signals appear 3–4 years after the eruptions.

The bands indicating the significance levels expand with increasing threshold due to the fewer eruptions included (Fig. 2). When only including larger volcanic events, exceeding an estimated NH forcing flux of >10 , a wine production decrease that is significant at the $p = 0.01$ level is only found the third year after the volcanic event – and only for Grevenmacher. Weaker, 220 less significant, decreases are also detected the first and second years after the volcanic event, with the Remich series showing a broad decrease for years 1–3 after such strong eruptions.

3.2 Effects of volcanic forcing on Moselle Valley pre-industrial climate

We have performed SEA from volcanic forcing events on temperature and precipitation series (local grid-cells when applicable) from the Moselle Valley (see Table 2). Figure 3 summarises the strengths of the connection by reporting the smallest p -values (two-tailed test) taken over years 0–5, i.e., from the year of the eruptions and the following 5 years. These p -values basically reflect the strength of the SEA signals.

We find that significant connections for the temperature series mainly in spring and summer. For example, p -values below 0.02 are found for the temperature series by Dobrovolný et al. (2010) and Luterbacher et al. (2004) only for the March–May and June–August seasons. For temperature in spring and summer, the lowest p -values are found for lag 0 or 1 year (numbers shown to the right in Fig. 3). The significant signals in the temperature series all correspond to coolings. Similar results are obtained using only eruptions larger than 5 kg/km^2 , while the situation for higher thresholds (and correspondingly fewer eruptions) is more muddled. For precipitation, significant signals corresponding to wetter conditions are found in annual and September–November reconstructions.

We emphasise the weak and insignificant volcanic signatures in the tree-ring based temperature reconstructions (EuroMed2k, NTREND, and Löttschental) for the Central Europe region. This less distinct volcanic signature stands in sharp contrast to the strong and clear wine production quantity reduction after volcanic forcing events, that reaches significance well below the $p = 0.01$ level for several years even when including minor forcing events (Fig. 2).

3.3 Effects of climate on Moselle Valley wine production

We have identified the 20 most extreme years (for each of the four seasons) for a selection of the climate series in Table 2. Thus, for temperature series we find the 20 most warm and cold years and for precipitation the 20 most wet and dry years. These extremes are identified from the raw series without detrending. We then conducted SEA to identify the responses in the two wine production series to these four kinds of extremes for each season. The results are shown in Table 3 which reports the p -values and the signs of the significant responses.

We find that the spring (MAM) stands out from the other seasons with significant responses for all four kinds of extremes. We also note that consistent directions of the responses are found; increasing wine production for warm/dry conditions and decreasing production for cold/wet conditions. The climate series from Luterbacher et al. (2004) and Pauling et al. (2006) show consistent results. Furthermore, we find almost identical results for the Remich and for Grevenmacher wine series. For the other seasons, fewer significant and less consistent responses are found, indicating that the significance in these seasons might partly be due to chance.

Through correlation analyses we explored the long-term relationships between wine production and seasonal and annual temperature, precipitation and soil moisture. We started the analyses with De Bilt instrumental series. The correlation between the De Bilt instrumental series and the Grevenmacher series, over their 45-year period of overlap 1742–1786, is $r = 0.46$ using 11-year high-pass filtered data for JJA temperature for both. For Remich the result is $r = 0.29$ (just below significance). Furthermore, annual mean temperature for De Bilt shows significant positive correlations with Grevenmacher and Remich of r

Table 3. The p -values for the SEA performed using as key years the 20 warmest/coldest/wettest/driest years, respectively, in the seasonal series of the Luterbacher et al. (2004) temperature reconstruction and the Pauling et al. (2006) precipitation reconstruction on the Remich and Grevenmacher wine production series. The p -values (two-tailed test) were calculated using repeated scrambling of years (bootstrap), see text for details, and p -values below 0.05 are typeset in bold. The lowest p -value (two-tailed test) in the year of the eruptions and the following 5 years are shown in the table. (+) and (–) designate the direction of the effect on the wine series, where the strongest excursion on the SEA curve inside the 0 to +5 years is used (which is often year 0 or +1).

Type	Climate series	Season	Wine series	p -value
Warmest	Luterbacher et al. (2004) temp.	DJF	Remich	0.079
Coldest	Luterbacher et al. (2004) temp.	DJF	Remich	0.031 (–)
Wettest	Pauling et al. (2006) precip.	DJF	Remich	0.042 (–)
Driest	Pauling et al. (2006) precip.	DJF	Remich	0.012 (–)
Warmest	Luterbacher et al. (2004) temp.	DJF	Grevenmacher	0.483
Coldest	Luterbacher et al. (2004) temp.	DJF	Grevenmacher	0.054
Wettest	Pauling et al. (2006) precip.	DJF	Grevenmacher	0.007 (–)
Driest	Pauling et al. (2006) precip.	DJF	Grevenmacher	0.164
Warmest	Luterbacher et al. (2004) temp.	MAM	Remich	0.005 (+)
Coldest	Luterbacher et al. (2004) temp.	MAM	Remich	0.008 (–)
Wettest	Pauling et al. (2006) precip.	MAM	Remich	0.019 (–)
Driest	Pauling et al. (2006) precip.	MAM	Remich	0.010 (+)
Warmest	Luterbacher et al. (2004) temp.	MAM	Grevenmacher	0.003 (+)
Coldest	Luterbacher et al. (2004) temp.	MAM	Grevenmacher	0.022 (–)
Wettest	Pauling et al. (2006) precip.	MAM	Grevenmacher	0.002 (–)
Driest	Pauling et al. (2006) precip.	MAM	Grevenmacher	0.006 (+)
Warmest	Luterbacher et al. (2004) temp.	JJA	Remich	0.079
Coldest	Luterbacher et al. (2004) temp.	JJA	Remich	0.016 (–)
Wettest	Pauling et al. (2006) precip.	JJA	Remich	0.323
Driest	Pauling et al. (2006) precip.	JJA	Remich	0.006 (+)
Warmest	Luterbacher et al. (2004) temp.	JJA	Grevenmacher	0.083
Coldest	Luterbacher et al. (2004) temp.	JJA	Grevenmacher	0.035 (–)
Wettest	Pauling et al. (2006) precip.	JJA	Grevenmacher	0.090
Driest	Pauling et al. (2006) precip.	JJA	Grevenmacher	0.129
Warmest	Luterbacher et al. (2004) temp.	SON	Remich	0.020 (+)
Coldest	Luterbacher et al. (2004) temp.	SON	Remich	0.058
Wettest	Pauling et al. (2006) precip.	SON	Remich	0.051
Driest	Pauling et al. (2006) precip.	SON	Remich	0.049 (–)
Warmest	Luterbacher et al. (2004) temp.	SON	Grevenmacher	0.447 (+)
Coldest	Luterbacher et al. (2004) temp.	SON	Grevenmacher	0.181
Wettest	Pauling et al. (2006) precip.	SON	Grevenmacher	0.131
Driest	Pauling et al. (2006) precip.	SON	Grevenmacher	0.060




Figure 4. Correlations between the (palaeo)climate series listed in Table 2 (shown vertically) and wine production quantity data from Grevenmacher and Remich (shown horizontally). The three panels are for different temporal filtering. From left to right: 11-year high-pass filtered data, linearly detrended data, and 11-year low-pass filtered data (after linearly detrending). Correlations significant at $p = 0.05$ with the phase-scrambling test are marked with a plus sign (+).

= 0.46 and $r = 0.42$, respectively, using 11-year high-pass filtered data. No other seasons, neither for linearly detrended data nor high-pass filtered data, reveal any significant correlation between wine production and the De Bilt instrumental temperature series.

Using reconstructed temperature data covering the entire, or the majority, of the length of the Grevenmacher and Remich wine production series shows similar, but more significant, correlation patterns as found using the short De Bilt instrumental series. Mostly significant correlations are found between temperature and wine production quantities between both 11-year high-pass filtered and linearly detrended data (Fig. 4). These correlations are, in general, slightly stronger for Grevenmacher than for Remich. The most significant association is the positive relationship between spring, summer and autumn temperature, as well as for the annual mean, and wine production (i.e., warm = high wine production and *vice versa*). Correlations are weak for winter temperatures being entirely outside the growing season (Fig. 4). Significant positive correlations are also

found between (palaeo)climate series and the Grevenmacher series using 11-year low-pass filtered data. While some of the correlations are slightly stronger than those for high-pass filtered or linearly detrended data, the fewer degrees of freedom render a lower number of them significant. Notably, for Remich not a single correlation using 11-year low-pass filtered data is significant, although most of the correlations are positive and of comparable strength as those for high-pass filtered and linearly
270 detrended data.

Negative relationships, in general less significant than those with temperature, are found in the hydroclimate between precipitation and soil moisture and wine production (i.e., wetter = lower wine production and *vice versa*) for both 11-year high-pass filtered data and linearly detrended data. Significant relationships with hydroclimate are mostly restricted to the summer season and to the annual mean. We note that while the correlation-based results provide the strongest signal for summer, the SEA-
275 based results using extreme years instead show the strongest signal in spring. Nevertheless, both analyses reinforce the finding that warm and dry years and growing seasons benefited the wine production, while cold and wet years had an adverse effect. For hydroclimate, no significant correlations are found with either the Grevenmacher or Remich series using 11-year low-pass filtered (palaeo)data.

4 Discussion

280 4.1 The volcanic and climatic signals embedded in the wine production series

Our results are robust to choices regarding the exclusion of weaker volcanic forcing events as well as to the exclusion of individual particular strong volcanic forcing events. Although wine tithes data is zero, or close to it, in the first year following certain major volcanic events (1481, 1601, 1674, 1767 and 1784), this is not necessarily meaning that the vine harvest was zero or close to zero. From grain tithes, for example in the early modern Swedish Realm (Leijonhufvud, 2001; Huhtamaa
285 et al., 2022), we know that sometimes no tithes were collected during years with extremely low harvests. Thus, zero wine tithes indicates very low vine harvests rather than necessarily no harvest at all. Furthermore, we cannot rule out the possibility that the respective error term is skewed towards a higher resistance to taxation in years of bad harvests. Such non-linearities are very difficult to assess and can similarly affect other types of records. Nevertheless, the long-term trend of the wine tithes data is more uncertain than the short-term (inter-annual to inter-decadal) variability. Since the SEA method focuses on a 5-year period
290 before and after each event, it is not sensitive to potential long-term trends in the data. The compositing in SEA constitutes an averaging process that serves as a filter enhancing the high-frequency response signal of interest while minimising noise and also accounts for long-term drifts (see, e.g., Rao et al., 2019). We have also ensured that the response statistics to volcanic forcing are not driven by just a few major eruptions by both investigating the effects of including all volcanic forcing events and only those exceeding different threshold values. The results are robust also to using forcings less than a threshold. We have
295 also tested the effects of excluding only the 1783 Laki eruption (not shown) and found that this volcanic forcing event is not dominating the results, although the forcing of Laki in the Northern Hemisphere is multiple times stronger than that of any other eruption.

Regarding the climate–wine production relationships, we can emphasise that we have tested both linear and non-linear correlation methods (that is Pearson correlation coefficient *versus* rank correlation methods such as Kendall and Spearman) although we only report the results of the first as they all give similar results. The similar results imply that the non-linearity that is indeed present in the wine data (which cannot go below zero) and various non-linearities present in palaeoclimate reconstructions have not prevented the linear Pearson correlation from rendering an accurate picture of how the series are correlated. The varying correlation strength between wine production and the different temperature reconstructions, even for the same season, is not surprising considering that the reconstructions have employed diverse input proxy data and different methods to combine the data and ‘calibrate’ the reconstructions against instrumental temperature measurements (see, for example, Christiansen and Ljungqvist, 2017; Anchukaitis and Smerdon, 2022).

Reduced wine production for up to several years following larger volcanic forcing events can, in part, be attributed to the fact that strong eruptions cause aerosols to linger for a longer period in the atmosphere, and induces feedback mechanisms in the climate system that prolongs cooling (e.g., sea-ice feedbacks, Miller et al., 2012; Toohey et al., 2016). In part, it can also be attributed to a biological memory effect, and cold and frost damage, of the vine stocks during years with very poor growth conditions (Meier et al., 2018; Pfister and Wanner, 2021). Note that while we do observe an immediate signal following larger volcanic events, the SEA results suggest that the maximum response is delayed. Biological memory effects are very well-documented for tree growth (Esper et al., 2015; Hartl-Meier et al., 2017). Especially, tree-ring width depends to some degree on growth conditions in the previous year and resulting climate reconstructions often show more auto-correlation than the instrumental target (Ljungqvist et al., 2020). As expected, auto-correlation values (AR1) reported for the time-series used in this study are higher, if biological archives are involved compared to documentary-based records (see Table 2). But while a limited immediate response to abrupt volcanic cooling has been reported for many tree-ring width records (Esper et al., 2015), as opposed to maximum latewood density (MXD) records (Hartl-Meier et al., 2017), this seems not to be the case for the wine production series. Thus, in this respect wine production shows a similar behaviour to MXD, potentially indicating that other physiological processes control fruit production than stem increment.

4.2 Further comparison between the volcanic effect on wine production and tree-ring growth

The wine production decline in Grevenmacher and Remich following volcanic forcing events contrasts to the weak volcanic response for Central Europe in the state-of-the-art gridded NTREND tree-ring-based temperature reconstruction (Anchukaitis et al., 2017), the mainly tree-ring-based updated EuroMed2k temperature reconstruction (Ljungqvist et al., 2019) as well as the MXD-based Löttschental temperature reconstruction (Büntgen et al., 2006). The NTREND reconstruction is for Europe primarily composed of MXD data from tree-rings, a temperature proxy commonly assumed to be little affected by biological memory (Anchukaitis et al., 2012; Esper et al., 2015). Despite remaining discussions regarding the precise quantification of volcanic signals in MXD-based reconstructions (Tingley et al., 2014; Edwards et al., 2022), there has been repeated evidence for significant cooling in MXD-based reconstructions following single events (Guillet et al., 2017; Hartl-Meier et al., 2017) and groups of eruptions at local (Rydval et al., 2017), regional (Esper et al., 2013b) and hemispheric (Wilson et al., 2016; Schneider et al., 2017) scales.

In contrast to a moderate volcanic cooling impact over Central Europe, NTREND and EuroMed2k reveal strong and widespread cooling in response to tropical eruptions over Scandinavia with a distinct transition between these two regions over the southern Baltic Sea region at around 55°N. However, NTREND and EuroMed2k do not contain tree-ring data in Europe between 50°N and 60°N making it difficult to determine the exact location of this transition. The gridded seasonal temperature reconstruction for Europe by Luterbacher et al. (2004), using predominantly documentary data, suggests significant summer cooling that extends as far south as the northern fringe of the Alpine arc (Fischer et al., 2007), which is better in agreement with the volcanic impact found here in wine production series from the Moselle Valley. The lack of climate-sensitive tree-ring records from Central Europe (north of the Alpine Arc) is also expressed in reduced values for explained variance at the Moselle grid-cell compared to the explained variance for, e.g., northern Fennoscandia.

Comparison between wine production declines and tree-ring growth declines is complicated by the limited number of major volcanic forcing events during the comparatively short period covered by the two wine production series. Numerous studies of temperature-sensitive tree-ring data, from northern Europe and elsewhere, have shown a maximum growth decline (growing season cooling) during the summer in the year following the volcanic forcing event (Esper et al., 2017). Such a distinct one year post-eruption cooling is most evident in MXD data, compared to TRW data, owing to the typically stronger correlation with temperature and the larger biological memory in TRW (Hartl-Meier et al., 2017). Thus, the post-volcanic decrease detected in this study in wine production quantities is very similar to the post-volcanic growth decline in MXD. Nevertheless, the most noteworthy feature in the wine production data is the strong and consistent response one year following the volcanic forcing event even for a low NH flux threshold (> 0 and > 5) that is absent in tree-ring based temperature reconstructions for Central Europe. At the mid-latitudes, where the wine production data diverges from the tree-ring based temperature reconstructions, no reconstructions show a significant summer temperature reduction in response to a low NH forcing flux threshold.

The Moselle Valley region is situated at the northern limit of viable viticulture areas in the same way as northern Scandinavia is situated close to the Arctic tree-line. Hence, both the Moselle Valley grapevines and the Scots pine trees in northern Scandinavia are thus very sensitive to temperature drops and presumably also to reduced sunlight after volcanic forcing events. The high climate sensitivity in both data types is clearly related to the marginally of the respective locations for grapevines and conifer trees. The higher climate sensitivity of the grapevines of Grevenmacher compared to Remich may also, rather than merely the small difference in latitude, be due to different soil conditions. We also suggest the explanation for the stronger, and more persistent, climate response in the wine production data than in the tree-ring data is due to physiological differences. In addition to the consistent signal one year following the volcanic forcing event, there is in the wine production data a 'late response' after big flux events (thresholds > 15 and > 20). This late signal is also present in the entirely tree-ring-based NTREND (Anchukaitis et al., 2017) and (weakly) in the Löttschental (Büntgen et al., 2006) temperature reconstructions (not shown). It is more difficult to come to a conclusion about this part of the signal because there are only few events falling into this category. While it would be interesting to study the climate (and volcanic) response on tree growth and wine production from the same locations, it would be challenging because temperature-sensitive and/or precipitation-sensitive tree-ring series rarely are available from vine growing regions with the exception of regions close to the Alps. Nevertheless, such studies could

provide insights into the decorrelation length of climate *versus* differences in the climate sensitivity of the vine growth and tree-ring growth.

5 Conclusions and outlook

We have systematically investigated the impacts of volcanic forcing events and climate variability on two of the longest wine
370 production quantity records (spanning 1444–1786 with gaps) in Europe deriving from the Moselle Valley close to the northern
limit of viticulture. We primarily used SEA analysis, which is particularly appropriate for episodic events. The statistical
significance of the SEA was estimated with a bootstrap method. We also assessed the long-term relationship between climate
and wine production using correlation analyses and calculating the significance with a conservative phase-scrambling test.

A strong negative impact of volcanic eruptions on wine production quantity was found in the year, and years, following a
375 volcanic forcing event (subsection 3.1). This finding is robust under removal of weaker volcanic eruptions. Both SEA based
on climate extremes and correlation analysis (11-year high- and low-pass filtered and linearly detrended) consistently show
that cold and wet conditions are detrimental for wine production (subsection 3.3). Furthermore, following volcanic eruptions
we see anomalously cold and wet conditions in the Moselle Valley region (subsection 3.2) which in the light of these results
from subsection 3.3 indicates negative impacts on the wine production. This is exactly what we find in subsection 3.1. For both
380 the volcanic impact on climate (subsection 3.2) and the impact of climate extremes on wine production (subsection 3.3), the
largest effect is found in spring and summer. Thus, taken together, these findings present a consistent picture of the effect of
volcanic eruptions on wine production and how it is mediated through climate.

We furthermore note that the detected volcanic signature in the Moselle Valley wine production is distinct and statistically
significant (subsection 3.1), while that of tree-ring based temperature reconstructions for Central Europe is neither (subsec-
385 tion 3.2). Based on our findings we conclude that long series of annual wine production quantity contains very valuable
biological and climatic information and we, thus, encourage further archival research to compile and publish additional wine
production quantity series from across the viticulture regions of Europe.

Code availability. We have used **IDL** and **R** (R Core Team, 2022) version 3.6.3 to program the analysis codes used in this work. In **R** we
used the package ‘corrplot’ (Wei and Simko, 2021) to generate the correlation matrices. Data were read from mixed files using ‘base’ libraries
390 for text files or ‘openxlsx’ (Schauberger and Walker, 2021) for Excel spreadsheet files. The superposed epoch analysis was performed with
R code, available upon reasonable request.

Data availability. The wine production data can be obtained from Appendix 1 in the article by Yante (1985), pp. 301–307. All used
(palaeo)climate data are digitally available from the National Oceanic and NOAA Paleoclimatology/World Data Center for Paleoclima-
tology:

395 <https://www.ncdc.noaa.gov/products/paleoclimatology>. The monthly De Bilt temperature series (van Engelen and Geurts, 1985) are available

from the KNMI Climate Explorer (Trouet and Van Oldenborgh, 2013):

<https://climexp.knmi.nl/getindices.cgi?WMO=KNMIData/labrijn&STATION=Tdebilt&TYPE=i>

Author contributions. F.C.L. designed the study together with B.C. and P.T. F.C.L., B.C. and P.T performed data analyses. B.C. and P.T conducted the SEA and the phase-scrambling significance tests. L.S. especially contributed with comparisons between the volcanic signature
400 in wine production data and tree-ring data. All authors interpreted the results and wrote the article together.

Competing interests. The authors have declared no conflicts of interest for this article.

Acknowledgements. F.C.L. acknowledges Visiting Researcher stays at the Institute of History, University of Bern, and at the Freiburg Institute for Advanced Studies (FRIAS), that allowed him time to work with this article. We express our appreciation to Asst. Prof. Michael Sigl, University of Bern, for lending us assistance regarding volcanic forcing datasets, and to Prof. em. Christian Pfister, University of Bern, for
405 stimulating and insightful discussions about wine production data and grapevine phenology. The authors thank the four reviewers whose useful comments helped improve this article.

Financial support. F.C.L. was supported by the Marianne and Marcus Wallenberg Foundation (grant no. MMW 2022-0114) and the Swedish Research Council (Vetenskapsrådet, grant no. 2018-01272 and grant no. 2023-00605). He conducted the work with this article as a Pro Futura Scientia XIII Fellow funded by the Swedish Collegium for Advanced Study through Riksbankens Jubileumsfond. B.C. and P.T. were
410 supported by the National Centre for Climate Research at the Danish Meteorological Institute. Open access publication funding for this article was provided by Stockholm University.

References

- Allen, R. C. and Unger, R. W.: The Allen-Unger Global Commodity Prices Database, *Research Data Journal for the Humanities and Social Sciences*, 4, 81–90, 2019.
- 415 Anchukaitis, K. J. and Smerdon, J. E.: Progress and uncertainties in global and hemispheric temperature reconstructions of the Common Era, *Quaternary Science Reviews*, 286, 107 537, 2022.
- Anchukaitis, K. J., Breitenmoser, P., Briffa, K. R., Buchwal, A., Büntgen, U., Cook, E. R., D'Arrigo, R. D., Esper, J., Evans, M. N., Frank, D., Grudd, H., Gunnarson, B., Hughes, M., Kirilyanov, A., Körner, C., Krusic, P., Luckman, B., Melvin, T. M. and Salzer, M., Shashkin, A., Timmermann, C., Vaganov, E., and RJS, W.: Tree rings and volcanic cooling, *Nature Geoscience*, 5, 836–837, 2012.
- 420 Anchukaitis, K. J., Wilson, R., Briffa, K. R., Büntgen, U., Cook, E. R., D'Arrigo, R., Davi, N., Esper, J., Frank, D., Gunnarson, B. E., Hegerl, G., Helama, S., Klesse, S., Krusic, P., Linderholm, H. W., Myglan, V., Osborn, T. J., Zhang, P., Rydval, M., Schneider, Lea, Z., and Eduardo: Last millennium Northern Hemisphere summer temperatures from tree rings: Part II, spatially resolved reconstructions, *Quaternary Science Reviews*, 163, 1–22, 2017.
- Andrade, C., Fonseca, A., Santos, J. A., Bois, B., and Jones, G. V.: Historic changes and future projections in Köppen–Geiger climate 425 classifications in major wine regions worldwide, *Climate*, 12, 94, 2024.
- Bock, A., Sparks, T. H., Estrella, N., and Menzel, A.: Climate-induced changes in grapevine yield and must sugar content in Franconia (Germany) between 1805 and 2010, *PLoS One*, 8, e69015, 2013.
- Brönnimann, S. and Krämer, D.: Tabora and the “Year Without a Summer” of 1816: A Perspective on Earth and Human Systems Science, Oeschger Centre for Climate Change Research, University of Bern, Bern, 2016.
- 430 Büntgen, U., Frank, D. C., Nievergelt, D., and Esper, J.: Summer temperature variations in the European Alps, AD 755–2004, *Journal of Climate*, 19, 5606–5623, 2006.
- Büntgen, U., Allen, K., Anchukaitis, K. J., Arseneault, D., Boucher, É., Bräuning, A., Chatterjee, S., Cherubini, P., Churakova, O. V., Corona, C., et al.: The influence of decision-making in tree ring-based climate reconstructions, *Nature Communications*, 12, 3411, 2021.
- Burke, A., Innes, H. M., Crick, L., Anchukaitis, K. J., Byrne, M. P., Hutchison, W., McConnell, J. R., Moore, K. A., Rae, J. W., Sigl, M., 435 and Wilson, R.: High sensitivity of summer temperatures to stratospheric sulfur loading from volcanoes in the Northern Hemisphere, *Proceedings of the National Academy of Sciences*, 120, e2221810 120, 2023.
- Chim, M. M., Aubry, T. J., Abraham, N. L., Marshall, L., Mulcahy, J., Walton, J., and Schmidt, A.: Climate projections very likely underestimate future volcanic forcing and its climatic effects, *Geophysical Research Letters*, 50, e2023GL103743, <https://doi.org/10.1029/2023GL103743>, 2023.
- 440 Chree, C.: Some Phenomena of Sunspots and of Terrestrial Magnetism at Kew Observatory, *Philos. Trans. Roy. Soc. London*, 212, 75–116, 1913.
- Chree, C.: Some phenomena of sunspots and of terrestrial magnetism. Part II, *Philos. Trans. Roy. Soc. London*, 213, 245–277, 1914.
- Christiansen, B. and Ljungqvist, F. C.: Challenges and perspectives for large-scale temperature reconstructions of the past two millennia, *Rev Geophys*, 55, 40–96, 2017.
- 445 Chuine, I., Yiou, P., Viovy, N., Seguin, B., Daux, V., and Ladurie Le Roy, E.: Grape ripening as a past climate indicator, *Nature*, 432, 289–290, 2004.
- Cook, B. I. and Wolkovich, E. M.: Climate change decouples drought from early wine grape harvests in France, *Nature Climate Change*, 6, 715–719, 2016.

- Cook, E. R., Seager, R., Kushnir, Y., Briffa, K. R., Büntgen, U., Frank, D., Krusic, P. J., Tegel, W., van der Schrier, G., Andreu-Hayles, L., Baillie, M., Baittinger, C., Bleicher, N., Bonde, N., Brown, D., Carrer, M., Cooper, R., Čufar, K., Dittmar, C., Esper, J., Griggs, C., Gunnarson, B., Günther, B., Gutierrez, E., Haneca, K., Helama, S., Herzig, F., Heussner, K.-U., Hofmann, J., Janda, P., Kontic, R., Köse, N., Kyncl, T., Levanič, T., Linderholm, H., Manning, S., Melvin, T. M., Miles, D., Neuwirth, B., Nicolussi, K., Nola, P., Panayotov, M., Popa, I., Rothe, A., Seftigen, K., Seim, A., Svarva, H., Svoboda, M., Thun, T., Timonen, M., Touchan, R., Trotsiuk, V., Trouet, V., Walder, F., Wazny, T., Wilson, R., and Zang, C.: Old World megadroughts and pluvials during the Common Era, *Sci Adv*, 1, e1500561, 2015.
- 450 D'Arrigo, R., Wilson, R., and Anchukaitis, K. J.: Volcanic cooling signal in tree ring temperature records for the past millennium, *Journal of Geophysical Research: Atmospheres*, 118, 9000–9010, 2013.
- D'Arrigo, R., Klinger, P., Newfield, T., Rydval, M., and Wilson, R.: Complexity in crisis: The volcanic cold pulse of the 1690s and the consequences of Scotland's failure to cope, *J. Volcanol. Geotherm. Res.*, 389, 106746, 2020.
- de Cortazar Atauri, I. G., Duchêne, E., Destrac, A., Barbeau, G., De Rességuier, L., Lacombe, T., Parker, A. K., Saurin, N., and Van Leeuwen, C.: Grapevine phenology in France: from past observations to future evolutions in the context of climate change, *OENO One*, 51, 115–126, 2017.
- 460 Dobrovolný, P., Moberg, A., Brázdil, R., Pfister, C., Glaser, R., Wilson, R., van Engelen, A., Limanówka, D., Kiss, A., Halfčková, M., Macková, J., Riemann, D., Luterbacher, J., and Böhm, R.: Monthly, seasonal and annual temperature reconstructions for Central Europe derived from documentary evidence and instrumental records since AD 1500, *Clim Change*, 101, 69–107, 2010.
- 465 Drappier, J., Thibon, C., Rabot, A., and Geny-Denis, L.: Relationship between wine composition and temperature: Impact on Bordeaux wine typicity in the context of global warming, *Critical Reviews in Food Science and Nutrition*, 59, 14–30, 2019.
- Droulia, F. and Charalampopoulos, I.: A review on the observed climate change in Europe and its impacts on viticulture, *Atmosphere*, 13, 837, 2022.
- Eddy, J. A.: The Maunder Minimum, *Science*, 192, 1189–1202, 1976.
- 470 Edwards, J., Anchukaitis, K. J., Gunnarson, B. E., Pearson, C., Seftigen, K., von Arx, G., and Linderholm, H. W.: The origin of tree-ring reconstructed summer cooling in northern Europe during the 18th century eruption of Laki, *Paleoceanography and Paleoclimatology*, 37, e2021PA004386, 2022.
- Esper, J., Büntgen, U., Luterbacher, J., and Krusic, P. J.: Testing the hypothesis of post-volcanic missing rings in temperature sensitive dendrochronological data, *Dendrochronologia*, 31, 216–222, 2013a.
- 475 Esper, J., Schneider, L., Krusic, P. J., Luterbacher, J., Büntgen, U., Timonen, M., Sirocko, F., and Zorita, E.: European summer temperature response to annually dated volcanic eruptions over the past nine centuries, *Bulletin of Volcanology*, 75, 1–14, 2013b.
- Esper, J., Schneider, L., Smerdon, J. E., Schöne, B. R., and Büntgen, U.: Signals and memory in tree-ring width and density data, *Dendrochronologia*, 35, 62–70, 2015.
- Esper, J., Krusic, P. J., Ljungqvist, F. C., Luterbacher, J., Carrer, M., Cook, E., Davi, N. K., Hartl-Meier, C., Kirilyanov, A., Konter, O., Myglan, V., Timonen, M., Treydte, K., Trouet, V., Villalba, R., Yang, B., and Büntgen, U.: Ranking of tree-ring based temperature reconstructions of the past millennium, *Quat. Sci. Rev.*, 145, 134–151, 2016.
- 480 Esper, J., Büntgen, U., Hartl-Meier, C., Oppenheimer, C., and Schneider, L.: Northern Hemisphere temperature anomalies during the 1450s period of ambiguous volcanic forcing, *Bulletin of Volcanology*, 79, 41, 2017.
- Fischer, E. M., Luterbacher, J., Zorita, E., Tett, S., Casty, C., and Wanner, H.: European climate response to tropical volcanic eruptions over the last half millennium, *Geophysical Research Letters*, 34, L05707, 2007.
- 485

- García de Cortázar-Atauri, I., Daux, V., Garnier, E., Yiou, P., Viovy, N., Seguin, B., Boursiquot, J., Parker, A., Van Leeuwen, C., and Chuine, I.: An assessment of error sources when using grape harvest date for past climate reconstruction, *The Holocene*, 20, 599–608, 2010.
- Garnier, E., Daux, V., Yiou, P., and García de Cortázar-Atauri, I.: Grapevine harvest dates in Besançon (France) between 1525 and 1847: Social outcomes or climatic evidence?, *Climatic Change*, 104, 703–727, 2011.
- 490 Guereau, A.: Climat et vendanges (XIV^e–XIX^e siècles): révisions et compléments, *Histoire & Mesure*, 10, 89–147, 1995.
- Guillet, S., Corona, C., Stoffel, M., Khodri, M., Lavigne, F., Ortega, P., Eckert, N., Sielenou, P. D., Daux, V., Churakova, O. V., et al.: Climate response to the Samalas volcanic eruption in 1257 revealed by proxy records, *Nature Geoscience*, 10, 123–128, 2017.
- Guillet, S., Corona, C., Ludlow, F., Oppenheimer, C., and Stoffel, M.: Climatic and societal impacts of a “forgotten” cluster of volcanic eruptions in 1108–1110 CE, *Scientific Reports*, 10, 1–10, 2020.
- 495 Guiot, J., Bernigaud, N., Bondeau, A., Bouby, L., and Cramer, W.: Viticulture extension in response to global climate change drivers – lessons from the past and future projections, *Climate of the Past*, 19, 1219–1244, 2023.
- Hartl-Meier, C., Büntgen, U., Smerdon, J. E., Zorita, E., Krusic, P. J., Ljungqvist, F. C., Schneider, L., and Esper, J.: Temperature covariance in tree ring reconstructions and model simulations over the past millennium, *Geophysical Research Letters*, 44, 9458–9469, 2017.
- Huhtamaa, H., Stoffel, M., and Corona, C.: Recession or resilience? Long-range socioeconomic consequences of the 17th century volcanic eruptions in northern Fennoscandia, *Climate of the Past*, 18, 2077–2092, 2022.
- 500 Jones, G. V., White, M. A., Cooper, O. R., and Storchmann, K.: Climate change and global wine quality, *Climatic Change*, 73, 319–343, 2005.
- Jungclaus, J., Bard, E., Baroni, M., Braconnot, P., Cao, J., Chini, L., Egorova, T., Evans, M., González-Rouco, J. F., Goosse, H., Hurr, G. C., Joos, F., Kaplan, J. O., Khodri, M., Klein Goldewijk, K., Krivova, N., LeGrande, A. N., Lorenz, S. J., Luterbacher, J., Man, W., Maycock, A. C., Meinshausen, M., Moberg, A., Muscheler, R., Nehrbass-Ahles, C., Otto-Bliesner, B. I., Phipps, S. J., Pongratz, J., Rozanov, E., Schmidt, G. A., Schmidt, H., Schmutz, W., Schurer, A., Shapiro, A. I., Sigl, M., Smerdon, J. E., Solanki, S. K., Timmreck, C., Toohey, M., Usoskin, I. G., Wagner, S., Wu, C.-J., Yeo, K. L., Zanchettin, D., Zhang, Q., and Zorita, E.: The PMIP4 contribution to CMIP6–Part 3: The last millennium, scientific objective, and experimental design for the PMIP4 past1000 simulations, *Geosci Model Dev*, 10, 4005–4033, 2017.
- 505 Kain, R.: Tithe as an index of pre-industrial agricultural production, *Agric. Hist. Rev.*, 27, 73–81, 1979.
- Keenan, D.: Grape harvest dates are poor indicators of summer warmth, *Theoretical and Applied Climatology*, 87, 255–256, 2007.
- Kiss, A., Wilson, R., and Bariska, I.: An experimental 392-year documentary-based multi-proxy (vine and grain) reconstruction of May–July temperatures for Kőszeg, West-Hungary, *International Journal of Biometeorology*, 55, 595–611, 2011.
- Körner, C.: The cold range limit of trees, *Trends in Ecology & Evolution*, 36, 979–989, 2021.
- 515 Labbé, T., Pfister, C., Brönnimann, S., Rousseau, D., Franke, J., and Bois, B.: The longest homogeneous series of grape harvest dates, Beaune 1354–2018, and its significance for the understanding of past and present climate, *Climate of the Past*, 15, 1485–1501, 2019.
- Lachiver, M.: Vins, vignes et vigneron. Histoire du vignoble français, Paris, Fayard, 1988.
- Ladurie Le Roy, E. and Baulant, M.: Grape harvests from the fifteenth through the nineteenth centuries, *The Journal of Interdisciplinary History*, 10, 839–849, 1980.
- 520 Landsteiner, E.: The crisis of wine production in late sixteenth-century central Europe: Climatic causes and economic consequences, *Clim Change*, 43, 323–334, 1999.
- Le Roy Ladurie, E. and Goy, J.: Tithe and Agrarian History from the Fourteenth to the Nineteenth Century: An Essay in Comparative History, Cambridge University Press, Cambridge, 1982.

- Le Roy Ladurie, E., Daux, V., and Luterbacher, J.: Le climat de Bourgogne et d'ailleurs XIV^e–XX^e siècle, *Hist. Econ. Soc.*, 25, 421–436, 2006.
- Leijonhufvud, L.: Grain tithes and manorial yields in early modern Sweden: Trends and patterns of production and productivity c. 1540–1680, Swedish University of Agricultural Sciences, Ulltuna, 2001.
- Ljungqvist, F. C., Seim, A., Krusic, P. J., González-Rouco, J. F., Werner, J. P., Cook, E. R., Zorita, E., Luterbacher, J., Xoplaki, E., Destouni, G., García-Bustamante, E., Aguilar, C. A. M., Seftigen, K., Wang, J., Gagen, M. H., Esper, J., Solomina, O., Fleitmann, D., and Büntgen, U.: European warm-season temperature and hydroclimate since 850 CE, *Environ Res Lett*, 14, 084 015, 2019.
- Ljungqvist, F. C., Thejll, P., Björklund, J., Gunnarson, B. E., Piermattei, A., Rydval, M., Seftigen, K., Støve, B., and Büntgen, U.: Assessing non-linearity in European temperature-sensitive tree-ring data, *Dendrochronologia*, 59, 125 652, 2020.
- Ljungqvist, F. C., Seim, A., and Huhtamaa, H.: Climate and society in European history, *Wiley Interdiscip Rev Clim Change*, 12, e691, 2021.
- Ljungqvist, F. C., Thejll, P., Christiansen, B., Seim, A., Hartl, C., and Esper, J.: The significance of climate variability on early modern European grain prices, *Climetrica*, 16, 29–77, 2022.
- Ljungqvist, F. C., Christiansen, B., Esper, J., Huhtamaa, H., Leijonhufvud, L., Pfister, C., Seim, A., Skoglund, M. K., and Thejll, P.: Climatic signatures in early modern European grain harvest yields, *Climate of the Past*, 19, 2463–2491, 2023.
- Ljungqvist, F. C., Seim, A., and Collet, D.: Famines in medieval and early modern Europe – Connecting climate and society, *Wiley Interdisciplinary Reviews: Climate Change*, 15, e859, 2024.
- Luterbacher, J., Dietrich, D., Xoplaki, E., Grosjean, M., and Wanner, H.: European seasonal and annual temperature variability, trends, and extremes since 1500, *Science*, 303, 1499–1503, 2004.
- Luterbacher, J., Werner, J. P., Smerdon, J. E., Fernández-Donado, L., González-Rouco, F., Barriopedro, D., Ljungqvist, F. C., Büntgen, U., Zorita, E., Wagner, S., Esper, J., Frank, D., Barriendos, M., Bertolin, C., Bothe, O., Brázdil, R., Camuffo, D., Dobrovolný, P., Gagen, M., García-Bustamante, E., Ge, Q., Gómez-Navarro, J., Guiot, J., Hao, Z., Hegerl, G., Holmgren, K., Jungclaus, J., Klimentko, V., Martín-Chivelet, J., McCarroll, D., Pfister, C., Roberts, N., Schindler, A., Schurer, A., Solomina, O., Toreti, A., von Gunten, L., Wahl, E., Wanner, H., Wetter, O., Xoplaki, E., Yuan, N., Zanchettin, D., Zhang, H., and Zerefos, C.: European summer temperatures since Roman times, *Environ Res Lett*, 11, 024 001, 2016.
- Malheiro, A. C., Santos, J. A., Fraga, H., and Pinto, J. G.: Climate change scenarios applied to viticultural zoning in Europe, *Climate Research*, 43, 163–177, 2010.
- Martin, P., Brown, T. A., George, T. S., Gunnarson, B., Loader, N. J., Ross, P., Wishart, J., and Wilson, R.: Climatic controls on the survival and loss of ancient types of barley on North Atlantic Islands, *Climatic Change*, 176, 4, 2023.
- Maurer, C., Hammerl, C., Koch, E., Hammerl, T., and Pokorny, E.: Extreme grape harvest data of Austria, Switzerland and France from AD 1523 to 2007 compared to corresponding instrumental/reconstructed temperature data and various documentary sources, *Theoretical and Applied Climatology*, 106, 55–68, 2011.
- Meier, M., Fuhrer, J., and Holzkämper, A.: Changing risk of spring frost damage in grapevines due to climate change? A case study in the Swiss Rhone Valley, *International Journal of Biometeorology*, 62, 991–1002, 2018.
- Meier, N., Rutishauser, T., Pfister, C., Wanner, H., and Luterbacher, J.: Grape harvest dates as a proxy for Swiss April to August temperature reconstructions back to AD 1480, *Geophysical Research Letters*, 34, L20 705, 2007.
- Miller, G. H., Geirsdóttir, Á., Zhong, Y., Larsen, D. J., Otto-Bliesner, B. L., Holland, M. M., Bailey, D. A., Refsnider, K. A., Lehman, S. J., Southon, J. R., Anderson, C., Björnsson, H., and Thordarson, T.: Abrupt onset of the Little Ice Age triggered by volcanism and sustained by sea-ice/ocean feedbacks, *Geophysical Research Letters*, 39, L02 708, 2012.

- Morales-Castilla, I., de Cortázar-Atauri, I. G., Cook, B. I., Lacombe, T., Parker, A., van Leeuwen, C., Nicholas, K. A., and Wolkovich, E. M.: Diversity buffers winegrowing regions from climate change losses, *Proceedings of the National Academy of Sciences*, 117, 2864–2869, 2020.
- 565 Možný, M., Brázdil, R., Dobrovolný, P., and Trnka, M.: April–August temperatures in the Czech Lands, 1499–2015, reconstructed from grape-harvest dates, *Climate of the Past*, 12, 1421–1434, 2016a.
- Možný, M., Brázdil, R., Dobrovolný, P., Trnka, M., Potopová, V., Hlavinka, P., Zahradníček, P., Štěpánek, P., and Žalud, Z.: Drought reconstruction based on grape harvest dates for the Czech Lands, 1499–2012, *Climate Research*, 70, 119–132, 2016b.
- Mullins, M. G., Bouquet, A., and Williams, L. E.: *Biology of the Grapevine*, Cambridge University Press, Cambridge, 1992.
- 570 Oppenheimer, C.: *Eruptions that Shook the World*, Cambridge University Press, Cambridge, 2011.
- Pauling, A., Luterbacher, J., Casty, C., and Wanner, H.: Five hundred years of gridded high-resolution precipitation reconstructions over Europe and the connection to large-scale circulation, *Climate Dynamics*, 26, 387–405, 2006.
- Pfister, C.: Die Fluktuationen der Weinmosterträge im schweizerischen Weinland vom 16. bis ins frühe 19. Jahrhundert: klimatische Ursachen und sozioökonomische Bedeutung, *Schweizerische Zeitschrift für Geschichte*, 31, 445–491, 1981.
- 575 Pfister, C.: *Klimageschichte der Schweiz 1525–1860. Das Klima der Schweiz und seine Bedeutung in der Geschichte von Bevölkerung und Landwirtschaft*, vol. 2, Bern: Verlag Paul Haupt, 1984.
- Pfister, C. and Wanner, H.: *Climate and Society in Europe: The Last Thousand Years*, Bern: Haupt Verlag, 2021.
- Pfister, C., Brönnimann, S., Altwegg, A., Brázdil, R., Litzemberger, L., Lorusso, D., and Pliemon, T.: 600 years of wine must quality and April to August temperatures in western Europe 1420–2019, *Climate of the Past*, 20, 1387–1399, 2024.
- 580 R Core Team: *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>, 2022.
- Rao, M. P., Cook, E. R., Cook, B. I., Anchukaitis, K. J., D'Arrigo, R. D., Krusic, P. J., and LeGrande, A. N.: A double bootstrap approach to Superposed Epoch Analysis to evaluate response uncertainty, *Dendrochronologia*, 55, 119–124, 2019.
- Robock, A.: Volcanic eruptions and climate, *Rev. Geophys.*, 38, 191–219, 2000.
- 585 Rydval, M., Loader, N. J., Gunnarson, B. E., Druckenbrod, D. L., Linderholm, H. W., Moreton, S. G., Wood, C. V., and Wilson, R.: Reconstructing 800 years of summer temperatures in Scotland from tree rings, *Climate Dynamics*, 49, 2951–2974, 2017.
- Schauberger, P. and Walker, A.: *openxlsx: Read, Write and Edit xlsx Files*, <https://CRAN.R-project.org/package=openxlsx>, r package version 4.2.5, 2021.
- Schneider, L., Smerdon, J. E., Büntgen, U., Wilson, R. J., Myglan, V. S., Kirilyanov, A. V., and Esper, J.: Revising midlatitude summer temperatures back to AD 600 based on a wood density network, *Geophysical Research Letters*, 42, 4556–4562, 2015.
- 590 Schneider, L., Smerdon, J. E., Pretis, F., Hartl-Meier, C., and Esper, J.: A new archive of large volcanic events over the past millennium derived from reconstructed summer temperatures, *Environmental Research Letters*, 12, 094 005, 2017.
- Sigl, M., Winstrup, M., McConnell, J. R., Welten, K. C., Plunkett, G., Ludlow, F., Büntgen, U., Caffee, M., Chellman, N., Dahl-Jensen, D., et al.: Timing and climate forcing of volcanic eruptions for the past 2,500 years, *Nature*, 523, 543–549, 2015.
- 595 Skoglund, M. K.: Farming at the margin: Climatic impacts on harvest yields and agricultural practices in Central Scandinavia, c. 1560–1920, *Agricultural History Review*, 71, 203–233, 2023.
- Squintu, A. A., van der Schrier, G., Brugnara, Y., and Tank, A. K.: Homogenization of daily temperature series in the European Climate Assessment and Dataset, *International Journal of Climatology*, 39, 1243–1261, 2019.
- Steinhilber, F., Beer, J., and Fröhlich, C.: Total solar irradiance during the Holocene, *Geophys Res Lett*, 36, L19 704, 2009.

- 600 Stoffel, M., Khodri, M., Corona, C., Guillet, S., Poulain, V., Bekki, S., Guiot, J., Luckman, B. H., Oppenheimer, C., Lebas, N., et al.: Estimates of volcanic-induced cooling in the Northern Hemisphere over the past 1,500 years, *Nature Geoscience*, 8, 784–788, 2015.
- Stoffel, M., Corona, C., Ludlow, F., Sigl, M., Huhtamaa, H., Garnier, E., Helama, S., Guillet, S., Crampsie, A., Kleemann, K., Camenisch, C., McConnell, J., and Gao, C.: Climatic, weather, and socio-economic conditions corresponding to the mid-17th-century eruption cluster, *Climate of the Past*, 18, 1083–1108, 2022.
- 605 Tingley, M. P., Stine, A. R., and Huybers, P.: Temperature reconstructions from tree-ring densities overestimate volcanic cooling, *Geophysical Research Letters*, 41, 7838–7845, 2014.
- Toohey, M. and Sigl, M.: Volcanic stratospheric sulfur injections and aerosol optical depth from 500 BCE to 1900 CE, *Earth System Science Data*, 9, 809–831, 2017.
- Toohey, M., Krüger, K., Sigl, M., Stordal, F., and Svensen, H.: Climatic and societal impacts of a volcanic double event at the dawn of the Middle Ages, *Climatic Change*, 136, 401–412, 2016.
- 610 Trouet, V. and Van Oldenborgh, G. J.: KNMI Climate Explorer: a web-based research tool for high-resolution paleoclimatology, *Tree-Ring Research*, 69, 3–13, 2013.
- Urhausen, S., Brienen, S., Kapala, A., and Simmer, C.: Climatic conditions and their impact on viticulture in the Upper Moselle region, *Climatic Change*, 109, 349–373, 2011.
- 615 van der Schrier, G., Jones, P., and Briffa, K.: The sensitivity of the PDSI to the Thornthwaite and Penman-Monteith parameterizations for potential evapotranspiration, *Journal of Geophysical Research: Atmospheres*, 116, D03 106, 2011.
- van Engelen, A. and Geurts, H.: Nicolaus Cruquius (1678–1754) and his Meteorological Observations, *Koninklijk Nederlands Meteorologisch Instituut*, 1985.
- van Leeuwen, C., Sgubin, G., Bois, B., Ollat, N., Swingedouw, D., Zito, S., and Gambetta, G. A.: Climate change impacts and adaptations of wine production, *Nature Reviews Earth & Environment*, 5, 258–275, 2024.
- 620 Venios, X., Korkas, E., Nisiotou, A., and Banilas, G.: Grapevine responses to heat stress and global warming, *Plants*, 9, 1754, 2020.
- von Storch, H. and Zwiers, F.: *Statistical Analysis in Climate Research*, Cambridge University Press, Cambridge, 1999.
- Wanner, H., Pfister, C., and Neukom, R.: The variable European Little Ice Age, *Quaternary Sci. Rev.*, 287, 107 531, 2022.
- Wei, T. and Simko, V.: R package ‘corrplot’: Visualization of a Correlation Matrix, <https://github.com/taiyun/corrplot>, (Version 0.92), 2021.
- 625 White, S., Moreno Chamarro, E., Stoffel, M., Zanchettin, D., Huhtamaa, H., Degroot, D., and Corona, C.: The 1600 CE Huaynaputina eruption as a possible trigger for persistent cooling in the North Atlantic region, *Climate of the Past*, 18, 739–757, 2022.
- Wilson, R., Anchukaitis, K., Briffa, K. R., Büntgen, U., Cook, E., D’Arrigo, R., Davi, N., Esper, J., Frank, D., Gunnarson, B., et al.: Last millennium northern hemisphere summer temperatures from tree rings: Part I: The long term context, *Quaternary Science Reviews*, 134, 1–18, 2016.
- 630 Xoplaki, E., Luterbacher, J., Paeth, H., Dietrich, D., Steiner, N., Grosjean, M., and Wanner, H.: European spring and autumn temperature variability and change of extremes over the last half millennium, *Geophysical Research Letters*, 32, L15 713, 2005.
- Yante, J.-M.: Grains et vins des terroirs mosellans de Remich et Grevenmacher (XV^e–XVIII^e siècles), *Revue belge de Philologie et d’Histoire*, 63, 273–309, 1985.
- Yiou, P., García de Cortázar-Atauri, I., Chuine, I., Daux, V., Garnier, E., Viovy, N., Van Leeuwen, C., Parker, A., and Boursiquot, J.-M.: Continental atmospheric circulation over Europe during the Little Ice Age inferred from grape harvest dates, *Climate of the Past*, 8, 577–588, 2012.
- 635