

# Assessing the most severe subsistence crisis of the 18th century in the Northwest of the Iberian Peninsula: a climatological perspective.

Maite deCastro <sup>1\*</sup>, José González-Cao<sup>1</sup>, Nicolás G. deCastro<sup>1</sup>, Juan J. Taboada<sup>2</sup>, José M. Vaquero<sup>3,4</sup>,  
Moncho Gómez-Gesteira<sup>1</sup>

<sup>1</sup> Centro de Investigación Mariña, Universidade de Vigo, Environmental Physics Laboratory (EphysLab), Campus da Auga, 32004, Ourense, Spain.

<sup>2</sup> MeeoGalicia, 15707 Santiago de Compostela, A Coruña, Spain.

<sup>3</sup> Departamento de Física, Universidad de Extremadura, 06800 Mérida, Spain

<sup>4</sup> Instituto Universitario de Investigación del Agua, Cambio Climático y Sostenibilidad (IACYS), Universidad de Extremadura, 06006 Badajoz, Spain

\*Correspondence to: mdecastro@uvigo.gal

## Abstract

The analysis of climate variability over centuries reveals how environmental forces shaped society and helps contextualize modern climate trends and future projections. The persistent and heavy rains across several regions of the Eastern Atlantic in 1768-1769 triggered the last and most severe agricultural crisis in Galicia and Northern Portugal, leading to high mortality. The atmospheric conditions of this historical episode were analyzed using the EKF400v2 paleo-reanalysis dataset, which spans from the 18th century to the early 21st century. From June 1768 to May 1769, the rainfall anomaly in Galicia and Northern Portugal was positive in 11 out of 12 months. Although the rainfall in Northern Portugal appeared less intense than in Galicia, June 1768 had the highest positive rain anomaly of the century, and September 1768 had the second-highest. This excess precipitation agrees with the occurrence of pro-Serenitate rogations and written testimonies indicating an unusually high number of rainy days between June 1768 and May 1769. The atmospheric synoptic patterns for the rainiest months show negative anomalies in both sea level pressure and 500 hPa geopotential height in the northeast Atlantic. These patterns are associated with troughs in the northeastern Atlantic that induce the formation of surface low-pressure systems and hinder the eastward progression of anticyclones into the region, resulting in more frequent episodes of rain and cold than usual.

**Keywords:** precipitation, paleo-reanalysis dataset, ecclesiastical rogations, atmospheric synoptic conditions, Atlantic Arc, agricultural crisis.

## 1. Introduction

The climate and weather conditions play a fundamental role in human health and in the development and evolution of societies, configuring some of their characteristics (Lamb, 1995). The impacts of climate and weather states on societies are complex and interconnected, affecting various aspects of human life. Seasonal variations and their extreme

35 patterns condition the daily lives of individuals, determining clothing, house construction,  
36 food production and consumption, water resources, and social well-being among others.  
37 When frequent deviations from the normal climatic pattern occur, illnesses, economic losses,  
38 and even deaths can result. Climate variability and extreme weather events can affect  
39 agricultural productivity and food availability (White et al., 2018a). Droughts, floods, and  
40 storms can damage crops and livestock, leading to food shortages and insecurity, particularly  
41 in vulnerable regions with limited access to resources.

42 In recent decades, the scientific community has become aware of the importance of going  
43 back in time to deepen our understanding of the climate, as longer data records lead to more  
44 reliable and consistent interpretations of climate (Degroot et al., 2021). Analyzing climate  
45 behavior over centuries allows us to investigate how environmental forces have historically  
46 shaped various sectors of society, analyzing the vulnerabilities generated in different  
47 socioeconomic sectors such as agriculture, transportation, energy, as well as the resilience  
48 and adaptability of society to weather anomalies and climatic dynamics (Fagan, 2001,  
49 Ljungqvist et al., 2020, Pfister, 2021). Given the absence of reliable local or regional details  
50 in climate projections for precipitation and changes in extreme events, identifying similar  
51 patterns from the pre-industrial era could aid in understanding the mechanisms underlying  
52 future extreme hydrometeorological events (e. g. Diodato et al., 2019; Diodado et al., 2020).  
53 Famines are often attributed to the interplay of climate-related and societal stressors within  
54 a framework of pre-existing environmental and social vulnerabilities (Slavin, 2016).  
55 Research on famine crises in medieval and early modern Europe provides a valuable, largely  
56 unexplored archive of societies that faced challenges akin to those of today (Ljungqvist et  
57 al., 2023a). Examining the famines of the 'Little Ice Age' (1300–1800) offers key insights  
58 into human-environment interactions, advancing our understanding of how past societies  
59 managed natural challenges and strengthening the foundation for future decision-making  
60 (Collet and Schub, 2018).

61 The analysis of historical climatic processes predating the industrial era is a highly  
62 challenging task, as it involves handling datasets of diverse origins, including instrumental  
63 data from *in situ* measurements and non-instrumental data obtained from proxies such as  
64 ecclesiastical rogations or written testimonies found in letters, diaries, and reports

65 (Brönnimann, 2015; White et al., 2018b). Additionally, these datasets often vary in terms of  
66 reliability, completeness, and spatial coverage, further complicating the analysis and  
67 interpretation of historical climate patterns. The complexity of this task is compounded by  
68 the need to carefully validate and reconcile disparate sources of historical climate data,  
69 ensuring consistency and accuracy in the analysis. Furthermore, interpreting historical  
70 climate records requires a deep understanding of the context in which the data were collected,  
71 including social, cultural, and environmental factors that may have influenced observations  
72 and recording practices over time. Despite these challenges, studying historical climatic  
73 processes offers valuable insights into long-term climate variability and helps contextualize  
74 modern climate trends and future projections (White et al., 2018b; Pfister et al., 2021).

75 Paleoclimatic reconstructions and modelling approaches (Moravec, 2019) have been  
76 employed over the two past decades to analyze climate history, primarily focusing on  
77 droughts and rainfall patterns across Europe (Murphy et al., 2020; Diodato et al., 2020;  
78 Vicente- Serrano et al., 2020; Noone et al., 2017; Noone et al., 2016; Spraggs et al., 2015;  
79 Brázdil et al., 2015; Todd et al., 2013; Potop et al., 2014) and on drying trends in the  
80 Mediterranean region (Nicault et al., 2008). Additionally, historical documentary data has  
81 enabled a millennium-long reconstruction of damaging hydrological events across Italy and  
82 the broader Mediterranean, revealing 674 events from 800 to 2017 (Diodato et al., 2019). In  
83 particular, numerous historical studies on the Iberian Peninsula have primarily focused on  
84 droughts (Dominguez-Castro et al., 2008; Dominguez-Castro et al., 2012; Dominguez-Castro  
85 et al., 2021; Bravo-Paredes et al., 2020), with relatively limited attention to extreme rainfall  
86 events (Dominguez-Castro et al., 2015). Thus, most of the studies linked to an excess rain  
87 refer to flood linked events (see Gonzalez-Cao et al., 2021; Fernandez-Novoa et al., 2023;  
88 Fernandez-Novoa et al., 2024, Beneyto et al., 20220; Benito et al., 2021; Peña et al., 2022;  
89 among others). Note that, according to the Köppen classification (Köppen, 1884), much of  
90 the southern and Mediterranean Iberian Peninsula experiences a temperate climate with hot,  
91 dry summers (Csa). In contrast, the northwestern Iberian Peninsula and the western coast of  
92 Portugal are classified as having a temperate climate with warm, dry summers (Csb) (see  
93 AEMET-IM, 2011 for further details). Annual precipitation is highly variable across the  
94 Iberian Peninsula (AEMET-IM, 2011). The highest precipitation levels, exceeding 2000 mm,  
95 occur in the mountainous regions of Serra do Gêres in north-eastern continental Portugal,

96 and in areas near the “Rias Baixas” in the southwestern Galicia (northwest of the Iberian  
97 Peninsula). Conversely, the lowest annual rainfalls, below 300 mm, is found in the southeast  
98 of Spain.

99 Multiple records highlight the connection between excessive rainfall and crop losses  
100 throughout history leading to famine both across Europe (Alfani & Ó Gráda 2017; White et  
101 al., 2018a; Ljungqvist et al., 2023a; Ljungqvist et al., 2023b; Slavin, 2016) and more  
102 regionally in Ireland (Ó Gráda, 2017), Great Britain (Hoyle, 2017), France (Béaur & Chevet,  
103 2017), Spain (Pérez-Moneda, 2017) and Northern Portugal (Amorín, 2017; Silva, 2019),  
104 among others. Particularly in Galicia, the biennium of 1768-1769 was characterized by  
105 incessant and heavy rains, resulting in the last and most significant agricultural crisis due to  
106 crop losses (Mejide-Pardo, 1965; Labrada, 1804; González-Fernández, 2000; Losada-  
107 Sanmartín, 2008; Martínez-Rodríguez, 2017), leading to a persistent famine that claimed  
108 human lives (Martín-García, 2001; Losada- Sanmartín, 2008; Silva 2019). This situation,  
109 which historically occurred several times, gave rise to a saying that “*in Galicia, hunger comes*  
110 *swimming*” (Fernández-Cortizo, 2005). The same author analyzes the Galician subsistence  
111 crisis during the 17<sup>th</sup> and 18<sup>th</sup> centuries, identifying that over 67% of the rogations during  
112 these centuries were attributed to an excess of precipitation. A similar situation was observed  
113 in Northern Portugal (Silva 2019). However, famine was not observed in the rest of the  
114 Iberian Peninsula, as documented by multiple sources of data collected in Table 3.2 of Pérez-  
115 Moneda (2017), which accounts for epidemic, death and famines occurring in the Iberian  
116 Peninsula from 1500- 1800, showing the years of excess mortality and famine in 60 small  
117 towns across Castile, Aragon and Extremadura.

118 Regions described above are included in the Atlantic Arc region which refers to a  
119 geographical area encompassing the western and northern coastal regions of Europe that  
120 border the Atlantic Ocean (<https://cpmr-atlantic.org/>). The Atlantic Arc encompasses the  
121 region III (Celtic Seas) and region IV (Bay of Biscay and Iberian Coast) of the OSPAR  
122 Maritime Area ([https://oap.ospar.org/en/ospar-assessments/quality-status-reports/qs-  
123 2023/synthesis-report/introduction/](https://oap.ospar.org/en/ospar-assessments/quality-status-reports/qs-2023/synthesis-report/introduction/)). This area, which typically includes countries such as  
124 Portugal, Spain, France, the United Kingdom, and Ireland, is characterized by its proximity

125 to the Atlantic Ocean and shares similar climatic, environmental, and economic  
126 characteristics due to this coastal influence.

127 The objective of this study is to analyze the atmospheric conditions in the Atlantic Arc from  
128 June 1768 to May 1769, which precipitated the most severe agricultural crisis in Galicia and  
129 Northern Portugal in the 18th century, resulting in high mortality. To achieve this,  
130 precipitation and atmospheric pressure data obtained from a paleo-reanalysis dataset  
131 spanning from the 18th century to the early 21st century will be utilized. Current climate data  
132 generated by ERA5 and precipitation data from a precipitation gauge at Santiago de  
133 Compostela will be used to corroborate that the synoptic conditions observed during that  
134 biennium are reproduced in the present day.

## 135 **2. 1768-1769 Event Identification and Databases**

### 136 **2.1. Identification of the 1768-1769 event**

137 The intense and persistent rainfall event of 1768-1769, which led to a famine in Galicia and  
138 Northern Portugal, resulting in human casualties in excess due to the complete devastation  
139 of crops, was identified through various sources of information with diverse characteristics  
140 and locations. In any case, historical sources verified that the event impacted not only the  
141 Atlantic coast of the Iberian Peninsula but also the entire Atlantic Arc. However, in other  
142 regions, the event did not have as severe consequences on contemporary society as it did in  
143 Galicia and Northern Portugal.


144 There are multiple sources confirming the biennium 1768-1769 as extraordinarily rainy in  
145 the Atlantic European region (red points in Figure 1). Particularly in England, Barker (1771)  
146 identifies 1768 as one of the three rainiest years in the period from 1683 to 1771 in London  
147 (Rutlandshire). Clarck (1999) analyzes the synoptic pattern preceding the major flood that  
148 occurred in Somerset on first of September 1768. Additionally, Macdonald and Sangster  
149 (2017) include the 1768 floods in the historical flood list, although they did not attribute it  
150 significant importance.

151 In France, there are both instrumental records and contemporary testimonies regarding the  
152 abundant rainfall in Bordeaux and in Vendée. Particularly, testimonies from the priest of La  
153 Limouzinière (Vendée) stating, “*This year 1768 has been one of the rainiest that we have*

154 *seen in living memory, the rains began in June and were almost continuous... (Cette année*  
155 *1768 a été une des plus pluvieuses qu'ont ait vu de mémoire d'homme, les pluies ont*  
156 *commencées au mois de juin et ont été presque toujours continuelles...)*“, from the priest of  
157 Lairoux (Vendée) mentioning, *“this year (1768) was remarkable for the abundance of water*  
158 *in the height of summer, which began to fall on the feast day of Saint Médard (June 8) ...*  
159 *(cette année (1768) fut remarquable par l'abondance des eaux au plus fort de l'été qui*  
160 *commencèrent à tomber la fete du dit Saint Médard (June 8)...)*”, and from the prior of Lasse  
161 (Maine-et-Loire) who wrote, *“In the current year (1768), the rains have been so continuous*  
162 *that in the memory of men, they have never seen the like... (Dans la présente année (1768)*  
163 *les pluies ont été si continues que de mémoires d'hommes on en avait vu de pareille...)*”.  
164 Finally, Le Roy Ladurie (2011) discusses the impact of climatic conditions on crops, stating,  
165 *“From 1768 onwards, due to unfavorable weather conditions, too cool and/or too wet, poor*  
166 *harvests and the rise in grain prices became prevalent... (À partir de 1768, en raison de*  
167 *circonstances météo défavorables, trop fraîches et/ou trop humides, les mauvaises récoltes*  
168 *et la hausse des prix frumentaires s'imposent ...)*“, although it is also mentioned that its effect  
169 on mortality was smaller than that observed in 1740.

170 In the Iberian Peninsula, Font-Tullot (1988) identifies 1768 as a particularly rainy year in the  
171 Galico-Cántabra region (northwest of Spain). In the specific case of Galicia, Perez-Constanti  
172 (1925) compiles information from several doctors in Santiago de Compostela on April 17,  
173 1769 who stated, *“...since the month of May of last year 1768, until the present time, it has*  
174 *almost always been raining... as it did in the months of February, March, and April of this*  
175 *year... (...desde el mes de Mayo del año pasado de 68, hasta el tiempo presente, está casi*  
176 *siempre lloviendo ... como lo hizo en los meses de febrero, marzo y abril del presente*  
177 *año...)*”. The same doctors also remarked, *“...it has been eighteen months since we have*  
178 *known the beneficial influences of the seasons, almost continuous rain and cold winds have*  
179 *confused summer, winter, autumn, and spring... (van pasados diez y ocho meses que no*  
180 *hemos conocido los influjos saludables de las estaciones del año, casi continua lluvia y*  
181 *vientos fríos han confundido verano, invierno, otoño y primavera...)*”. The coincidence of  
182 these testimonies with the earlier ones described by the French priests is striking. Lastly, in  
183 Northern of Portugal, Silva (2019), through an annual precipitation index (Fig. 23 of his  
184 thesis), indicated that the end of summer and the fall of 1768 were characterized by high

185 amounts of rain, serving the prelude to a severe agrarian crisis. Additionally, Amorín (2017)  
186 identifies severe floods in the Porto region due to continuous rains in 1768-1769.



187  
188 **Figure 1:** Atlantic Arc region, encompassing Portugal, Spain, France, England and Ireland.  
189 Red points indicate all instrumental or documentary testimony collected regarding the  
190 extraordinary rainy event of 1768-1769.

191 These specific climatic conditions were reflected in numerous ecclesiastical rogations “*pro*  
192 *Serenitate*” held in various locations in Galicia and Northern Portugal (Silva, 2019; González  
193 Fernández, 2000; Losada-Sanmartín, 2008). These authors have referred to the crisis of 1768-  
194 1769 as one of the two most severe in the 18<sup>th</sup> century, accompanied by episodes of hunger  
195 and excess mortality throughout the region, as documented in numerous studies (Ávila and  
196 LaCueva, 1852; Mejjide-Pardo, 1965).

197 Other studies emphasize the impact of the extraordinary climatic conditions on cereal harvest  
198 (Pérez-Costanti, 1925; Mejjide-Pardo, 1965; Martínez-Rodríguez, 2017), which is also  
199 reflected in tithe records (Eiras, 1978).

## 200 **2.2. Databases**

201 Historical data of precipitation, sea level pressure (SLP), and geopotential height at 500 hPa  
202 (GPH) at a monthly scale were obtained from the EKF400v2 paleo-reanalysis database with  
203 approximately 2° spatial resolution (Valler et al., 2022). According to these authors, the  
204 EKF400v2 utilizes atmospheric-only general circulation model simulations (CCC400). The

205 30 ensemble members were generated with the ECHAM5.4 general circulation model. These  
206 simulations are augmented by a significantly expanded observational network comprising  
207 early instrumental temperature and pressure data, documentary evidence, and proxy records  
208 derived from tree-ring width and density. Additionally, new types of observations, including  
209 monthly precipitation amounts, the frequency of wet days, and coral proxy records, have  
210 been incorporated into the assimilation process. In this version 2 system, the assimilation  
211 procedure has undergone methodological enhancements, notably the estimation of the  
212 background-error covariance matrix through a blending technique involving both time-  
213 dependent and climatological covariance matrices. The EKF400v2 model simulations cover  
214 the period from the beginning of 18th century to the beginning of the 21st century. For further  
215 de details, the reader is referred to Valler et al. (2020).

216 Two additional long-term regional precipitation series were considered. For Ireland, the  
217 Island of Ireland 1711 (IoI\_1711) series, was used, providing continuous monthly  
218 precipitation data from 1711 to 2016 (Murphy et al., 2018). The post-1850 series was  
219 constructed using quality-assured monthly precipitation records compiled by Noone et al.  
220 (2016), while the pre-1850 series was derived from instrumental and documentary sources  
221 compiled by the UK Met Office (Jenkinson et al., 1979). The monthly IoI series was accessed  
222 from PANGAEA (<https://doi.pangaea.de/10.1594/PANGAEA.887593>). For Wales-  
223 England, the England and Wales Precipitation (EWP) series (Alexander and Jones, 2001;  
224 Simpson and Jones, 2014) were considered. These series represent an area-averaged  
225 precipitation record derived from five rainfall regions representing England and Wales. It  
226 provides a continuous monthly precipitation record from 1766 and is regularly updated by  
227 the UK Met Office (UKMO) Hadley Centre, from whom monthly data were accessed  
228 (<https://www.metoffice.gov.uk/hadobs/hadukp/>). Both data sets were combined by Murphy  
229 et al (2020) to reconstruct and analyze monthly precipitation in England and Wales, Scotland  
230 and Ireland, and to reevaluate historical droughts over the period 1748-2000. Notably, the  
231 overlooked drought of 1765–1768, which impacted the British-Irish Isles, was identified as  
232 the most significant event in their 250-year reconstruction. This event can serve as a valuable  
233 benchmark for stress- testing current systems to ensure resilience.




234 *In situ* monthly precipitation data were obtained from precipitation gauges located at Lyndon  
235 and Cornwall in England, and Bordeaux in France. The precipitation series at Lyndon spans  
236 from 1737 to 1770, while at Bordeaux it covers from 1751 to 1770, and in Cornwall from  
237 1767 to 1771. Moreover, the cumulative number of rainy days in Exeter, England, from 1755  
238 to 1775 was obtained from Exeter weather diaries, accessible at  
239 [https://digital.nmla.metoffice.gov.uk/IO\\_11c660bd-60c1-4d59-a079-64fdbdb20144](https://digital.nmla.metoffice.gov.uk/IO_11c660bd-60c1-4d59-a079-64fdbdb20144).

240 Current daily precipitation data in Galicia were obtained from a rain gauge located at  
241 Santiago de Compostela (42° 53' 17''N, 8° 24' 30''W), available at  
242 [https://www.aemet.es/es/datos\\_abiertos](https://www.aemet.es/es/datos_abiertos). This rain gauge is one with the longest precipitation  
243 series in Galicia from 1944 to 2023. Additionally, monthly sea-level pressure and  
244 geopotential height at 500 hPa were retrieved from ERA5 database  
245 ([https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels-monthly-](https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels-monthly-means?tab=form)  
246 [means?tab=form](https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels-monthly-means?tab=form) ) at a spatial resolution of 0.5° covering the region from 60°N to 25°N and  
247 from 5°E to 45°W for the period 1940-2023.

248 The cumulative annual precipitation for each month was calculated by considering the  
249 preceding six months and the subsequent five months relative to the month under study. The  
250 same methodology was applied for calculating the number of rainy days per month. This  
251 approach enables the determination of both the cumulative precipitation and the number of  
252 rainy days per year without relying on calendar years.

### 253 **3. Results**

254 The precipitation anomaly was calculated using EKF400v2 data for the period of maximum  
255 cumulative rainfall 6/1768-5/1769, documented in the data sources described in previous  
256 section, relative to the annual mean for the period 1755- 1785 (Fig. 2). The precipitation  
257 anomaly reached values of 200 mm in French Brittany and southern England, and values of  
258 approximately 150 mm in Galicia and North Portugal, where typical annual precipitation  
259 ranges from 1000 to 1200 mm (AEMET-IM, 2011).




260

261 **Figure 2:** Precipitation anomaly (mm) during the period of maximum cumulative rainfall  
 262 (6/1768-5/1769) relative to the annual mean for period 1755- 1785.

### 263 3.1 Analyzing the historical persistent and heavy rainfall event

264 The cumulative annual precipitation provides by EKF400v2 data in Galicia (Fig. 3a) and  
 265 Northern Portugal (Fig. 3b) over the period 1755 to 1785 shows a peak during the final six  
 266 months of 1768 and the initial six months of 1769, with values exceeding 1200 mm.



267

268 **Figure 3:** Cumulative annual precipitation (mm) in Galicia (a) and Northern Portugal (b).  
 269 Data obtained from EKF400v2 paleo-reanalysis database. Red arrow marks the 1768- 1769  
 270 precipitation peak.

271 This persistent and heavy rainfall event was also observed in neighboring regions such as  
 272 Ireland (Fig. 4a), Wales-England (Fig. 4b) and Normandy and French Brittany (Fig. 4c),  
 273 where similar peaks in cumulative precipitation occurred simultaneously.


274 Figure 4a illustrates the precipitation in Ireland for the decades just before and after the  
 275 extreme rain event observed from mid-1768 to mid-1769 in NW Iberian Peninsula. The  
 276 drought period of 1765- 1768, as analyzed by Murphy et al. (2020), immediately precedes

277 the peak rainfall of 1768-1769. While the period 1768-1769 appears significantly rainy, it is  
278 comparable to other rainy events identified in preceding and subsequent decades. These  
279 findings align with those derived from Murphy's reconstructed database for Ireland, Murphy  
280 et al., (2020), as depicted in Figure 5a.

281

282

283




284 **Figure 4:** Cumulative annual precipitation (mm) in Ireland (a), Wales-England (b) and  
285 Normandy and French Brittany (c). Data obtained from EKF400v2 paleo-reanalysis  
286 database. Red arrow marks the 1768- 1769 precipitation peak.

287 In Wales-England (Fig. 4b), similar to the case of Ireland, a peak in cumulative precipitation  
288 between 1768 and 1769 is observed following the drought of 1765-1768. Once again, the

289 precipitation peak is comparable to that observed at the beginning of the decade. As was the  
290 case with Ireland, the results are comparable to those from the reconstructed database for  
291 Wales-England (Alexander and Jones, 2001; Simpson and Jones, 2014) used in Murphy et  
292 al., (2020) as depicted in Figure 5b.


293 In France (Fig. 4c), it is also evident that the peak in cumulative precipitation spanning from  
294 1768 to 1769 is the highest of the period under study (1755-1785), although it is not as  
295 pronounced as the peak identified in Galicia and Northern Portugal.

296



297 **Figure 5:** Cumulative annual precipitation (mm) in Ireland (a) and Wales-England (b). Data  
298 from Murphy et al., (2018) for Ireland and from Alexander and Jones, (2001) and Simpson  
299 and Jones (2014) for Wales-England. Red arrow marks the 1768- 1769 precipitation peak.


300 Similar information can be obtained from local *in situ* precipitation gauges, such as illustrated  
301 in Figure 6 for Lyndon and Cornwall in England, (Figs. 6a and b, respectively) and Bordeaux  
302 in France (Fig. 6c).



303 **Figure 6:** Cumulative annual precipitation (mm) in (a) Lyndon (England), (b) Cornwall  
 304 (England), and (c) Bordeaux (France). Data derived from local *in situ* precipitation gauges.  
 305 Red arrow marks the 1768- 1769 precipitation peak.

306 *In situ* data corroborate the presence of a peak in rainfall at these locations between 1768 and  
 307 1769. Unfortunately, the limited duration of the precipitation series restricts our ability to  
 308 gain a broader perspective.


309 In the same line, the cumulative number of rainy days calculated from *in situ* precipitation  
 310 data at Exeter (England) from 1755 to 1775 demonstrates a notable positive anomaly of  
 311 between 20 and 40 days between mid- 1768 and mid-1769 (Fig. 7).



312

313 **Figure 7:** Cumulative number of rainy days at Exeter (England). Data derived from local *in*  
 314 *situ* precipitation gauges.

315 The origin of this precipitation anomaly pattern can be analyzed in terms of the anomaly in  
 316 SLP (Fig. 8a) and in GPH (Fig. 8b), for the region under study during the period from 6/1768  
 317 to 5/1769, relative to the annual mean for the period 1755- 1785.



318 **Figure 8:** (a) SLP anomaly (hPa) and (b) 500 GPH anomaly (m) during the period 6/1768-  
 319 5/1769, relative to the annual mean for the period 1755- 1785. Data obtained from EKF400v2  
 320 paleo-reanalysis database.

321 Both subplots depict a negative anomaly minimum of approximately 3 hPa in the SLP (Fig.  
 322 8a) and of 30 m in the GPH (Fig. 8b), locating the area of strongest anomaly (negative) in

323 the Bay of Biscay and covering the westernmost part of Europe inside the low anomalies  
324 area.

### 325 **3.2 Analyzing current persistent and heavy rainfall events**

326 After identifying the synoptic conditions that led to the extraordinary rainfall during the  
327 period 6/1768 to 5/1769, the next step will be to analyze whether similar patterns have been  
328 observed over the past 80 years, during which abundant instrumental records facilitate the  
329 identification of unusual rainfall events. Considering that documentary records point out the  
330 presence of incessant rains over the period 1768-1769, the number of rainy days per month  
331 was calculated from 1944 to 2023 using data from the Santiago de Compostela rain gauge.  
332 A day was considered rainy when at least 1 mm of precipitation was collected (AEMET-IM,  
333 2011). The number of days corresponding to the mean and the 50<sup>th</sup> (median), 90<sup>th</sup>, and 95<sup>th</sup>  
334 percentiles is shown in Table 1. Note that Santiago de Compostela was one of the places most  
335 affected by the 1768-1769 and has one of longest meteorological series in the area of study,  
336 which makes it an optimal candidate to analyze how current patterns can be related to those  
337 obtained almost three centuries ago.

338 **Table 1.** Number of days corresponding to mean and the 50<sup>th</sup> (median), 90<sup>th</sup>, and 95<sup>th</sup>  
339 percentiles of rainiest days per month from 1944 to 2023 using data from the Santiago de  
340 Compostela rain gauge.


Month	Mean	50 <sup>th</sup>	90 <sup>th</sup>	95 <sup>th</sup>
1	16.5	17	25	26
2	14.1	14	23	25
3	14.3	15	24	25
4	13.7	14	20	23
5	12.4	12	19	21
6	8.0	8	14	15
7	5.3	5	10	10
8	6.8	7	12	14
9	9.2	9	16	17
10	13.9	14	21	24
11	15.5	15	23	26
12	16.0	16	25	27

341

342 The 50<sup>th</sup> percentile serves as a reference on the number of rainy days per month in a normal  
343 year, while the 90<sup>th</sup> and 95<sup>th</sup> percentile provide information on the number of rainy days per  
344 month in extreme years. In fact, the total number of days in Santiago de Compostela during  
345 a normal year (50<sup>th</sup> percentile) with precipitation equal to or greater than 1 mm is 146 days.  
346 This is similar to the value reported in the Iberian Climatic Atlas (AEMET-IM, 2011), which  
347 states that the highest number of days with precipitation equal to or greater than 1 mm (over  
348 150 days) in the Iberian Peninsula occurs, among other regions, in the northeastern Galicia.  
349 Additionally, the 95<sup>th</sup> percentile is exceeded by 23 days of precipitation per month in seven  
350 months, which include January to April and October to December (JFMAOND). In the  
351 summer months, the 95<sup>th</sup> percentile for June, July, and August is 15, 10, and 14 days,  
352 respectively.

353 The precipitation observed during the rainiest months (95<sup>th</sup>) over the recent period 1944 to  
354 2023 was analyzed using composite maps. Initially, the mean composite map for SLP and  
355 GPH was calculated from 1944 to 2023. Subsequently, the composites maps (SLP and GPH)  
356 for extreme rainy conditions were determined as follows: i) the monthly 95<sup>th</sup> percentiles of  
357 the rainiest days presented in Table 1 served as threshold values; ii) for each month, its  
358 composite maps were generated by averaging SLP or GPH only for the years when the  
359 number of rainy days in that month exceeded the threshold; iii) the monthly composite maps  
360 were then averaged to obtain the annual composite maps corresponding to rainy months.  
361 Finally, the mean annual composite map was subtracted from the rainy composite map to  
362 yield the anomaly. The mean composite from 1944 to 2023 and anomaly maps for SLP and  
363 GPH are illustrated in Figure 9, with the left column representing SLP and the right column  
364 GPH. The mean composite map for SLP (Fig. 9a) was subtracted from the SLP composite  
365 map for rainy months (Fig. 9c) to obtain the SPL anomaly (Fig. 9e). Similarly, the GPH  
366 anomaly (Fig. 9f) was obtained by subtracting Fig. 9b from Fig. 9d.





367 **Figure 9:** (a) Annual SLP composite (hPa) from 1944 to 2023, (b) Annual GPH composite  
 368 (m) from 1944 to 2023, (c) Annual SLP composite during the rainiest months (exceeding the  
 369 95<sup>th</sup> percentile for that month), (d) Annual GPH composite during the rainiest months  
 370 (exceeding the 95<sup>th</sup> percentile of that month), (e) Annual SLP composite of anomalies,  
 371 calculated as the difference between subplots c and a, (f) Annual GPH composite of  
 372 anomalies, calculated as the difference between subplots d and b.

373 The synoptic patterns shown in Figure 9 are similar to the ones obtained during the 1768-  
 374 1769 rainy event (Fig. 8) with the Iceland low anomaly low and displaced southeastward  
 375 over the Bay of Biscay.

#### 376 4. Discussion

377 The period from June 1768 to May 1769 was characterized by incessant and heavy rains in  
378 the northwestern region of the Iberian Peninsula, resulting in the last and most significant  
379 agricultural crisis due to crop losses and leading to a persistent famine that claimed human  
380 lives. During these years, Spain was a country immersed in Bourbon reformism and, in  
381 particular, in the reforms led by King Charles III, which were characterized by enlightened  
382 ideas, as long as these did not endanger his absolute power and the traditional social order  
383 (enlightened absolutism). In 1766, a strong crisis occurred that triggered the so-called  
384 "Esquilache Riot", largely motivated by a subsistence crisis as a result of a very sharp rise in  
385 the price of bread. This rise in the price of bread was motivated by a combination of poor  
386 harvests and the promulgation of a decree in 1765 that liberalized the grain market  
387 (Domínguez Ortiz, 2005).

388 The poverty and low level of socioeconomic development in the northwestern region of the  
389 Iberian Peninsula were also contributing factors to the absence of instrumental  
390 measurements, which were already incipient at other European locations during the period  
391 of interest. The first instrumental readings of the weather in Galicia were located in El Ferrol  
392 in 1788 (Domínguez-Castro et al. 2014). Additionally, the first instrumental meteorological  
393 readings in northern Portugal were made by Joao da Veiga in Lamego, from 1770 until 1784  
394 (Alcoforado et al. 2012). This lack of instrumental information was partially mitigated by  
395 utilizing other documentary sources such as rogation ceremonies, convent diaries, letters,  
396 which allowed for the categorization of meteorological events following the method  
397 proposed by Pfister (1984, 1992). The ecclesiastical rogations "*pro-Serenitate*" constitute a  
398 fundamental source of information used to characterize the historical rainy event in Galicia  
399 and Northern Portugal (Fernandez-Cortizo, 2005; Silva, 2019) complementing the written  
400 testimonies previously described (Silva, 2019). In particular, the ecclesiastical rogation  
401 database corresponding to Santiago de Compostela, possibly the ground zero of the event in  
402 terms of deaths and socio-economic impact, contains 283 rogation masses over the period  
403 1670-1804 (approximately 2 per year), among which 70 were for rain (*pro-Pluvia*) and 181  
404 for fair weather (*pro-Serenitate*). This strongly contrasts with observations in other parts of  
405 the Iberian Peninsula, where *pro-Serenitate* rogations are less common (Dominguez-Castro

406 et al., 2008; Dominguez-Castro et al., 2012; Dominguez-Castro et al., 2021; Bravo-Paredes  
 407 et al., 2020) due to particular climate conditions that characterize the NW corner of the  
 408 Iberian Peninsula. For comparison purposes, Table 2 exhibits the current annual precipitation  
 409 levels at the most rainfall-prone locales within the Atlantic Arc, encompassing Santiago de  
 410 Compostela (Spain) and Porto (Portugal), both situated within the designated area of interest.  
 411 Table 2: Current annual rainfall at the rainiest locations in the Atlantic-Arc averaged over the  
 412 period 1991-2021. Source <https://es.climate-data.org/>.

City	Annual Rainfall (mm)
Brest (France)	941
Cardiff (Wales, UK)	1071
Manchester (England, UK)	1047
Londonderry (Northern Ireland, UK)	1102
Galway (Ireland)	1117
Santiago de Compostela (Spain)	1242
Porto (Portugal)	1285

413

414 During the decade encompassing the event (from 1761 to 1770), 20 *pro-Serenitate* and 6 *pro-*  
 415 *Pluvia* rogations were celebrated in Santiago, which aligns with the average during the longer  
 416 period (1670-1804) mentioned above. However, the summer of 1768 stood out for the  
 417 frequency of rogations for fair weather, with four ceremonies held in Santiago from June to  
 418 August of that year. In Pontevedra, located 60 km south of Santiago, *pro-Serenitate* rogations  
 419 occurred in May and September 1768 and May 1769. Similarly, rogations for fair weather  
 420 were documented in Braga (North Portugal) in September and October 1768 (Silva, 2019).  
 421 The same author (refer to page 214) indicates that 1768 witnessed the highest number of *pro-*  
 422 *Serenitate* rogative processions in the 17th-18th centuries for northern Portugal. Another  
 423 noteworthy aspect highlighting the intensity of the 1768-1769 event is that in the city of  
 424 Santiago de Compostela during the historical record (1670-1804), *pro-Serenitate* rogations  
 425 took place only on six occasions in two summer months of the same year, 1768 (June and  
 426 August) being one of those years.

427 This high precipitation event was not confined exclusively to this area of the Iberian  
 428 Peninsula but extended to other areas of France, Wales, England, and Ireland, although in

429 these regions, it did not lead to agricultural and demographic crises. This may be attributed  
430 to the proactive measures taken in other areas such as France, following the "Great Winter"  
431 of 1709, where strategies like product substitution were adopted. As a result, wheat was  
432 replaced by less prized substitutes such as buckwheat, rye, and chestnuts (Béaur and Chevet,  
433 2017). The introduction of buckwheat in western France is believed to have contributed to  
434 the region's relatively mild impact during the great famines of the eighteenth century  
435 (Nassiet, 1998). Similarly, in the UK, some authors (Hoyle 2017) suggest that the climatic  
436 variability of the early eighteenth century may have prompted the cultivation of root crops in  
437 fields as an emergency fodder crop. By the late 1720s, potatoes had become a common part  
438 of the diet among the poor. Nevertheless, there remains the possibility that famine was  
439 averted because potatoes, like oats, provided the option for people to switch to cheaper, albeit  
440 less desirable, foodstuffs during years of high prices. In Ireland, potato had become the base  
441 of the diet as the popular saying stated "*potatoes in the morning, potatoes at noon; and if I*  
442 *rose at midnight, it would still be potatoes (ditty prátaí ar maidin, prátaí um nóin; is dá n-*  
443 *éireoinn meánoíche, prátaí gheobhainn)*".

444 In Galicia and North Portugal, all sources indicate a severe famine. To comprehend the  
445 diverse implications of the historical rainy event on the societies on these regions, it is  
446 imperative to understand the socio-economic context of Galicia and North Portugal at that  
447 time. One reason for the famine in this area stemmed from the predominant reliance on wheat  
448 and rye crops during that period. Traditionally, the most agriculturally productive regions of  
449 the Iberian Peninsula were the south and center, where Mediterranean agriculture thrived,  
450 particularly with the cultivation of wheat. Conversely, the north faced challenges due to its  
451 humid climate, which posed difficulties for staple crops such as olive trees and vines to adapt.  
452 The introduction of crops from the Americas significantly transformed the agricultural and  
453 commercial landscape. For the northern regions, the emergence of potatoes and corn  
454 provided a solution to their historical agricultural constraints. However, by 1768-1769, these  
455 new crops had not yet been widely adopted. Corn, native to the Americas, arrived in Europe  
456 around 1604, initially being cultivated in Cantabria (NW, Spain). Despite its early  
457 introduction, corn initially faced resistance and was primarily used as fodder. Similarly, the  
458 potato, encountered by Spanish conquistadors in the Andean regions in the mid-16th century,  
459 was initially disregarded as food and used primarily for animals and ornamental purposes

460 until the early 18th century. Consequently, the widespread acceptance and culinary use of  
 461 potatoes, as exemplified by the Spanish potato omelet, did not occur until the late 18th  
 462 century, 1798 (López Linage, 2008). Table 3 provides details on the planting and harvesting  
 463 seasons for various crops in the current area of interest.

464 Table 3: Planting and harvesting periods for various crops are exemplified by the provinces  
 465 of A Coruña and Pontevedra in Galicia, which were the most affected areas  
 466 ([https://www.mapa.gob.es/es/estadistica/temas/estadisticas-agrarias/01-calendariosiembra-](https://www.mapa.gob.es/es/estadistica/temas/estadisticas-agrarias/01-calendariosiembra-nuevo-sencilla-1_tcm30-514260.pdf)  
 467 [nuevo-sencilla-1\\_tcm30-514260.pdf](https://www.mapa.gob.es/es/estadistica/temas/estadisticas-agrarias/01-calendariosiembra-nuevo-sencilla-1_tcm30-514260.pdf)).

<b>Crop</b>	<b>Planting</b>	<b>Harvesting</b>
Wheat	Apr	Jul
Rye	Oct-Nov	May-Jun
Corn	Apr-May	Sep-Oct
Fodder Corn	Jun	Sep
Potato	Oct-Nov	Jan-Apr
	Dec-Feb	Apr-Jun
	Mar-Apr	Sep-Oct
	Jul-Aug	Nov-Dec

468

469 It is apparent that an agricultural system reliant exclusively on cereals (wheat and rye) is  
 470 more prone to encountering subsistence crises in comparison to one integrating  
 471 supplementary crops. Such a system, predominantly centered on wheat and rye, exhibits  
 472 heightened vulnerability to heavy rainfall during late spring and early summer.

473 Nearly contemporary authors like Labrada, (1804) highlight that it was the famines of 1768  
 474 and subsequent years that forced the poorest peasants to sow and eat potatoes, which were  
 475 previously only consumed by pigs. Additionally, Meijide-Pardo (1965), recounts that the  
 476 copious and continuous rains during the summer of 1768 ruined almost the entire wheat and  
 477 rye crops in all the provinces existing in Galicia at that time. This situation worsened at the  
 478 beginning of autumn when the corn harvest, which was the main resource in rural areas,  
 479 failed. Consequently, by mid-May 1769, the price increase compared to that of the previous  
 480 3 years was 141% for wheat, 181% for rye and 173% for corn highlighting that the local

481 authorities aid was less than expected (Martínez-Rodríguez, 2017). Furthermore, the  
482 situation exacerbated due to the lack of repaired roads or adequate means of transportation  
483 to distribute foreign grain. All food transportation was carried out using rudimentary carts  
484 and horses. It should also be noted that part of the livestock was fed with grain, which led to  
485 a cascading effect. This agricultural crisis resulted in an influx of poor people, as documented  
486 by Ávila and LaCueva (1852), “*Since the beginning of the year 1769, there was a great*  
487 *famine due to the scarcity of grain resulting from the heavy rains of the previous year, from*  
488 *which countless poor people descended from the mountains to this City; many died of extreme*  
489 *necessity (desde principios del año de 1769, se padeció una muy grande hambre por la*  
490 *escasez de frutos de todos los granos que hubo en el año anterior a causa de las muchas*  
491 *llubias que sobrevinieron en él, de cuyas resultas bajaron de las montañas a esta Ciudad*  
492 *infinidad de pobres; murieron muchos de suma necesidad)”. The mortality crisis that*  
493 *occurred as a result of this agricultural crisis in Galicia is also documented by Martín-García*  
494 *(2001), who states, “The famous crisis of 1768-1769 affected practically all of Galicia and*  
495 *had its prelude in the poor harvests of 1768, caused by incessant rains, which were the*  
496 *breeding ground for famines and epidemics (La famosa crisis de 1768-1769 castigó a la*  
497 *práctica totalidad de Galicia y tuvo su prólogo en las pésimas cosechas de 1768, provocadas*  
498 *por las incesantes lluvias, que fueron el caldo de cultivo de hambrunas y epidemias)” and by*  
499 *Silva (2019) in northern Portugal.*

500 Data assimilation techniques have gained popularity in the field of climate reconstruction, as  
501 they estimate historical climate states by integrating observational data and model  
502 simulations. The EKF400v2 paleo-reanalysis database (Valler et al, 2022) spanning from the  
503 18th century to the early 21st century enabled us to reconstruct the historical rainy event of  
504 1768-1769. Rain anomalies relative to the century (1701-1800) can provide valuable insights  
505 into the singularity of the event.

506 In Galicia, during the period from June 1768 to May 1769, the rain anomaly was positive in  
507 11 out of 12 months, with March 1769 being the only exception. Additionally, June 1768  
508 exhibited the highest positive rain anomaly of the century, and September 1768 had the  
509 second-highest positive rain anomaly. These findings align well with the occurrence of *pro-*  
510 *Serenitate* rogations in Santiago de Compostela. Similarly, in North Portugal, over the same

511 period, the rain anomaly was also positive in 11 out of 12 months, with March 1769 being  
512 the only exception. However, the rain event appeared to be less intense, with only February  
513 1769 presenting the second- highest positive anomaly of the century for that month, and  
514 September 1768 corresponding to the fifth-highest positive anomaly. Furthermore, June 1768  
515 had the highest positive rain anomaly of the century, and September 1768 had the second-  
516 highest positive rain anomaly. According to Silva (2019), rogation ceremonies took place in  
517 September and October 1768 in Braga (North Portugal). The same author created a  
518 classification by assigning numerical values between 0 and  $\pm 1$  to each season of the year,  
519 with +1 indicating an excess (rainy season), -1 indicating a deficit (dry season), and 0  
520 indicating "normal" seasons. Consequently, the summer and autumn of 1768 and the spring  
521 of 1769 are classified with an index of 1.

522 Written testimonies indicate an unusually high number of rainy days between June 1768 and  
523 May 1769, however the lack of instrumental historical data in Galicia and North Portugal  
524 hinders our ability to estimate the number of rainy days. To discern the significance of an  
525 unusually high number of rainy days, the number of days corresponding to the 50<sup>th</sup>, 90<sup>th</sup> and  
526 95<sup>th</sup> percentiles of rainiest days per month were analyzed using data from the Santiago de  
527 Compostela rain gauge from 1944 to 2023. Remarkably, from the analysis of the current  
528 precipitation data, it is evident that over an eighty-year period, there were three natural years  
529 with more than five months experiencing precipitation exceeding the 90<sup>th</sup>. These natural  
530 years include November to December 1950, February, May, and August 1951; July,  
531 September, and November 1965, to January, February, April, and June 1966, and finally,  
532 April, November, and December 2000 to January and March 2001. This fact clearly  
533 demonstrates that using a limited record (only 80 years), the chances of having extreme rainy  
534 years, with several months experiencing a high number of rainy days, are not negligible.  
535 Additionally, the composite of the rainiest months, those that exceed the 95<sup>th</sup> percentile of  
536 that month, exhibits synoptic patterns similar to those obtained during the 1768-1769 rainy  
537 event. Synoptic patterns obtained from the ERA5 database for the wettest months (Figures  
538 8a and b) show negative anomalies in both SLP and 500GPH in the northeast Atlantic. This  
539 type of anomaly is normally associated with a circulation in which the jet stream adopts a  
540 very meridional mode. These meridional modes exhibit greater persistence compared to the  
541 zonal ones. This persistence leads to the association with significant anomalies, as observed

542 in this study. Regions situated within the colder sector of the circulation experience  
543 continuous influx of low-pressure systems traveling along the jet stream. This is evident in  
544 Figure 8, particularly over the NW Iberian Peninsula, accounting for the notable surplus in  
545 rainy days. Conversely, areas farther east or west may experience prolonged periods of  
546 anticyclonic influence, resulting in reduced rainfall.

547 Figures 8a and 8b, depicting SLP and 500GPH data extracted from the EKF400v2 paleo-  
548 reanalysis database for the 1768-1769 rainy event, closely resemble those obtained from  
549 contemporary records (Figures 9e and 9f). This similarity allows us to interpret the  
550 atmospheric circulation dynamics during this event. It is likely that a pronounced planetary  
551 circulation pattern, predominantly influenced by meridional modes in the northern  
552 hemisphere, contributed to the frequent occurrence of troughs in the northeastern Atlantic.  
553 These troughs, characterized by cold air in the mid-to-upper atmospheric layers, induce the  
554 formation of surface low-pressure systems. Additionally, they hinder the eastward  
555 progression of anticyclones into the region, resulting in more frequent episodes of rain and  
556 cold than usual.

557 Moreover, broadening the analysis to encompass other geographical regions on the map  
558 enables us to elucidate why this event primarily impacted areas in Portugal, northwest Spain,  
559 parts of France, and the British Isles, while sparing other regions in Europe. The trough  
560 depicted in Figures 9c and 9f encompasses all the affected areas during this event. However,  
561 as elucidated in the preceding paragraph, regions lying beyond its influence are not subjected  
562 to the frequent arrival of low-pressure systems and thus remain unaffected by excessive  
563 rainfall. This is exemplified by the central and eastern regions of Spain.

564

## 565 **5. Conclusions**

566 The incessant and heavy rainfall in several regions of the Atlantic Arc (Ireland, England,  
567 France, Galicia and Northern Portugal) over the period June 1768- May 1769 precipitated  
568 the last and most severe agricultural crisis in Galicia and Northern Portugal, resulting in high  
569 mortality. The atmospheric conditions that led to this historical episode were analyzed using



570 the EKF400v2 paleo-reanalysis dataset, which spans from the 18th century to the early 21st  
571 century. The following main conclusions were obtained:

572 - The rainfall anomaly in Galicia and North of Portugal from June 1768 to May 1769 was  
573 positive in 11 out of 12 months, with March 1769 being the only exception. Although the  
574 rainfall event in North Portugal appeared to be less intense than in Galicia, June 1768  
575 exhibited the highest positive rain anomaly of the century, and September 1768 had the  
576 second-highest positive rain anomaly.

577 - This excess precipitation aligns well with the occurrence of *pro-Serenitate* rogations in  
578 Santiago de Compostela and Braga, and with written testimonies indicating an unusually high  
579 number of rainy days between June 1768 and May 1769. Additionally, the excess mortality  
580 in 1769 and 1770, which is documented in different sources, highlights the unusual nature of  
581 the event.

582 - The atmospheric synoptic patterns for the rainiest months show negative anomalies in both  
583 SLP and 500GPH in the northeast Atlantic. These patterns are associated with a pronounced  
584 planetary circulation predominantly influenced by the meridional mode of the jet stream in  
585 the northern hemisphere. This circulation contributes to the frequent occurrence of troughs  
586 in the northeastern Atlantic, which induce the formation of surface low-pressure systems and  
587 hinder the eastward progression of anticyclones into the region, resulting in more frequent  
588 episodes of rain and cold than usual.

589

#### 590 *Credit Author Statement*

591 **Maite deCastro:** Conceptualization, Methodology, Formal Analysis, Writing – Original  
592 Draft, Writing – Review, Editing, Elaboration of tables, visualization. **José González-Cao:**  
593 Writing – Review, Editing, Formal Analysis. **Nicolás G. deCastro:** Software, Validation and  
594 comparison between different data sources, elaboration and revision of tables, Writing –  
595 Review. **Juan J. Taboada:** Writing – Review, Editing, Formal Analysis. **Jose M. Vaquero:**  
596 Writing – Review, Formal Analysis. **Moncho Gómez-Gesteira:** Conceptualization,  
597 Methodology, Supervision, Software, Graphing.

#### 598 *Acknowledgments*

599 The authors would like to acknowledge funding from the Xunta de Galicia under project  
600 ED431C 2021/44 (Grupos de Referencia Competitiva), and from the Interreg POCTEP  
601 program under project RISC\_PLUS (0031\_RISC\_PLUS\_6\_E).

602 During the preparation of this work the authors used ChatGPT-4 in order to improve language  
603 and readability. After using this tool/service, the authors reviewed and edited the content as  
604 needed and take full responsibility for the content of the publication.

605

606 **References**

- 607 AEMET-IM Iberian Climate Atlas. Agencia Estatal de Meteorología, Instituto de  
608 Meteorología de Portugal, 2011.  
609 [https://www.aemet.es/documentos/es/conocerlas/recursos\\_en\\_linea/publicaciones\\_y](https://www.aemet.es/documentos/es/conocerlas/recursos_en_linea/publicaciones_y_estudios/publicaciones/Atlas-climatologico/Atlas.pdf)  
610 [estudios/publicaciones/Atlas-climatologico/Atlas.pdf](https://www.aemet.es/documentos/es/conocerlas/recursos_en_linea/publicaciones_y_estudios/publicaciones/Atlas-climatologico/Atlas.pdf).
- 611 Alcoforado, M.J. et al. Early Portuguese meteorological measurements (18th century).  
612 *Climate of the Past* 8, 353–371, doi: 10.5194/cp-8-353-2012, 2012
- 613 Alexander, L.V., and Jones, P.D. Updated precipitation series for the U.K. and discussion of  
614 recent extremes, *Atmospheric Science Letters* 1(2), 142–150,  
615 doi:10.1006/asle.2001.0025, 2001.
- 616 Amorín, I.. As cheias do rio Douro no Porto (Portugal) do Século XVIII in *Sémata: Ciências*  
617 *Sociais e Humanidades*, 2017.
- 618 Ávila y LaCueva, F. *Historia Civil y Eclesiástica de la Ciudad de Tuy y su Obispado*, 1852.
- 619 Barker, T. A Letter from Thomas Barker, Esq; Of Lyndon in Rutlandshire, to James West,  
620 Esq; Pres.R. S. concerning Observations of the Quantities of Rain Fallen at That Place  
621 for Several Years. *Philosophical Transactions (1683-1775)*, 61, 221–226, 1771.
- 622 Béaur, G., and Chevet, J.M. Chapter 4 France in Famines in European History. Editors  
623 Alfani, G. and Ó Gráda, C. Cambridge University Press, ISBN: 9781316841235, doi:  
624 <https://doi.org/10.1017/9781316841235>, 2017.
- 625 Beneyto, C., Aranda, J.A., Benito, G., and Francés, F. New Approach to Estimate Extreme  
626 Flooding Using Continuous Synthetic Simulation Supported by Regional Precipitation  
627 and Non-Systematic Flood Data. *Water* 12, 3174, doi:10.3390/w12113174, 2020.
- 628 Benito, G., Castillo, O., Ballesteros- Cánovas, J.A., Machado, M., and Barriendos, M.  
629 Enhanced flood hazard assessment beyond decadal climate cycles based on centennial  
630 historical data (Duero basin, Spain). *Hydrol. Earth Syst. Sci.*, 25, 6107–6132  
631 <https://doi.org/10.5194/hess-25-6107-2021>, 2021.
- 632 Brázdil, R., Trnka, M., Mikšovský, J., Řezníčková, L., and Dobrovolný, P. Spring-summer  
633 droughts in the Czech Land in 1805–2012 and their forcings. *International Journal of*  
634 *Climatology*, 35, 1405–1421, 2015.
- 635 Bravo-Paredes, N., Gallego, M.C., Domínguez-Castro, F., García, J.A., and Vaquero, J.M.  
636 Pro-pluvia rogation ceremonies in Extremadura (Spain): Are they a good proxy of winter  
637 NAO? *Atmosphere*, 11(3), 282, 2020.
- 638 Brönnimann, S. *Climatic Changes Since 1700*. Springer, doi: 10.1007/978-3-319-19042-6,  
639 2015.
- 640 Ceballos y Fernández de Córdoba, L. *El maíz y la dura, informe y noticias históricas sobre*  
641 *la introducción del cultivo del maíz en Galicia*. Editorial Maestre, 1953.

- 642 Clark, C. The Great Flood of 1768 at Bruton, Somerset. *Weather*, 54, 108-113,  
643 <https://doi.org/10.1002/j.1477-8696.1999.tb06436.x>, 1999.
- 644 Collet, D., and Schuh, M., eds. *Famines During the ‘Little Ice Age’ (1300–1800): Socio-*  
645 *natural Entanglements in Premodern Societies*, Springer, Berlin/Heidelberg,  
646 <https://doi.org/10.1007/978-3-319-54337-6>, 2018.
- 647 Degroot, D., Anchukaitis, K., and Bauch, M. et al. Towards a rigorous understanding of  
648 societal responses to climate change. *Nature* 591, 539–550, doi: 10.1038/s41586-021-  
649 03190-2, 2021.
- 650 deCastro, M., Lorenzo, M.N, Taboada, J.J., Sarmiento, M., Alvarez, I., and Gomez-Gesteira,  
651 M. Influence of teleconnection patterns on precipitation variability and on river flow  
652 regimes in the Mino River basin (NW Iberian Peninsula). *Climate Res.* 32, 63–73, doi:  
653 10.3354/cr032063, 2006.
- 654 de Sallier Dupin, G. *Bulletin 28 mémoires 2000, les amis de Lamballe et Penthièvre*, 2000.
- 655 Diodato, N., Ljungqvist, F.C., and Bellocchi, G., A millennium-long reconstruction of  
656 damaging hydrological events across Italy. *Scientific Reports*, 9, 9963.  
657 <https://doi.org/10.1038/s41598-019-46207-72>, 2019.
- 658 Diodato, N., Ljungqvist, F.C., and Bellocchi, G. Fingerprint of climate change in  
659 precipitation aggressiveness across the central Mediterranean area. *Scientific Reports*,  
660 10, 22062. <https://doi.org/10.1038/s41598-020-7885>, 2020.
- 661 Domínguez-Castro, F., Santisteban, J.I., Barriendos, M., and Mediavilla, R. Reconstruction  
662 of drought episodes for central Spain from rogation ceremonies recorded at the Toledo  
663 Cathedral from 1506 to 1900: A methodological approach. *Global Planet. Change*, 63,  
664 230–242, <https://doi.org/10.1016/j.gloplacha.2008.06.002>, 2008.
- 665 Domínguez-Castro, F., *et al.* Assessing extreme droughts in Spain during 1750-1850 from  
666 rogation ceremonies. *Clim. Past*, 8, 705–722, <https://doi.org/10.5194/cp-8-705-2012>,  
667 2012.
- 668 Domínguez -Castro, F. *et al.* Early Spanish meteorological records (1780–1850). *Int. J.*  
669 *Climatol.* 34, 593–603, doi: 10.1002/joc.3709, 2014.
- 670 Domínguez-Castro, F., Ramos, A.M., García-Herrera, R., and Trigo, R.M. Iberian extreme  
671 precipitation 1855/1856: an analysis from early instrumental observations and  
672 documentary sources. *Int. J. Climatol.* 35, 142–153, <https://doi.org/10.1002/joc.3973>,  
673 2015.
- 674 Domínguez-Castro, F., *et al.* Dating historical droughts from religious ceremonies, the  
675 international pro pluvia rogation database. *Scientific Data*, 8(1), 186, doi:  
676 10.1038/s41597-021-00952-5, 2021.
- 677 Domínguez Ortiz, A. *Carlos III y la España de la Ilustración*. Madrid: Alianza Editorial,  
678 ISBN: 9788491044468, 2005.

- 679 Eiras, R.A. Estudios sobre agricultura y población en la España Moderna. Santiago, 1990.
- 680 Fagan, B. The Little Ice Age. How Climate Made History, 1300–1850. Nueva York: Basic  
681 Books, ISBN 0465022723, 9780465022724, 2001.
- 682 Fernandez-Cortizo, C. ¿En Galicia, el hambre entra nadando? Rogativas, clima y crisis de  
683 subsistencias en la Galicia Litoral suboccidental en los siglos XVI-XVIII. SÉMATA,  
684 Ciencias Sociais e Humanidades, 17, 259-298, <http://hdl.handle.net/10347/4459>, 2005.
- 685 Fernandez-Novoa, D., Gonzalez-Cao, J., Figueira, J.R., Catita, C., García-Feal, O., Gomez-  
686 Gesteira, M., and Trigo, R.M. Numerical simulation of the deadliest flood event of  
687 Portugal: Unravelling the causes of the disaster. Sci. Total Environ. 896, 165092,  
688 <https://doi.org/10.1016/j.scitotenv.2023.165092>, 2023.
- 689 Fernández-Nóvoa, D., Ramos, A.M., González-Cao, J., García-Feal, O., Catita, C., Gómez-  
690 Gesteira, M., and Trigo, R.M.. How to mitigate flood events similar to the 1979  
691 catastrophic floods in the lower Tagus. Nat. Hazard Earth Sys , 24(2), 609–630,  
692 <https://doi.org/10.5194/nhess-24-609-2024>, 2024.
- 693 Font Tullot, I. Historia del Clima en España: Cambios climáticos y sus causas. Instituto  
694 Nacional de Meteorología. Madrid, ISBN 8450571782, 9788450571783, 1988.
- 695 Gonzalez-Cao, J., Fernandez-Novoa, D., García-Feal, O., Figueira, J.R., Vaquero, J.M,  
696 Trigo, R.M., and Gomez-Gesteira, M. Numerical reconstruction of historical extreme  
697 floods: The Guadiana event of 1876. J Hydrol. 599, 126292,  
698 <https://doi.org/10.1016/j.jhydrol.2021.126292>, 2021.
- 699 González-Fernández, M.A. Las crisis de subsistencias y epidémicas en las villas de Vigo y  
700 Bouzas (1620-1890), Boletín del Instituto de Estudios Vigueses, 6, 87–105, ISSN 1135-  
701 1810, 2000.
- 702 Hoyle, R. Chapter 7 Britain in Famines in European History. Editors Alfani, G. and Ó  
703 Gráda, C. Cambridge University Press, ISBN: 9781316841235, doi:  
704 <https://doi.org/10.1017/9781316841235>, 2017.
- 705 Jenkinson, A. F., Shackleton, W., Lawson, S. Monthly and annual rainfall for Ireland, 1711–  
706 1977. UK Met Office Branch Memorandum 77, Meteorological Office, Bracknell, 1979.
- 707 Köppen, W., Die Wärmezonen der Erde, nach der Dauer der heissen, gemässigten und kalten  
708 Zeit und nach der Wirkung der Wärme auf die organische Welt betrachtet (The thermal  
709 zones of the earth according to the duration of hot, moderate and cold periods and to the  
710 impact of heat on the organic world). – Meteorol. Z. 1, 215–226. (translated and edited  
711 by VOLKEN E. and S. BRÖNNIMANN. – Meteorol. Z. 20 (2011), 351–360).  
712 doi:10.1127/0941-2948/2011/105, 1884.
- 713 Labrada, J.L. Descripción económica del Reyno de Galicia, Ed. Galaxia, 1804.
- 714 Lamb, H.H. Climate, History and the Modern World, Second edition, Routledge. London  
715 and New York, ISBN:9780415127349, 0415127343, 1995.

- 716 Le Roy Ladure, E. Sur les fluctuations du climat de la France septentrionale et centrale  
717 depuis le XVIIe siècle. Academie des Sciences Morales et Politiques, ISBN-13  
718 978-2213654249, 2011.
- 719 Ljungqvist, F. C., Seim, A., and Huhtamaa, H. Climate and society in European history,  
720 Wiley Interdisciplin. Rev. Clim. Change, 12, e691, <https://doi.org/10.1002/wcc.691>,  
721 2020.
- 722 Ljungqvist, F. C., Seim, A., and Collet, D.: Famines in medieval and early modern Europe  
723 – connecting climate and society, Wiley Interdisciplin. Rev. Clim. Change, 15, e859,  
724 <https://doi.org/doi.org/10.1002/wcc.859>, 2023a.
- 725 Ljungqvist, F. C., Christiansen, B., Esper, J., Huhtamaa, H., Leijonhufvud, L., Pfister, C.,  
726 Seim, A., Skoglund, M. K., and Thejll, P. Climatic signatures in early modern European  
727 grain harvest yields, Clim. Past, 19, 2463–2491, [https://doi.org/10.5194/cp-19-2463-](https://doi.org/10.5194/cp-19-2463-2023)  
728 [2023](https://doi.org/10.5194/cp-19-2463-2023), 2023b.
- 729 López Linage, J. La patata en España. Historia y agroecología del tubérculo andino. Ed.  
730 Ministerio de Medio Ambiente y Medio Rural y Marino, ISBN: 84-491-0855-1, 2008.
- 731 Losada-Sanmartín, M.L. Chapter 6. Documentación histórica e clima en Historia da  
732 meteoroloxía e da climatoloxía de Galicia. Ed. Francisco Díaz-Fierros Viqueira, ISBN  
733 978-84-96530-72-0, 2008.
- 734 Macdonald, N., and Sangster, H. High-magnitude flooding across Britain since AD 1750.  
735 Hydrol. Earth Syst. Sci., 21, 1631–1650, 2, <https://doi.org/10.5194/hess-21-1631-2017>,  
736 2017.
- 737 Martín-García, A. Ordinary and Extraordinary Death Rate in El Ferrol at the End of the  
738 Ancien Regime. Stud His Ha Mod. 23, 249–273, 2001.
- 739 Martínez-Rodríguez, E. Santiago council and the crises of subsistences between 1741 and  
740 1770. SÉMATA, Ciencias Sociais e Humanidades 29, 219–237, 2017.
- 741 Meijide-Pardo, A. El hambre de 1768-1769 en Galicia y la obra asistencial del estamento  
742 eclesiástico compostelano. Compostellanum: revista de la Archidiócesis de Santiago de  
743 Compostela, 10, 213–256, ISSN 0573-2018, (1965).
- 744 Moravec, V., Markonis, Y., Rakovec, O., Kumar, R. and Hanel, M. A 250-year European  
745 drought inventory derived from ensemble hydrologic modeling. Geo Res Letters,46,  
746 5909–5917, <https://doi.org/10.1029/2019GL082783>, 2019.
- 747 Murphy, C., *et al.*, A 305-year continuous monthly rainfall series for the Island of Ireland  
748 (1711-2016). Clim Past, 14, 413–440, <https://doi.org/10.5194/cp-14-413-2018>, 2018.
- 749 Murphy, C., *et al.*, The forgotten drought of 1765–1768: Reconstructing and re-evaluating  
750 historical droughts in the British and Irish Isles. Int J Climatol. 40, 5329–5351,  
751 <https://doi.org/10.1002/joc.6521>, 2020.

- 752 Nassiet, M. La diffusion du blé noir en France à l'époque moderne. *Histoire et Sociétés*  
753 *Rurales* 9, 59 – 77, 1998.
- 754 Nicault, A., Alleaume, S., Brewer, S., Carrer, M., Nola, P. and Guiot, J. Mediterranean  
755 drought fluctuation during the last 500 years based on tree-ring data. *Clim Dynam*, 31(2–  
756 3), 227–245. <https://doi.org/10.1007/s00382-007-0349-3>, 2008.
- 757 Noone, S., Murphy, C., Coll, J., Matthews, T., Mullan, D., Wilby, R.L. and Walsh, S.  
758 Homogenization and analysis of an expanded long-term monthly rainfall network for the  
759 Island of Ireland (1850–2010). *Int. J. Climatol.*, 36(8), 2837–2853,  
760 <https://doi.org/10.1002/joc.4522>, 2016.
- 761 Noone, S., Broderick, C., Duffy, C., Matthews, T., Wilby, R.L. and Murphy, C. A 250-year  
762 drought catalogue for the Island of Ireland (1765–2015). *Int. J. Climatol.*, 37, 239–254.  
763 <https://doi.org/10.1002/joc.4999>, 2017.
- 764 Ó Gráda, C. Chapter 8, Ireland in Famines in European History. Ed. Alfani, G. and Ó Gráda,  
765 C. Cambridge University Press, ISBN: 9781316841235, doi:  
766 <https://doi.org/10.1017/9781316841235>, 2017.
- 767 Peña, J.C., *et al.*, Low-Frequency Atmospheric Variability Patterns and Synoptic Types  
768 Linked to Large Floods in the Lower Ebro River Basin. *J Climate* 35, 2351–2371, doi:  
769 10.1175/JCLI-D-20-0394.1, 2022.
- 770 Pérez-Costanti, P. *Notas Viejas Galicianas*. Vigo, 1925.
- 771 Pérez-Moreda, V. (2017). Chapter 3, Spain in Famines in European History. Editors Alfani,  
772 G. and Ó Gráda, C. Cambridge University Press, ISBN: 9781316841235, DOI:  
773 <https://doi.org/10.1017/9781316841235>.
- 774 Pfister, C. *Das Klima der Schweiz von 1525-1860 und seine Bedeutung in der Geschichte*  
775 *von Bevölkerung und Landwirtschaft*. Bern: Paul Haupt, 1984.
- 776 Pfister, C. Monthly temperature and precipitation patterns in Central Europe 1525-1979:  
777 quantifying documentary evidence on weather and its effects. In Bradley, R.S.; Jones,  
778 P.D. (ed.). *Climate since A.D. 1500*. London/New York: Routledge, 118-142, 1992.
- 779 Potop, V., Boroneant, C., Možný, M., Štěpánek, P. and Skalák, P. Observed spatio temporal  
780 characteristics of drought on various time scales over The Czech Republic. *Theor and*  
781 *Appl Climatol*, 115, 563–581, doi 10.1007/s00704-013-0908-y, 2014.
- 782 Simpson, I.R. and Jones, P.D. Analysis of UK precipitation extremes derived from met  
783 office gridded data. *Int. J. Climatol.*, 34, 2438–2449, <https://doi.org/10.1002/joc.3850>,  
784 2014.
- 785 Silva, L.P. *O clima do Noroeste de Portugal (1600-1855): dos discursos aos impactos*.  
786 Thesis, 2019.

- 787 Slavin, P. Climate and famines: A historical reassessment, Wiley Interdisciplin. Rev. Clim.  
788 Change, 7, 433–447, <https://doi.org/10.1002/wcc.395>, 2016.
- 789 Spraggs, G., Peaver, L., Jones, P. and Ede, P. Re-construction of historic drought in the  
790 Anglian region (UK) over the period 1798–2010 and the implications for water  
791 resources and drought management. J. Hydrol., 526, 231–252,  
792 <https://doi.org/10.1016/j.jhydrol.2015.01.015>, 2015.
- 793 Todd, B., Macdonald, N., Chiverrell, R.C., Caminade, C. and Hooke, J.M. Severity, duration  
794 and frequency of drought in SE England from 1697 to 2011. Climatic Change, 121, 673–  
795 687, doi 10.1007/s10584-013-0970-62013.
- 796 Trigo, R.M., Pozo-Vázquez, D., Osborn, T.J., Castro-Díez, Y., Gámiz-Fortis, S., and  
797 Esteban-Parra, M.J. North Atlantic Oscillation influence on precipitation, river flow and  
798 water resources in the Iberian Peninsula. Int. J. Climatol. 24, 925–944,  
799 <https://doi.org/10.1002/joc.1048>, 2004.
- 800 Vicente-Serrano, S., *et al.*, Long-term variability and trends in meteorological droughts in  
801 Western Europe (1851–2018). Int J Climatol 41 E690–E717, doi: 10.1002/joc.6719,  
802 2020.
- 803 Valler, V., Franke, J., Brugnara, Y., and Brönnimann, S. An updated global atmospheric  
804 paleo- reanalysis covering the last 400 years. Geosci Data J. 9: 89–107,  
805 <https://doi.org/10.1002/gdj3.121>, 2022.
- 806 White, S., Brooke, J., and Pfister, C. Climate, Weather, agriculture, and food, in: The  
807 Palgrave Handbook of Climate History, edited by: White, S., Pfister, C., and  
808 Mauelshagen, F., 331–353, Springer, Berlin/Heidelberg, ISBN 978-1-137-43019-9,  
809 <https://doi.org/10.1057/978-1-137-43020-5> , 2018a.
- 810 White, S., Pfister, C., and Mauelshagen, F., eds. The Palgrave Handbook of Climate History.  
811 Palgrave Macmillan London. <https://doi.org/10.1057/978-1-137-43020-5>, 2018b.
- 812