

# Holocene land cover change in North America: continental trends, regional drivers, and implications for vegetation-atmosphere feedbacks

Andria Dawson\*<sup>†1</sup>, John W. Williams<sup>†2</sup>, Marie-José Gaillard<sup>3</sup>, Simon J Goring<sup>2</sup>, Behnaz Pirzamanbein<sup>4</sup>, Johan Lindstrom<sup>5</sup>, R. Scott Anderson<sup>6</sup>, Andrea Brunelle<sup>7</sup>, David Foster<sup>8</sup>, Konrad Gajewski<sup>9</sup>, Daniel G. Gavin<sup>10</sup>, Terri Lacourse<sup>11</sup>, Thomas A Minckley<sup>12</sup>, Wyatt Oswald<sup>13</sup>, Bryan Shuman<sup>12</sup>, Cathy Whitlock<sup>14</sup>

<sup>1</sup>Department of Mathematics and Computing, Department of Biology, Mount Royal University, Calgary, AB T3E6K6

<sup>2</sup>Department of Geography and Center for Climatic Research, University of Wisconsin-Madison, Madison, WI 53706

<sup>3</sup>Department of Biology and Environmental Science, Linnaeus University, SE 39231 Kalmar, Sweden

<sup>4</sup>Department of Statistics, School of Economics and Management, Lund University, SE-220 07 Lund, Sweden

<sup>5</sup>Division of Mathematical Statistics, Centre for Mathematical Sciences, Lund University, SE-221 00 Lund, Sweden

<sup>6</sup>School of Earth and Sustainability, Northern Arizona University, Flagstaff, AZ 86011

<sup>7</sup>Geography Department, University of Utah, Salt Lake City, UT 84112

<sup>8</sup>Harvard Forest, Harvard University, Petersham, MA 01366

<sup>9</sup>Département de Géographie, Environnement et Géomatique, Université d'Ottawa, Ottawa, ON K1N 6N50

<sup>10</sup>Department of Geography, University of Oregon, Eugene, OR 97403

<sup>11</sup>Department of Biology and Centre for Forest Biology, University of Victoria, Victoria, BC V8P 5C2

<sup>12</sup>Department of Geology and Geophysics, University of Wyoming, Laramie, WY 82071

<sup>13</sup>Marlboro Institute for Liberal Arts and Interdisciplinary Studies, Emerson College, Boston, MA 02116 and

<sup>14</sup>Department of Earth Sciences, Montana State University, Bozeman, MT 59717

\*Correspondence email: [adawson@mtroyal.ca](mailto:adawson@mtroyal.ca)

<sup>†</sup>These authors contributed equally to the development of this manuscript

## 30 Abstract

31 Land cover governs the biogeophysical and biogeochemical feedbacks between the land surface and  
32 atmosphere. Holocene vegetation-atmosphere interactions are of particular interest, both to understand the  
33 climate effects of intensifying human land use and as a possible explanation for the Holocene  
34 [Temperature Conundrum](#), a widely studied mismatch between simulated and reconstructed  
35 temperatures. Progress has been limited by a lack of data-constrained, quantified, and consistently  
36 produced reconstructions of Holocene land cover change. As a contribution to the Past Global Changes  
37 (PAGES) LandCover6k Working Group, we present a new suite of land cover reconstructions with  
38 uncertainty for North America, based on a network of 1445 sedimentary pollen records and the  
39 REVEALS pollen-vegetation model coupled with a Bayesian spatial model. These spatially  
40 comprehensive land cover maps are then used to determine the pattern and magnitude of North American  
41 land cover changes at continental to regional scales. Early Holocene afforestation in North America was  
42 driven by rising temperatures and deglaciation, and this afforestation likely amplified early Holocene  
43 warming via the albedo effect. A continental-scale mid-Holocene peak in summergreen trees and shrubs  
44 (8.5 to 4 ka) is hypothesized to represent a positive and understudied feedback loop among insolation,  
45 temperature, and phenology-[seasonality](#). A last-millennium decrease in summergreen trees and shrubs  
46 with corresponding increases in open land likely was driven by a spatially varying combination of  
47 intensifying land use and neoglaciation. Land cover trends vary within and across regions, due to  
48 individualistic taxon-level responses to environmental change. Major species-level events, such as the  
49 mid-Holocene decline of [Tsuga canadensis](#) (eastern hemlock), may have altered regional climates. The  
50 substantial land-cover changes reconstructed here support the importance of biogeophysical [and](#)  
51 [biogeochemical](#) vegetation feedbacks to Holocene climate-[carbon](#) dynamics. However, recent model  
52 experiments that invoke vegetation feedbacks to explain the Holocene [Temperature Conundrum](#) may  
53 have overestimated ~~the~~ land cover forcing by replacing Northern Hemisphere grasslands >30°N with  
54 forests; an ecosystem state that is not supported by these land cover reconstructions. These Holocene  
55 reconstructions for North America, along with similar LandCover6k products now available for other  
56 continents, serve the Earth system modeling community by providing better-constrained land cover  
57 scenarios and benchmarks for model evaluation, ultimately making it possible to better understand the  
58 regional- to global-scale processes driving Holocene land-~~cover~~, [carbon-cycle](#), [and climate](#) dynamics.  
59

## 1. Introduction

Vegetation is the great mediator of biogeophysical and biogeochemical interactions between the land surface and the atmosphere (Bonan and Doney, 2018; Gaillard et al., 2010; Harrison et al., 2020; Pongratz et al., 2010). Enhanced carbon uptake and sequestration by terrestrial ecosystems is an essential component to contemporary ~~net-negative-net~~ CO<sub>2</sub> emission scenarios needed to stabilize the climate system and mitigate the dangerous impacts already emerging (Rogelj et al., 2018; van Vuuren et al., 2017). During the Holocene, as cryosphere-ocean-atmosphere feedbacks waned and anthropogenic land use intensified (Ruddiman, 2013; Stephens et al., 2019), vegetation-atmosphere feedbacks and forcings increased in importance, particularly in regions where climate variability interacted with major changes in vegetation structure. Examples include soil and vegetation feedbacks that amplified precessional-driven variations in monsoonal rainfall intensity in North Africa and Asia (Chen et al., 2021; Chandan and Peltier, 2020), and increases in high-latitude tree cover, which decreased wintertime albedo and increased temperatures. [Carbon sequestration due to Northern Hemisphere postglacial afforestation is estimated at 13.9 GtC between 9 and 6 ka and a 3.5 GtC loss after 3 ka.](#) Intensified human land use and resulting greenhouse gas emissions may have delayed Northern Hemisphere late-Holocene cooling and glaciation (Ruddiman, 2003). However, initial models of global anthropogenic land cover change (ALCC) (Kaplan et al., 2009; Klein Goldewijk et al., 2011, 2010) over the Holocene were largely unconstrained by paleoecological and archaeological observations and so differed widely in their estimated size and scope of the anthropogenic footprint. More recently, Holocene increases in vegetation cover have been invoked to explain the Holocene ~~Temperature Conundrum~~, a discrepancy between proxy and model-estimated temperature during the early- to mid-Holocene (Thompson et al., 2022; Kaufman and Broadman, 2023), ~~but global simulations of vegetation-climate feedbacks during the Holocene are not constrained by observational data.~~ At subcontinental scales, data-constrained studies of Holocene climate-vegetation feedbacks in Europe indicate that mid-Holocene vegetation changes relative to pre-Industrial baselines could have warmed winters in some areas by 4–6°C in northeastern Europe (Strandberg et al., 2022, 2014). [Global simulations of vegetation-climate feedbacks during the Holocene, however, remain unconstrained by observational data.](#)

Hence, ~~there is an on-going need for~~ comprehensive and accurate proxy-based reconstructions of past land cover at regional to global extents [are needed](#) (Gaillard and LandCover6k Steering Group, 2015; Gaillard et al., 2018, 2010). These reconstructions can ~~then~~ be used with Earth system models (ESMs) to test hypotheses about the physical, biological, and anthropogenic processes that drove Holocene climate variability (Harrison et al., 2020). Fossil pollen records offer the primary observational constraint on past vegetation composition and structure, with thousands of records now available globally. Efforts to

systematically map late-Quaternary land cover using fossil pollen data and well-defined rulesets began in the late 1990s with the Biome6000 project (Prentice et al., 2000, 2011). Since then, the continental-scale pollen databases launched in the 1980s and 1990s (Grimm et al., 2013) have coalesced along with other paleoecological data into the Neotoma Paleoecology Database (Neotoma), an international, multi-proxy, curated data resource that helps tame issues of data heterogeneity through community curation by experts (Williams et al., 2018). Neotoma thus enables global-scale analyses of past vegetation and climate change (Mottl et al., 2021; Herzschuh et al., 2023).

Multiple pollen-vegetation models (PVMs) have been developed to make quantitative inferences about past vegetation. Some PVMs involve relatively simple but effective transfer functions, such as the modern analogue technique (Williams et al., 2011b) or rule-based systems for classifying land cover (Prentice et al., 2000; Cruz-Silva et al., 2022; Fyfe et al., 2010). Others are process-based proxy system models (Evans et al., 2013) that attempt to represent the processes governing the atmospheric transport and deposition of pollen, such as the REVEALS and LOVE (Sugita, 2007a, b) or STEPPS (Dawson et al., 2019). Efforts continue to test and refine the parameterizations of these models through paired analyses of pollen assemblages and forest composition at local to landscape scales (Liu et al., 2022).

In response to these scientific needs and opportunities, the Past Global Changes (PAGES) LandCover6k working group was launched as an international effort (Gaillard and LandCover6k Steering Group, 2015) to reconstruct vegetation globally for the Holocene. LandCover6k, led by experts typically working at continental scales, has had the explicit aim of creating vegetation reconstructions that can better constrain past [land-use](#) histories ~~of anthropogenic land use~~ in ESMs and is mostly based on networks of fossil pollen records. To facilitate the use of these vegetation reconstructions in ESMs, the REVEALS PVM has been used for all LandCover6k reconstructions, with standard model parameterizations and standard protocols for pollen data handling. LandCover6k gridded REVEALS reconstructions at the continental scale have been published so far for Europe (Githumbi et al., 2022a, b; Serge et al., 2023; Trondman et al., 2015) and China (Li et al., 2023). However, no comparable REVEALS-based land cover reconstructions are available for North America, despite a comparable density of fossil pollen records to these other regions (Stegner and Spanbauer, 2023) and prior regional-scale applications of REVEALS in North America (Sugita et al., 2010; Chaput and Gajewski, 2018).

REVEALS uses pollen counts, pollen productivity estimates, pollen fall speeds, atmospheric conditions, and sedimentary basin type and size to estimate vegetation composition for a given time period. REVEALS accounts for the processes of differential pollen production (determined by pollen productivity estimates) and dispersal-deposition (determined by the pollen fall speeds, atmospheric conditions, and sedimentary basin type and size).

While REVEALS reconstructions usually combine information from multiple pollen records, ~~the~~  
~~mode~~REVEALS is not explicitly spatial, and so does not support the interpolation of inferences to places  
with no pollen records. To address this issue, ~~other~~ researchers have developed a statistical approach to  
spatially interpolate REVEALS-based land cover estimates from individual grid cells to all cells within  
the grid, including estimates of uncertainty (Pirzamanbein et al., 2014, 2018). The approach uses a  
Bayesian hierarchical model with spatial dependence specified according to a Gaussian Markov Random  
Field (GMRF; Lindgren et al. (2011)); we refer to the two-step process of REVEALS-based estimation  
followed by this spatial interpolation as the REVEALS-GMRF approach. REVEALS-GMRF has been  
used to develop spatially continuous gridded vegetation reconstructions in Europe to assess vegetation-  
climate feedbacks resulting from natural and anthropogenic land cover change (Strandberg et al., 2022)  
and to evaluate ALCC models (Kaplan et al., 2017) and dynamic vegetation models (Pirzamanbein et al.,  
2020; Dallmeyer et al., 2023; Zapolska et al., 2023).

Here we adopt the REVEALS-GMRF approach to reconstruct land cover changes in North  
America during the Holocene. This work relies on 1445 Holocene pollen records drawn from Neotoma  
and its constituent database, the North American Pollen Database (NAPD), with a targeted data-  
mobilization campaign employed to add more records to the NAPD for western North America. Using  
these data, we reconstruct the fractional cover of 32 plant taxa and three land cover types (LCTs):  
evergreen trees and shrubs (ETS), summergreen trees and shrubs (STS), and open vegetation/land (OVL,  
including grasses, herbs, and low shrubs) from 12,000 years ago (12 ka) to present. We present the  
Holocene vegetation reconstructions by working across scales, first describing continental-scale trends in  
land cover, then shifting to several regional-scale case studies to show how the continental-scale trends  
emerge from taxon-level dynamics that vary within and among regions, with respect to key taxa, drivers,  
and resultant land-cover changes. We then zoom out to discuss the continental-scale drivers of Holocene  
land cover change in North America and possible biophysical implications of these changes for Holocene  
vegetation-atmosphere interactions and the Holocene [Temperature c](#)onundrum. Lastly, we discuss the  
potential limitations of the REVEALS-GMRF approach and the opportunities now available for well-  
constrained hemispheric- to global-scale studies of vegetation-atmosphere interactions.

## 2. Data and methods

### 2.1 Pollen data and data mobilization for western North America

Western North America has traditionally been underrepresented in the NAPD and Neotoma, but the  
density of fossil pollen records in western North America has steadily increased in recent decades, as  
multiple teams have worked to collect new records, often focusing on interactions among past vegetation,

fire, climate, and human dynamics (Anderson et al., 2008; Gavin and Brubaker, 2014; Marlon et al., 2012; Iglesias et al., 2018; Alt et al., 2018). Many of these datasets were contributed to Neotoma (Williams et al., 2018) when originally published, while others were contributed to Neotoma for an open-data mobilization campaign conducted for this paper and PAGES LandCover6k (Gaillard et al., 2018).

After this effort, 1582 North American Holocene pollen records were downloaded from Neotoma (Supp. Table 1). Each record included pollen count data at a series of depths. For methodological consistency (Flantua et al., 2023), we used the Bayesian age-depth model Bchron to refit all age-depth models using a custom-built workflow (<https://github.com/andydawson/bulk-bchron>). This workflow that assessed chronological constraints and ~~then used the~~ mse chronological constraints and IntCal20NH (Reimer et al., 2020) to fit the Bchron age-depth model (Parnell et al., 2008). This resulted in 1445 records with age-depth models. Ages of the youngest and oldest chronological constraints were used to determine the reliable age range for each record; we limited extrapolation of pollen sample ages beyond the youngest or oldest constraints to 1000 years. For discussion, evaluation, and comparison of widely used age-depth models, we refer readers to Lacourse and Gajewski (2020), McKay et al. (2021), and Trachsel and Telford (2017).

Finally, pPollen types were aggregated to taxa using the taxa list in the North American Modern Pollen Database (Whitmore et al., 2005). ~~In this work, we first translated the Neotoma taxonomy to the Whitmore taxonomy (ref);~~ This resulted in a list of 47 taxa. We identified the taxa that were most abundant and ~~also~~ indicators of land cover type, of which there were 33 (Supp. Table S2). There were corresponding pollen productivity ~~PPE~~ and fallspeed values for all of these taxa. Among the ~~The~~ set of excluded taxa (Nyssa, Platanus, Apiaceae, Liquidambar, Ephedra, Cornus, Chrysopsis, Eriogonum, Cactaceae, Taxus, Larrea), the fraction of taxon pollen grains with respect to total pollen grain count was at most 0.16% (Nyssa), while the total fraction of excluded pollen taxa was about 0.5% ~~represents about results in a total of about 0.5% of the total pollen grains counted (for the North American Holocene); with the most abundant excluded.~~ We identified ~~ed~~ used a subset of 332 taxa for this analysis, choosing the most abundant taxa, several open land indicators, and taxa with available estimates of relative pollen productivity (Tables 2, 5). There were several taxa that were excluded because there were no published pollen productivity estimates (Nyssa, Platanus, Apiaceae, Liquidambar, Ephedra, Cornus, Chrysopsis, Eriogonum, Cactaceae, Taxus, Larrea). However, the fraction of taxon pollen grains with respect to total pollen grain count was at most 0.16% (Nyssa), while the total fraction of excluded pollen taxa was about 0.5% for even the most abundant excluded taxon, Nyssa, the represented only 0.16% of total pollen

## 2.2 REVEALS (regional reconstructions)

191 Pollen-based land cover reconstructions were performed using the REVEALS pollen-vegetation model  
 192 (Sugita, 2007a), and based on the standard protocol for PAGES LandCover6k (Trondman et al., 2015;  
 193 Githumbi et al., 2022a). This model estimates the relative abundance of plant taxa, along with the  
 194 standard error of these estimates, given pollen counts and input parameters that represent sedimentary  
 195 basin size and type, pollen productivity estimates (PPEs), pollen fall speeds, and atmospheric conditions.  
 196 REVEALS was developed to operate at the regional scale (Hellman et al., 2008a, b; Sugita, 2007a);  
 197 inferences of plant relative abundance represent the background vegetation over large areas (suggested as  
 198 100 km x 100 km by Hellman et al. (2008b), but this scale is variable). REVEALS traditionally has been  
 199 used to infer plant relative abundance from records from large lakes (>50 ha), but has been tested and  
 200 applied to regions with records from a number of smaller lakes. The REVEALS model accounts for both  
 201 differential productivity and dispersal among taxa. Differential productivity is determined by taxon-  
 202 specific PPEs, while dispersal is modeled according to a Gaussian plume (Sutton, 1953) or Lagrangian  
 203 dispersal-deposition model (Kuparinen et al., 2007), both of which require the specification of  
 204 atmospheric conditions including wind speed. REVEALS accounts for differential dispersibility among  
 205 taxa using pollen fall speeds. See Githumbi et al. (2022a), and Sugita (2007a) for a more detailed and  
 206 theoretical description of REVEALS. We implement REVEALS using the REVEALSinR R package  
 207 (Theuerkauf et al., 2016). REVEALS estimates for other regions included in the LandCover6k effort  
 208 (Githumbi et al., 2022a; Li et al., 2023) were developed using more traditional implementations of this  
 209 model (LRA.REVEALS.v6.2.4.exe (Sugita, unpublished) and LRA R package (Abraham et al., 2014)),  
 210 which differ slightly in their calculation of the standard errors of relative abundances.

211 Pollen source areas and the relative representation of plant taxa in the REVEALS dispersal-  
 212 deposition model are affected by sedimentary basin type (e.g. lake, mire) and area (Sugita, 2007a;  
 213 Trondman et al., 2016). Basin type is typically included in Neotoma metadata for pollen datasets, but not  
 214 all datasets include metadata on basin area. To determine basin area for these datasets, we developed a  
 215 standard workflow (Goring, 2024); [https://github.com/NeotomaDB/neotoma\\_lakes](https://github.com/NeotomaDB/neotoma_lakes). First, we used  
 216 hydrological databases (National Hydrography Dataset (United States Geological Survey, 2022), National  
 217 Hydro Network (Natural Resources Canada, 2022)) to assign basin areas to datasets whose coordinates  
 218 fell within a water-body polygon. Second, for dataset coordinates that landed outside a water-body  
 219 polygon, we used Google Earth Engine to identify the basin. These basins were traced using the polygon  
 220 tool in Google Earth Engine, and basin area was calculated from polygon area. Not all basins could be  
 221 identified, however, particularly for pre-GPS sites in Neotoma with imprecise coordinates. In total we  
 222 determined basin area for 545 of the 1445 records. For the remaining ~~Third, for~~ sites still without basin  
 223 area, we assigned a size of 50 ha (900 of 1445 records). In the context of the REVEALS model, this  
 224 represents a medium-sized lake. This decision avoids the potential biases from assigning large or small

lake areas, although site-level reconstructions may over- or under-represent taxa if basin area (and hence pollen source area) is inaccurate (Jackson, 1990; Davis, 2000, p.200; Liu et al., 2022). All basin areas recovered in the first and second steps were added as site-level metadata to Neotoma, along with dataset notes.

We used PPE and fall speed datasets from Wieczorek and Herzschuh (2020) for the Northern Hemisphere extra-tropics. Specifically, we used the North America continental-scale datasets, which include PPE (with grass as the reference taxon) and fall speed values for 30 of the 32 taxa we consider in this work. For *Ambrosia* (ragweed) and *Tsuga* (hemlock), which were not included in Wieczorek and Herzschuh (2020), we use PPE and fall speed values from a previously compiled North America dataset (Dawson et al., 2016; Trachsel et al., 2020). Additionally, for the *Larix* (larch) fall speed, we used the value for *Larix laricina* from Niklas (1984), which is the dominant species in eastern and northern North America. This estimate is an order of magnitude smaller than the *Larix* taxon-level fall-speed estimate included in Wieczorek and Herzschuh (2020), which originates from Bodmer (1922). We experimented with fall-speed datasets that included these larger *Larix* fall-speed estimates; these vegetation reconstructions ~~appeared to indisputably~~ overrepresented *Larix* larch. These challenges associated with dealing with the heavy Larix pollen are described in Sjögren et al. (2008).

We used the Gaussian plume dispersal model with a wind speed of 3 m/s and neutral atmospheric conditions (vertical diffusion coefficient  $c_z=0.12$ ; turbulence parameter  $n=0.25$ ; wind speed  $u=3$  m/s; see Jackson and Lyford (1999)), to be consistent with the dispersal model specified in the LandCover6k protocol for Northern Hemisphere reconstructions (Dawson et al., 2018; Githumbi et al., 2022a). We set the region cutoff to the REVEALSinR function default of 100 km; this specifies the maximum distance that most pollen will originate from (Sugita, 2007a; Theuerkauf et al., 2016).

We reconstructed land cover for 25 time intervals that cover the Holocene. These time intervals were defined according to the LandCover6k working group protocol (Trondman et al., 2015) (SI Table 4). Time intervals are specified in kiloyears before present (ka), where present is defined as 1950 CE, but for time intervals <1ka we also note the Common Era (CE) timescale. Intervals in the period from 11.7 to 0.7 ka have a 500-year temporal grain (11.7 to 11.2 ka; 11.2 to 10.7 ka; etc.), while the three most recent intervals have a finer temporal grain (0.7-0.35 ka [1250-1600 CE]: 350 years; 0.35-0.1 ka [1600-1850 CE]: 250 years; 0.1-(-0.074) ka: 174 years [1850-2024 CE]) in order to better capture the changes associated with intensifying anthropogenic land use over the last five centuries. Pollen samples were assigned to a time interval based on their mean calibrated radiocarbon age. If multiple samples for a record fell within the same interval, pollen counts were summed by taxon, so that each record would have at most one set of pollen counts per time interval. We used a grid resolution of  $1^\circ \times 1^\circ$ , also as specified by the LandCover6k protocol. Within a grid cell, REVEALS reconstructions for multiple sites were



259 averaged, as is standard practice (Sugita, 2007a). Taxon-level reconstructions were aggregated to three  
260 land cover categories (Supp. Table 25): summergreen trees and shrubs (STS), evergreen trees and shrubs  
261 (ETS), and open vegetation/land (OVL), which includes grasses, herbs, and low-stature shrubs. Note that  
262 bare ground cannot be detected by pollen-based land cover mapping and so it is not included as a  
263 potential land cover type; presumably this missing land cover type is usually misclassified as OVL. All  
264 trees and shrubs (ATS) is calculated as the sum of STS and ETS.

265

### 266 2.3 REVEALS-GMRF (spatial modeling and interpolation with uncertainty)

267 ~~Here, We~~ we use the REVEALS-GMRF Bayesian hierarchical model (Pirzamanbein et al., 2018) to  
268 spatially interpolate the REVEALS-based land cover reconstructions. REVEALS-GMRF exploits the  
269 spatial dependence in land cover using a Gaussian Markov Random Field, and permits the  
270 characterization of uncertainty given the empirical land cover product. As in Pirzamanbein et al. (2018),  
271 we use elevation as a covariate. Although including simulations of land cover as an additional covariate  
272 can further improve the reliability of resulting land cover maps (Pirzamanbein et al., 2020), the intent of  
273 our work is to develop a spatial land cover product suitable for validation of and assimilation with models  
274 of dynamic vegetation, land use, and the Earth system models. Hence, to maintain independence, we  
275 refrain from using simulated land cover as a covariate. To quantify overall uncertainty of a grid cell for a  
276 specified time period, we computed the area of confidence regions (CR; Pirzamanbein et al. (2018)).  
277 Smaller CR values indicate higher confidence, while larger CR values indicate more uncertainty. As in  
278 Pirzamanbein et al. (2018), a CR threshold is determined using the complete set of CR values.  
279 Accordingly, any grid cells with a CR greater than a threshold of 9 were omitted from spatial  
280 reconstructions (Githumbi et al., 2022a). The result of this spatial interpolation is a set of empirically-  
281 based land cover maps for North America and the LandCover6k time intervals, with uncertainty.

282

### 283 2.4 Calculation of proportional changes and mapped anomalies

284 To summarize continental-scale land-cover dynamics through time, we built time series of 1) land both  
285 mean relative cover relative to ice-free area for the time interval being mapped (here called relative-to-  
286 time-t mean land cover) and 2) area-weighted relative land cover relative to ice-free land area at 0.25 ka  
287 (here called relative-to-modern mean land cover). First we set a fixed domain size by including only grid  
288 cells for which there were reconstructions for all time periods; this removes confounding effects that may  
289 arise from a changing number of included grid cells. We Then we used recently updated ice sheet maps  
290 (Dalton et al., 2020) to identify glaciated and unglaciated grid cells for each time period, then calculated  
291 the fraction of unglaciated land cover for each time period. We did the same for pro-glacial and other  
292 lakes (Dyke and Prest, 1987; Peltier, 1994). We recognize that shorelines have changed over the

Formatted: Not Highlight

Formatted: Not Highlight

Holocene, but ultimately determined that the effect of this at the considered spatial resolution would be minimal (PALSEA Group, 2010). We then calculated the relative-to-time-t mean land relative cover of each land cover type for each time period as the mean, across all unglaciated grid cells at that time period. This metric of relative-to-time-t ~~Mean relative cover~~ usefully summarizes the relative proportions of land cover types changes within ice-free regions for any given time period, but, because both the numerator and denominator change over time, this metric does not track the overall increases in vegetated land area in North America during the last deglaciation. To calculate relative-to-modern mean area-weighted land relative cover, for each time period, we multiplied the relative cover of each grid cell by the cell area, and then summed the relative cover of each land cover type across space for unglaciated land grid cells. We then divided ~~these summed area-weighted cover~~ this summed relative cover values for each time period by the sum of total unglaciated land area at 0.025 ka. Because this second metric uses a constant denominator that represents late-Holocene deglaciated land area, a number of unglaciated land grid cells increases changes through time, while the denominator is constant, the relative-to-modern this area-weighted metric of relative cover captures both the deglacial is affected by both increase in available land in North America and changes in land cover proportion within that available land, resulting in a proportional metric that is sensitive to continental scale increases in vegetation cover.


For each pair of time intervals, we calculated land cover change as grid-cell scale differences for each of the three land cover classes. After visually identifying areas of large land cover changes (Supp. Fig. S3), we hand-selected identified several regions for further investigation, choosing areas with large changes and/or few prior regional reviews: the northeastern US and southeastern Canada (NEUS/SEC), eastern Canada (ECAN), western Canada and Alaska (WCAN/AK), and the Pacific Coast, Cascades, and Sierra Nevada (PCCS) ~~based on areas of greatest change and pollen site density~~. We then assessed vegetation change for each region at both the taxon and land cover scales, using the REVEALS grid-cell estimates of taxon mean abundances and the REVEALS-GMRF interpolated estimates of mean cover for the land cover types.

### 3. Results


#### 3.1 Data coverage

The assembled dataset of 1445 fossil pollen records has good coverage across the continental US and Canada (Fig. 1a, Supp. Fig. S1). Areas of relatively high site density include the Great Lakes region of the US and Canada, the northeastern US, the Rocky Mountains, the Pacific Coast, and central Alaska. Given good data-mobilization efforts for the US and Canada, this distribution reasonably approximates the true distribution of fossil pollen records, so spatial gaps usually represent lack of available sites (e.g.


325 few lakes or wetlands in arid regions) or inaccessibility (e.g. high Arctic). Conversely, a lower site  
326 density in Mexico and Central America is partially due to less extensive open-data mobilization efforts in  
327 these regions. Temporally, the distribution of oldest samples (an indicator of record length) is smooth  
328 (Fig. 1b), with rapid accumulation of sites between 11.7 and 11 ka (33% of sites have an oldest sample in  
329 this interval) and between 0.5 and -0.074 ka (74% of sites have a youngest sample in this interval).  
330



331




**Figure 1:** The spatiotemporal distribution of fossil pollen data used here. A) Map indicating the spatial distribution of sites recovered from the Neotoma Paleoecology Database, as well as the case study regions (WCAN/AK: Western Canada and Alaska; PCCS: Pacific Coast, Cascades, and Sierra Nevada; ECAN: Eastern Canada; and NEUS/SEC: North-Eastern US and South-Eastern Canada). B) Temporal extent of all sites, in which each site is represented by a horizontal gray line between the site's oldest and youngest Holocene samples. Sites are ordered by age of the oldest sample, reported as ka. Samples older than 11.7 ka are not shown.



344 **Figure 2:** Interpolated REVEALS-based estimates of fractional cover for evergreen trees and shrubs  
 345 (ETS), summergreen trees and shrubs (STS), and open land (OVL). Laurentide and Cordilleran Ice Sheet  
 346 extents (Dalton et al., 2020) are indicated by gray polygons. Estimates are presented on a 1°x1° grid, for  
 347 selected time periods, with ages reported as ka. Map ordering follows the geological convention of oldest  
 348 maps at bottom. The spatial domain for interpolation includes all unglaciated locations in North America  
 349 between 17 and 79°N.

350  
 351  
 352



353  
 354 **Figure 3:** (a) Holocene trends in relative-to-modern mean land cover for weighted cover of all trees  
 355 and shrubs (ATS) in North America (pink curve with 95% uncertainty envelope), expressed relative to  
 356 present unglaciated land area (see Methods), and the fraction of unglaciated land relative to continental  
 357 land area (black). ATS is the sum of the evergreen and summergreen trees and shrubs. (b)

Trends in the ~~relative-to-modern mean landarea-weighted~~ cover of evergreen (ETS), summergreen (STS) and open vegetation/land (OVL) for North America. For both plots, land cover estimates are based on ~~the~~ interpolated data for the spatial domain shown in Fig. 2.

### 3.2 Continental-scale trends in Holocene land cover

At a continental scale, the spatial configuration of land cover in North America has been broadly stable during the Holocene (Fig. 2). Persistent features include belts of high evergreen tree cover in the mountainous West and Canada, high summergreen tree cover in the eastern US, moderate to high cover of summergreen trees and shrubs in Alaska and northern Canada, and high proportions of open vegetation in the Great Plains, western Alaska, and Arctic Canada (Fig. 2).

However, despite this broadly stable spatial configuration, there were large continental-scale changes in the ~~relative-to-modern area-weighted~~ fractional ~~cover of~~ land cover types in North America during the Holocene, particularly during the early Holocene (Fig. 3a, Supp. Fig. ~~S3S5~~). From 11.7 to 7 ka, the ~~relative-to-modern area-weighted~~ fraction of forested land cover increased from about ~~4256~~ to ~~691~~%. This increase closely tracked the increase in available land surface area, as the Laurentide Ice Sheet retreated (Fig. 3a) (Dalton et al., 2020). During the early Holocene, the largest gains in forest cover were across deglaciated western Canada and ice-marginal areas in eastern Canada (Fig. 2, Supp. Fig. ~~SS53~~). Forest cover continued to expand between 7.5 and 4.5 ka (Fig. 3a), even though the Laurentide Ice Sheet had disappeared by ~6 ka, with the largest afforestation in the northern Great Plains of central Canada and in recently deglaciated regions of eastern Canada (Fig. 2, Supp. Fig. ~~S3~~ ~~SS5~~). Continental-scale forest cover remained stable from 4.5 to 1.5 ka, then declined after 1.5 ka (Fig. 3a). The late-Holocene decline in forest cover was most pronounced in the eastern US and in Arctic and boreal Canada, where open vegetation began increasing after 4 ka (Fig. 2, Supp. Fig. ~~S3S5~~, ~~SS64~~).

Within these Holocene trends, the three land cover types followed differing trajectories (Fig. 3b). All three show a strong increase between 11.7 and 7.5 ka in their ~~relative-to-modern landarea-weighted~~ ~~relative~~ cover, again tracking ice retreat. After 7.5 ka, however, the three trajectories diverged. Evergreen trees and shrubs continued to rise slowly but steadily from 7.5 to 2 ka, then declined slightly (Fig. 3b). Summergreen trees and shrubs reached peak ~~extentarea-weighted cover~~ at 4.5 ka, then declined, with an accelerated decline after 1.5 ka. The proportion of open lands remained largely stable from 7.5 to 4.5 ka, then increased, with an accelerating increase after 2 ka. A close examination of the continental-scale anomaly maps (Supp. Figs. ~~S3S5~~, ~~S4S6~~) ~~indicates suggests~~ a fairly complex spatial mosaic for each land cover type, with the continental-scale trends emerging from a welter of regional-scale phenomena. For example, widespread gains in evergreen trees and shrubs across much of Canada

392 and Alaska between 6 and 4 ka were partially offset by large losses in the eastern US and southeastern  
393 Canada over the same period (Supp Fig. [SS5](#)). Because different plant species predominate in different  
394 regions of North America, these continental-scale trends in land cover were the emergent outcomes of  
395 individualistic species-level responses to changing climates and, in some places, intensifying land use  
396 (Williams et al., 2004).  
397

### 398 3.3 Regional case studies

#### 399 3.3.1 Northeastern US & Southeastern Canada (NEUS/SEC)

400 For the mostly forested NEUS/SEC, after initial afforestation and loss of open lands during the early  
401 Holocene (10 to 8 ka), the ~~principal primary-central~~ dynamic has been shifts in the relative cover of  
402 evergreen and summergreen trees and shrubs (Fig. 4a, Supp. Fig. [SS75](#)). The ~~relative-to-time-t~~ fractional  
403 cover of evergreen trees and shrubs declined throughout the early to middle Holocene (10 to 4 ka) in the  
404 southern part of the domain, which intensified to widespread loss across the NEUS/SEC (6 to 4 ka), but  
405 then reversed after 4 ka, with recovery and regrowth of evergreen trees and shrubs (Fig. 4a). The western  
406 increase in open lands between 8 and 4 ka is caused by the eastward expansion of prairie due to drier  
407 conditions (Williams et al., 2009b), while the decrease in open lands from 4 to 0.5 ka is due to increased  
408 moisture availability that resulted in a westward shift of the prairie-forest border and increase in  
409 summergreen forest taxa in the eastern Midwest (Umbanhowar et al., 2006). Overall, the low proportions  
410 of open vegetation in the NEUS/SEC from 9 ka until European settlement (Fig. 4B) likely represents  
411 these western prairies and local wetlands, which expanded in the late Holocene in parts of the NEUS/SEC  
412 (Brugam and Swain, 2000; Ireland and Booth, 2010), rather than grasslands or other open lands in eastern  
413 deciduous forests (Faison et al., 2006).

414 At a taxon level, most changes in evergreen cover can be attributed to regional declines of cold-  
415 tolerant conifers such as *Picea glauca*, *P. mariana*, and *P. rubens* (white, black, and red spruce); *Pinus*  
416 *banksiana*, *P. resinosa*, and *P. strobus* (jack, red, and white pine), and *Abies balsamea* (balsam fir)  
417 between 10 and 6 ka (Fig. 4c, Supp. Fig. [SS86](#)) (Spear et al., 1994; Jackson and Whitehead, 1991;  
418 Jackson et al., 1997). These changes, combined with the zonal pattern of evergreen expansion in the  
419 northern NEUS/SEC and declines in the southern part, suggest that much of the evergreen tree and shrub  
420 cover changes during the early to middle Holocene can be attributed to postglacial northward shifts in tree  
421 distributions in response to rising temperatures and deglaciation.

422 A second major feature is the well-known and dramatic expansion, collapse, and re-expansion of  
423 *Tsuga canadensis* (eastern hemlock), a cool-temperate conifer that typically occupies warmer climates  
424 than *Abies balsamea* (Thompson et al., 1999). Investigations continue into understanding the abrupt and  
425 widespread collapse in *Tsuga canadensis*, which differed from the overall evergreen trend. The collapse





426 occurred in less than 10 years at some sites (Allison et al., 1986) and is linked to regional shifts in water  
427 availability and temperature gradients (Booth et al., 2012; Foster et al., 2006; Haas and McAndrews,  
428 1999; Shuman et al., 2023). Initial hypotheses that a pest or pathogen such as *Lambdina fiscellaria*  
429 (hemlock looper) (Bhry and Filion, 1996; Davis, 1981; Anderson et al., 1986) caused the *Tsuga hemlock*  
430 decline have not been supported by recent investigations (Oswald et al., 2017), although insect remains  
431 are scarce in lacustrine archives. Understanding the causes of the hemlock collapse is outside the scope of  
432 this paper; however, this paper shows its importance, in that a single-species collapse fundamentally  
433 altered the functional composition, ecosystem phenology, and land cover of the NEUS/SEC for thousands  
434 of years.

435 Among summergreen taxa, these REVEALS reconstructions indicate a strong growth in *Acer*  
436 (maple) cover in the NEUS/SEC until 4 ka and declines thereafter. *Acer saccharum* (sugar maple) is  
437 probably the dominant taxon driving this curve, with *A. rubrum* (red maple) increasing in the late  
438 Holocene (Finkelstein et al., 2006). *Quercus* (oak) abundances were high between 9 and 4.5 ka, then  
439 steadily declined, while *Betula* (birch) abundances remained relatively stable. *Fagus grandifolia*  
440 (American beech) abundances continued to steadily increase throughout this period until 3 ka, then began  
441 declining after 1.5 ka, along with *Tsuga canadensis*, while *Abies balsamea* abundances increased.  
442 Disturbance-related taxa and indicators of open vegetation such as *Ambrosia* (~~ragweed~~) and *Rumex*  
443 (sorrel, not shown) begin increasing at ca. 0.35 ka (1600 CE). These late-Holocene changes in forest  
444 composition can ~~be~~ plausibly be attributed to regional cooling (explaining the increase in *Abies balsamea*)  
445 and perhaps also intensified human land use and disturbance, with the relative importance of these drivers  
446 varying at subregional scales and among taxa (Oswald et al., 2020; Mottl et al., 2021).

447 An intriguing feature of these REVEALS reconstructions is the inference of *Abies balsamea* and  
448 *Acer* ~~spp.~~ as the most common evergreen and summergreen tree taxa in the NEUS/SEC, given that *Picea*,  
449 *Pinus*, and *Quercus* are more abundant in pollen and macrofossil records and prior site-level and regional  
450 syntheses of Holocene pollen records have emphasized their dynamics (Jackson et al., 1997; Jackson and  
451 Whitehead, 1991; Payette et al., 2022; Spear et al., 1994). Witness-tree data in the NEUS also indicate  
452 that *Fagus* and *Quercus* were the most abundant broadleaved taxa at time of European settlement  
453 (Thompson et al., 2013). One possible reason for the higher levels of *Abies* and *Acer* reported here is that  
454 our NEUS/SEC domain extends a bit farther north than prior reconstructions that have focused more on  
455 central and southern New England. A second possibility is that the parameterizations for *Abies* and *Acer*  
456 are incorrect, causing the coverages of these taxa to be overestimated (see Discussion).

457





**Figure 4:** (a) Land cover anomaly maps for the northeastern US and southeastern Canada (NEUS/SEC) case-study region. Maps show the anomalies in fractional cover for each land cover class for pairs of indicated time intervals. Spatial resolution is 1°x1° and time units are ka. (b) Holocene trends in the

466 | ~~mean~~ relative-to-time-t mean fractional cover of the three land cover types (ETS, STS, OVL) and the  
467 | ATS sum for the uninterpolated REVEALS grid cell estimates across the NEUS/SEC region. (c)  
468 | Holocene trends for the REVEALS relative-to-time-t abundance estimates for the six most commonly  
469 | occurring taxa in the region. Line color indicates assignment of individual taxa to land cover types. For  
470 | taxon-level maps, see Supp. Fig. S86.  
471 |

### 472 | 3.3.2 Eastern Canada (ECAN)


473 | In Eastern Canada (ECAN), immediately north of the NEUS/SEC, the relative-to-time-t fraction of  
474 | evergreen trees and shrubs increased ~~expanded~~ across the entire region until 4 ka, with slower and more  
475 | spatially heterogeneous expansion of evergreen cover between 4 and 2 ka (Fig. 5a, Supp. Fig. S897).  
476 | After 4 ka, open vegetation expanded (Fig. 5a, 5b). This expansion of evergreen trees and shrubs can be  
477 | traced to separate phases of expansion for *Abies balsamea* and *Picea*, with *A. balsamea* expansion mostly  
478 | between 11.5 and 7.5 ka (in the southern portion) and *Picea* expansion mostly between 7.5 and 3 ka,  
479 | across the area and especially in the central portion (Fig. 5c, Supp. Fig. S8S10). Increasing Cyperaceae  
480 | abundance after 3.5 ka signals the expansion of forest-tundra. Major hardwood summergreen taxa  
481 | include *Betula* and *Alnus*. *Alnus* abundances expanded until 6 ka, then slowly declined, while birch  
482 | abundances expanded between 8.5 and 6 ka, then also declined. These declines in *Alnus* and *Betula* after  
483 | 6 ka co-occurred with expansions of *Picea* and Cyperaceae, suggesting that the regional vegetation  
484 | shifted from more of a mixed forest or woodland in the early Holocene, to evergreen coniferous forest,  
485 | forest-tundra, or tundra in the middle to late Holocene. Note that *Populus* is not included in these  
486 | REVEALS reconstructions, but it is an important taxon in ECAN (Peros et al., 2008), and its omission  
487 | may cause an underestimate of summergreen cover.



488 | These vegetation changes in ECAN can be attributed to a combination of deglaciation, changing  
489 | temperatures, and fire history. The Laurentide Ice Sheet collapsed at 8.4 ka and the last remnants of the  
490 | Labrador Dome disappeared from northern Quebec by 5.7 ka (Dalton et al., 2020). The replacement of  
491 | open land by forest cover during the early Holocene was driven primarily by increases in evergreen trees  
492 | and shrubs. The open forests at this time lack modern analogues and varied spatially in taxonomic  
493 | composition, with more Abies fir and Betula birch to the east and Picea spruce toward the west (Richard  
494 | et al., 2020). By 7.5 ka, closed forests developed in response to warmer temperatures. This was followed  
495 | by a decrease in summergreen taxa in the southern portion of the boreal forest as well as a decline in  
496 | Abies fir in the lichen woodland and feathermoss forest toward the west (Fréchette et al., 2021; Richard et  
497 | al., 2020). In the lichen woodlands of the north, shrub Betula birch and Betula papyrifera (paper birch)  
498 | decreased after 6 ka, although remaining high in the southwest (Fréchette et al., 2018). Near treeline,  
499 | Picea spruce forests were more open between 7 and 4 ka, with *Alnus* and shrub *Betula* more abundant

500 | (Fréchette et al., 2018; Gajewski, 2019). In ECAN, ~~Picea spruce~~ abundances reached a maximum  
501 between 4 and 2 ka (Fréchette et al., 2018), with the southern portion of the forest tundra becoming lichen  
502 woodland at this time. After 2 ka, *Picea* abundances at northern sites declined and open lands increased  
503 in response to cooling, but the timing and rate of *Picea* losses varied among sites and is governed in part  
504 by fire history (Gajewski et al., 2021; Gajewski, 2019).

505

506





**Figure 5:** (a) Land cover difference maps for ETS, STS, and OVL for the eastern Canada (ECAN) case study. (b) Holocene trends in the relative-to-time-t mean land relative-cover of the three land cover types (ETS, STS, OVL) and the ATS sum for the uninterpolated REVEALS grid cell estimates across the


515 ECAN region. (c) Holocene trends for the REVEALS abundance estimates for the six most commonly  
516 occurring taxa in the region. For both panels, figure conventions follow Figure 4. For taxon-level maps,  
517 see Supp. Fig. [8S10](#).



### 519 3.3.3 Western Canada and Alaska (WCAN/AK)

520 In western Canada and Alaska (WCAN/AK), [the relative-to-time-t mean land cover for](#) evergreen trees  
521 and shrubs rapidly expanded between 11 and 8 ka, while the coverage of summergreen trees and shrubs  
522 and open lands decreased (Fig. 6a, 6b, Supp. Fig. [S9S11](#)). Most of Alaska was not glaciated during the  
523 Last Glacial Maximum; however, much of western Canada, the Brooks Range in Alaska, and south-  
524 coastal Alaska were covered by ice (Dalton et al., 2020). Evergreen tree species such as *Picea glauca*  
525 (white spruce) persisted in this region in local microrefugia (Anderson et al., 2006), then expanded their  
526 ranges during end-Pleistocene warming and deglaciation. Other evergreen taxa such as *Pinus contorta*  
527 ([lodgepole pine](#)) expanded northward with deglaciation (MacDonald and Cwynar, 1991). Trends in taxon  
528 abundances differ substantially between the western and eastern subregions (Fig. 6c, Supp. Fig. [SS129](#)).  
529 In the eastern subregion, changes in taxon abundances were muted, with a modest expansion in *Picea* and  
530 *Tsuga* between 11.5 and 7.5 ka, and a slow decline in *Betula* between 10 and 6 ka. *Tsuga* (mostly *Tsuga*  
531 *heterophylla*; western hemlock) occurs primarily along the coast, was abundant along the south coastal  
532 areas by 11 ka (Lacourse et al., 2012; Lacourse and Adeleye, 2022), arrived by 9.5 ka into southeast  
533 Alaska (Hansen and Engstrom, 1996; George et al., 2023), increased in abundance along the south-coastal  
534 areas by 8 ka, and then expanded during the mid- to late Holocene into south-central Alaska (Anderson et  
535 al., 2017) and the inland mesic forests of British Columbia (Rosenberg et al., 2003; Gavin et al., 2021).  
536 In the western part of this domain (mainly Alaska), changes in summergreen hardwood (mostly shrub)  
537 taxa predominate, with a major expansion of *Alnus* between 11.5 and 6.5 ka (Anderson and Brubaker,  
538 1994; Cwynar and Spear, 1995), and a modest expansion of *Betula* between 11.5 and 9 ka. These  
539 expansions were accompanied by declines in Poaceae and Cyperaceae, with continued decline in  
540 Cyperaceae until 6 ka. *Picea* abundances steadily increased until 4 ka. The net effect was a decline in  
541 open vegetation and expansion of evergreen and summergreen trees and shrubs during the early to middle  
542 Holocene, with apparent region-wide stability after 4 ka (Anderson et al., 2019).

Formatted: Font: Not Italic







**Figure 6:** (a) Land cover difference maps for the western Canada / Alaska (WCAN/AK) case study region. (b) Holocene trends in the relative-to-time-t mean fractional relative cover of the three land cover types (ETS, STS, OVL) and the ATS sum for the uninterpolated REVEALS grid cell estimates across the WCAN/AK region, with grid cells averaged separately for eastern and western subregions (dividing line

shown in (a) as ~~vertical~~ dashed line). (c) Holocene trends for the REVEALS abundance estimates for the six most commonly occurring taxa in the region, with grid cells averaged separately for eastern and western subregions. For both panels, figure conventions follow Figure 4. For taxon-level maps, see Supp. Fig. S129.

#### 3.3.4 Pacific Coast, Cascade, and Sierra Nevada Ranges (PCCS)

In contrast to the other regions, evergreen trees have dominated much of the land cover in the PCCS region since 12 ka (Fig. 7a, 7b, Supp. Fig. S143). In the northern subregion (often referred to as the Pacific Northwest), the ~~relative-to-time-t~~ proportions of cover types were fairly constant until 9 ka, after which summergreen and open land declined until 6 ka (Fig 7a, 7b). However, taxon-level changes were very dynamic. From 11.7 to 10.5 ka, the REVEALS-estimated abundance of *Pseudotsuga menziesii* (Douglas fir) more than doubled, reflecting its arrival, expansion, and northward migration (Gugger and Sugita, 2010), and replacing true firs (*Abies*) and *Picea* (Fig 7c, Supp. Fig. S142). In the coastal ranges, frequent forest fires in the early Holocene contributed to a *Pseudotsuga-Alnus rubra* (red alder) association in the mountains (Gavin et al., 2013; Long et al., 1998, 2007) and a ~~Quercus~~ *oak* savanna (not shown) that was common in the lowlands (Walsh et al., 2008, 2010; Giuliano and Lacourse, 2023). Summergreen taxa (mostly *Alnus rubra*) and open-land pollen indicators declined from 8 to 6 ka (Fig 7b), replaced by shade-tolerant *Abies*, *Tsuga*, and Cupressaceae, which is mostly western redcedar in this region (Fig. 7b) (Lacourse, 2009; Gavin et al., 2013; Worona and Whitlock, 1995). After 6 ka, the overall abundance of conifer taxa remained constant (Fig 7a, 7b), although the proportion of ~~Tsuga hemlock~~ and Cupressaceae ~~cedar~~ increased steadily through the late Holocene while *Pseudotsuga* declined, consistent with less fire and the persistence of old-growth forest (Whitlock, 1992; Lacourse and Adeleye, 2022). ~~until the logging of the last century, which manifested as a decline in evergreen forests~~ conifer and is associated with an increase in summergreen ~~trees and shrubs~~ forests and open land in the coastal ~~regions~~ forests (Fig. 7a) (Davis, 1973; Whitlock et al., 2018).

The most common taxon in these REVEALS reconstructions, *Abies*, varied little over the last 9 ka, but this genus represents several shade-tolerant species (*A. lasiocarpa* [subalpine fir], *A. grandis* [grand fir], *A. amabilis* [Pacific silver fir], *A. procera* [noble fir]) that collectively are common throughout the region. As noted for the NEUS/SEC, the reconstructed values of *Abies* are likely too high, because the REVEALS estimate of fall speed ~~is~~ ~~may be~~ overestimated. *Pinus* also represents several species and was most abundant in the dry eastern areas where it was consistently abundant during the Holocene (Minckley et al., 2007). *Pseudotsuga* and *Larix* have indistinguishable pollen morphologies and are therefore grouped in the reconstructions; however, *Larix* is limited to the eastern edge of this region.

Formatted: Font: 11 pt

Formatted: Font: Italic


586           The southern subregion (the Sierra Nevada, Klamath Mountains and California Coast ranges and  
587 interior Great Basin) supported roughly equal amounts of conifer forest and open land that overall had  
588 minor relative changes over the Holocene (Fig. 7a). Summergreen trees and shrubs have been infrequent  
589 contributors to land cover in this subregion. At the taxon-level, *Abies* is again the most common  
590 component of reconstructed evergreen land cover across the region, although its abundance may be  
591 overrepresented in this parameterization of REVEALS (Fig. 7b). *Pinus* and *Quercus* were most abundant  
592 in the early Holocene, forming open forests and woodlands, especially in the Klamath Mountains and  
593 Sierra Nevada (Anderson, 1996; Briles et al., 2008) fire-prone forests east of the Cascade Range (Walsh  
594 et al. 2015). Most of the eastern portion of this subregion is the Great Basin shrub steppe, where the few  
595 pollen records that exist are from high-elevation sites and low-elevation wetlands that are sensitive to  
596 fluctuating water tables, recorded as large fluctuations in Cyperaceae pollen. Increases in open-land cover  
597 taxa (*Cyperaceae*, *Poaceae*, *Asteraceae*) between 10-7.5 ka and after 3 ka in this region may reflect either  
598 local expansion of alpine meadows and wetlands or expansion of steppe more broadly (Mensing et al.,  
599 2008; Brugger and Rhode, 2020; Minckley et al., 2007; Thompson, 1992). ~~Note, However, that~~ desert,  
600 steppe, and other open-land arid ecosystems are likely to be underrepresented in these reconstructions,  
601 due to a scarcity of dryland sites. Furthermore, bare soil and rock cannot be detected using pollen-based  
602 land-cover mapping, and so this land cover type is missing in these reconstructions.

Formatted: Font: Italic

Formatted: Font: Italic



Formatted: Font: Italic

605



606

607



**Figure 7:** (a) Land cover difference maps for the Pacific Coast, Cascade and Sierra Nevada (PCCS) case study region. (b) Holocene trends in the mean relative cover of the three land cover types (ETS, STS, OVL) and the ATS sum for the uninterpolated REVEALS grid cell estimates across the Pacific Coast.

[Cascade and Sierra Nevada \(PCCS\) case study](#) ~~PCCS~~ region, with grid cells averaged separately for northern and southern subregions (dividing line shown in ~~(ca)~~ as vertical dashed line). ~~(be)~~ Holocene trends for the REVEALS abundance estimates for the ~~ninesix~~ most commonly occurring taxa in the region, with grid cells averaged separately for northern and southern subregions. For both panels, figure conventions follow Figure 4. For taxon-level maps, see Supp. Fig. [S142](#). [\(c\) Land cover difference maps for the PCCS region.](#)

## 4. Discussion

### 4.1 Drivers of Holocene land cover change in North America: scaling from taxon-level regional dynamics to continental-scale trends

These continental-scale Holocene changes in land cover (Figs. 2-3) are an emergent outcome of the individualistic plant responses to deglaciation and multiple environmental changes, including seasonal temperatures, effective moisture, atmospheric carbon dioxide, [sea level rise](#), soil development, fire and other disturbance regimes, and, at some locations during the late Holocene, human land use. As shown here (Figs. 4-7), this interplay differed spatially across North America, often with opposing trends (e.g. simultaneous increases in evergreen cover in some regions and decreases elsewhere) that partially offset at the continental scale.

The first-order continental-scale drivers of Holocene land cover change in North America were changing climates and the retreat and disappearance of the Laurentide Ice Sheet, particularly during the early Holocene (11.7 to 7.5 ka) and lasting until 5.7 ka, with the disappearance of remnant ice in the Labrador Dome in northern Quebec (Dalton et al., 2020). Deglaciation and an overall increase in land availability explains why all land cover types show an increase in ~~relative-to-modern-area-weighted~~ cover during the early Holocene, with evergreen taxa showing the greatest gain (Fig. 3). Continental temperatures also closely tracked the decline in ice area (Marsicek et al., 2018), and vegetation responded to both factors. Superimposed upon the early-Holocene increase in ~~relative-to-modern-area-weighted~~ cover for all PFTs are gains and losses for each PFT at subregional scales (Figs. 4-7), which can be linked to climate-driven changes in plant abundances at local to landscape scales that scaled upwards to continental-scale shifts in plant distributions, with both within-range shifts in dominance and leading-edge and trailing-edge range dynamics for individual plant taxa (Payette et al., 2022; Williams et al., 2004; George et al., 2023; Dallmeyer et al., 2022; Anderson et al., 2017).

Leading-edge dynamics and range expansion of tree taxa appear to have been primarily controlled by the increasing availability of deglaciated land area, end-Pleistocene warming, the declining influence of ice sheet on regional climates and moisture availability (Alder and Hostetler, 2015; Bartlein

et al., 2014), and on-going expansion of plant taxa into areas of increased climate suitability (Payette et al., 2022; Williams et al., 2004; George et al., 2023; Dallmeyer et al., 2022). Declining ice extent favored moisture advection into eastern areas where most of the increase in summergreen tree taxa took place and reduced it in midcontinental North America, where most of the open vegetation increase developed (Shuman and Marsicek, 2016; Shuman et al., 2002; Liefert and Shuman, 2020). These changes in the patterns of moisture availability simultaneously favored the large increase from 10 to 6 ka in summergreen trees and shrubs in eastern North America south of the former Laurentide Ice Sheet and the increase in open land in the mid-continent (Supplementary Fig. 52-36).

Declines in abundance and trailing-edge dynamics of tree taxa were likely governed by a combination of declining climate suitability and, in some places, fire regimes, in which many local populations failed to re-establish after one or more fire events. For example, in the high northern latitudes of Quebec and Labrador, the expansion of open lands over the last three thousand years (Fig. 5) can be attributed first to declining summer insolation and temperatures, caused by precessional changes in the Earth's orbit (Payette, 2021), but at local to landscape and centennial scales, loss of forest cover was asynchronous and caused by individual fire events (Payette et al., 2008; Gajewski et al., 2021). Similarly, in the southern Great Lakes region and NEUS, the late-Pleistocene to early Holocene transition from conifer-dominated forests and parklands to summergreen forests (Fig. 4) was driven by rising temperatures and changes in effective moisture (Shuman et al., 2002, 2019), but accelerated locally by intensified fire regimes, with the timing and rate of conversion varying among sites (Jensen et al., 2021; Clark et al., 1996). The expansion and then retreat of the Great Plains prairies (Fig. 4) was driven by early Holocene aridification and middle- to late-Holocene increases in moisture availability, while fire may have facilitated prairie expansion, but delayed its retreat (Williams et al., 2009b; Nelson et al., 2006; Umbanhowar et al., 2006). Of course, throughout North America, forests and fire dynamically co-existed and interacted during the Holocene (Iglesias and Whitlock, 2020; Kelly et al., 2013), so whether fire causes transformative shifts in ecosystem type depends on its synergistic interactions with directional changes in climate and other factors (Napier and Chipman, 2021). Regardless of the role of fire or other disturbances, regional shifts in temperature and moisture availability are usually the first-order predictors of Holocene changes in vegetation composition (Dean et al., 1984; MacDonald, 1989; Calder and Shuman, 2017; Shuman et al., 2004; Nelson et al., 2006).

The relative proportion of evergreen and summergreen cover types may have been affected by changes in the length and intensity of the growing season during the Holocene, which is regulated by precessional variations in insolation. In particular, the broad peak in summergreen tree cover between 8.5 and 3.5 ka (Fig. 3) is consistent with the hypothesis that summergreen tree and shrub abundances in North America are partially regulated by summer insolation and temperatures (Delcourt and Delcourt, 1994;



Williams et al., 2001). Summer insolation reached higher peak intensity but has a shorter seasonal duration in the early Holocene than during the late Holocene (Huybers, 2006; Jackson et al., 2009), which may have favored plants with summer deciduous phenology that could more effectively exploit available energy during a briefer but more intense growing season (Delcourt and Delcourt, 1994; Williams and Jackson, 2007; Edwards et al., 2005). Indeed, pollen-reconstructed and CCSM3-simulated changes in growing-degree days also peaked during the mid-Holocene when forest cover was greatest, and several millennia after peak summer temperatures, likely because of differential responses of maximum summer temperatures and total growing season warmth to orbital and other forcings (Marsicek et al., 2018). This may in turn suggest a seasonal-scale feedback loop between summer insolation, deciduous phenology, and total summer warmth (see Section 4.2).

During the late Holocene, the growth of Indigenous populations and intensification of human land use in the Americas~~intensified had, with~~ increasing effects upon land cover. U~~and u~~nderstanding the interactions among past climate change, fire regime, and ~~human~~ land use is a highly active area of research. Rates of vegetation change worldwide began to increase between 4.6 and 2.9 ka (Mottl et al., 2021), consistent with growing intensification and extent of land use (Stephens et al., 2019). The 7 to 10 ppm decrease in global CO<sub>2</sub> between 1570 and 1620 CE has been attributed to ~~the~~ land abandonment and reforestation due to mass mortality of Indigenous populations in the Americas, caused by the spread of multiple pathogens (Lewis and Maslin, 2015). The best evidence for dense human populations and extensive land clearance in the Americas comes from Central and tropical South America (Islebe et al., 1996; McMichael, 2020).

In the land cover reconstructions presented here, the increase in prevalence of open lands after 0.5 ka in North America and some subregions (Figs. 3, 4, 7) is attributable to land use, but the role of land use for earlier time periods remains unclear. In North America, the effects of human~~II~~and use prior to EuroAmerican settlement clearly altered land cover are detectable at some sites in North America, but land use effects are not easily detected at the regional to continental scales addressed here. In North America, the distribution of radiocarbon dates from archaeological contexts suggest that population levels remained relatively low ~~and increasing slowly~~ until ~7 ka, with ~~further~~ increases between 7 and 2 ka, and then rapidly increase~~d~~ after 2 ka (Peros et al., 2010). Indigenous cultures in North America clearly engaged in activities that, in some areas, modified the composition and structures of forested and unforested landscapes for millennia (Ellis, 2021; Delcourt et al., 1986; Leopold and Boyd, 1999; Munoz et al., 2014). In densely populated regions, such as lands of the Iroquois Confederacy, southern Ontario, or the Cherokee Nation, Cahokia and the American Riverbottom, land use altered vegetation structure and composition at local to landscape scales, through management of fire regimes (Roos et al., 2018; Anderson and Carpenter, 1991), land clearance for agricultural crops (McAndrews and Turton, 2007), and

silviculture, e.g. favoring the spread of nut-bearing trees (Munoz et al., 2014; Black et al., 2006). In western North America, the effects of anthropogenic activity on vegetation and fire is an active area of research, with several paleoecological studies indicating changes consistent with human influences on forest composition (Walsh et al., 2015; Knight et al., 2022; Lacourse et al., 2007). Because the scale of human action was strongest at local to landscape scales and varied in intensity within and among regions, its detection often requires highly-focused, local- to regional-scale studies (Oswald et al., 2020; Roos, 2020; Lacourse et al., 2007; Knight et al., 2022; Munoz and Gajewski, 2010). These studies clearly indicate a high intersite variance in the level and detectability of human impact. At the continental scale, (Gajewski et al., 2019) did not find clear correlations between human population abundance and pollen abundance, including taxa of economic use or disturbance taxa, and concluded impacts ~~occurred~~<sup>were</sup> at local to regional scales. Thus, at the regional to continental scales considered here, it is an open question whether the rich relationships and diverse activities engaged by communities within their homelands led to detectable changes in land cover types. We expect that pollen-based land cover reconstructions will continue to contribute to interdisciplinary approaches to address anthropogenic fire and land cover change during the Holocene (Snitker et al., 2022).

In contrast, after EuroAmerican arrival and expansion from 1492 onwards, North American ecosystems were massively transformed (Stegner and Spanbauer, 2023) by multiple anthropogenic processes. These include widespread forest conversion to agricultural and pastoral lands, intensive forest harvesting, massive hydrological change through dam construction and wetland drainage, introduction of exotic species and pests, and both greatly increased fire activity and fire suppression (Klein Goldewijk et al., 2011; Foster and Aber, 2004). These land cover changes can be detected as an increase in open land from 0.5 ka onwards (Figs. 3, 4, 7).

## 4.2 Biophysical and biogeochemical vegetation-atmosphere feedbacks to Holocene climates

### 4.2.1 Holocene vegetation-atmosphere feedbacks: prior work and need for data constraints

This paper and the other REVEALS-based reconstructions for the Northern Hemisphere (Githumbi et al., 2022a; Li et al., 2023) are now laying the foundation for a next generation of Holocene vegetation-atmosphere research with well-constrained land surface data. Many studies have explored the potential impacts of vegetation-atmosphere feedbacks on Holocene climate dynamics at hemispheric to global scales, but these studies have generally not employed well-constrained land cover reconstructions. Treeline shifts have been ~~identified~~<sup>recognized</sup> as an important regulator of Holocene vegetation-atmosphere feedbacks in the high northern latitudes (Kutzbach et al., 1996). ~~Earth-system model experiments with prescribed vegetation scenarios have shown that, in the northern latitudes (45–90N), changes in forest cover are the largest contributor to changes in net land surface radiation through snow~~

749 ~~masking and effects on surface albedo (Bathiany et al., 2010).~~ In mid-Holocene atmosphere-vegetation  
750 model simulations, strong snow masking resulted in warming three times higher than those with weak  
751 snow masking (Otto et al., 2011). ~~Additionally, empirical estimates of surface albedo feedbacks in high~~  
752 ~~latitudes are stronger than predicted by climate models (Hogg, 2022).~~ More recent work has highlighted  
753 the importance of the type and density of forest cover in determining the albedo feedback and magnitude  
754 of snow masking (Lorantý et al., 2014; Alessandri et al., 2021). Hence, because snow masking and  
755 surface albedo is an important regulator of surface-atmosphere feedbacks in climate models, it is also an  
756 important source of uncertainty in Holocene climate simulations, because of the limited availability of  
757 well-constrained reconstructions of past changes in vegetation type, structure, and density. Vegetation  
758 structure and topographic complexity also jointly govern surface roughness, which affects lower  
759 atmosphere temperature, humidity, wind speed, and soil moisture (Bonan, 2015) and is the dominant  
760 land-cover influence on micrometeorological processes (Chen and Dirmeyer, 2016). The effects of  
761 changing surface roughness on past climate variability remains largely unexplored. This new generation  
762 of land cover reconstructions provided by LandCover6k is intended to support future a new generation of  
763 Earth system model experiments into past vegetation-atmosphere feedbacks, with first results now  
764 available for Europe (Strandberg et al., 2022).

765 The role of vegetation feedbacks on terrestrial carbon sequestration and Holocene climate  
766 variations also remains largely unexplored, despite ice-core evidence that Holocene CO<sub>2</sub> increased  
767 from between 260 to and 285 ppm over the period from between 8 to and 1 ka (ca. 53 GtC) and no clear  
768 understanding of the source(s) of that carbon the source of where that carbon came from. Initial work,  
769 using carbon isotopic evidence and carbon cycle models, indicated that changes in terrestrial biomass  
770 were responsible for about two-thirds of this increase (Indermühle et al., 1999). More recent Earth-  
771 system model simulations indicate an ocean source through a process of elimination, given simulations  
772 that both the atmosphere and terrestrial biosphere were gaining carbon between 8 and 2 ka (Brovkin et al.,  
773 2019). Prior data-based reconstructions of terrestrial carbon sequestration, mostly restricted ~~so~~ to the  
774 northern middle and upper latitudes, indicate net carbon sequestration in these regions for most of the  
775 Holocene. Peatlands in the upper latitudes of the Northern Hemisphere appear to have been continuously  
776 sequestering carbon throughout the Holocene, with a total sequestration across the Holocene of  
777 500±100 GtC (Loisel et al., 2017; Yu, 2012). Prior reconstructions from pollen data and the LPJ dynamic  
778 vegetation model suggest a terrestrial sequestration of 13.9 GtC between 9 and 6 ka and a 3.5 GtC release  
779 after 6 ka, for aboveground biomass in the Northern Hemisphere >45°N (Williams et al., 2011a).  
780 Regional reconstructions for eastern Canada indicate a peak in terrestrial biomass at 7 ka, with a ca. 8.5  
781 GtC sequestration due to afforestation between 11 and 7 ka and a ca. 2 GtC release after 7 ka (Blarquez  
782 and Aleman, 2015), suggesting that postglacial boreal forest dynamics in this region were a primary

contribution to Northern Hemisphere aboveground carbon sequestration. In the upper Midwest, aboveground biomass doubled between 8 and 0 ka, sequestering 1.8 GtC (Raiho et al., 2022). Given these various well-constrained Holocene sinks in the atmosphere and terrestrial biosphere, the key unknown is the corresponding sources, with potential options including changes in ocean alkalinity (Brovkin et al., 2019), subtropical wetland reductions with waning Northern Hemisphere monsoons (Guo et al., 2012), and intensifying land use (Ruddiman, 2003). Balancing the Holocene carbon cycle in the atmosphere-biosphere-ocean system is a major problem in Earth system science. This new generation of LandCover6k vegetation reconstructions, combined with Earth system models, are ~~the~~ our best avenue forward—this promise to sharpen our understanding of the role played by surface-atmosphere feedbacks in Holocene climate dynamics, both biogeophysical and biogeochemical, with first results from Europe already underway (Strandberg et al., 2022).

#### 4.2.4.2 Assessing recent modeling studies of the Holocene ~~t~~Temperature ~~c~~Conundrum

The Holocene ~~t~~Temperature ~~c~~Conundrum has been a major focus of paleoclimatic research over the last decade (Kaufman and Broadman, 2023; Liu et al., 2014), and ~~recent~~ research has invoked vegetation-atmosphere feedbacks to resolve this conundrum (Thompson et al., 2022). The Holocene ~~t~~Temperature ~~c~~Conundrum involves a discrepancy between proxy-based reconstructions of global temperature changes, mostly based on marine records, which indicate a mid-Holocene maximum and mid- to late-Holocene cooling (Marcott et al., 2013), while transient model simulations show small but steady warming throughout the Holocene (Liu et al., 2014). Many explanations for this discrepancy have since been proposed (Kaufman and Broadman, 2023). Recent prescribed-vegetation experiments in climate models produce an early Holocene warming to a Holocene Thermal Maximum, followed by cooling towards the pre-industrial Holocene (Thompson et al., 2022), consistent with paleoclimatic proxies. Conversely, experiments that include drivers such as dust, ice cover, orbital forcing, and greenhouse gasses without accounting for Northern Hemisphere vegetation changes do not result in a mid-Holocene thermal maximum (Thompson et al., 2022).

Our reconstructions suggest, however, that for North America, the prescribed vegetation maps used by Thompson et al. (2022) overstate the magnitude of Holocene vegetation change. These prescribed-vegetation experiments for 9 and 6 ka fully replace all C<sub>3</sub> grasses with boreal forest for all locations north of 50°N (Thompson et al. (2022), Supp. Fig. ~~S97B~~). This pattern is qualitatively consistent with the early Holocene afforestation reported here (Fig. 2) but is inconsistent with the demonstrated persistence of tundra throughout the early and middle Holocene, particularly in WCAN/AK (Figs. 2, 6) and Canadian High Arctic (Fig. 2). Similarly, the prescribed full replacement of C<sub>3</sub> grasslands with temperate deciduous forest for the 9 and 6 ka experiments (Thompson et al. (2022), Supp.

Fig. S97B) is inconsistent with clear palynological evidence ~~for~~ the establishment of the Great Plains grassland by the early Holocene and prairie expansion during the early to middle Holocene (Fig. 2, Supp. Fig. S45) (Williams et al., 2009a). Hence, the Thompson et al. (2022) simulations should be viewed as useful experiments that show the potential sensitivity of Holocene climates to large vegetation changes, but these experiments likely overestimate the contribution of vegetation feedbacks to resolving the Holocene ~~Temperature~~ Conundrum.

#### 4.24.3 Understudied Holocene vegetation-atmosphere feedbacks in North America

Our reconstructions also highlight several major features of North American vegetation dynamics that may have underappreciated effects on Holocene vegetation-atmosphere feedbacks and climate dynamics at regional to continental scales. First, the mid-Holocene decline and recovery of *Tsuga canadensis* (~~eastern hemlock~~) in the northeastern United States is an example of how single-species dynamics can drive fundamental changes in vegetational structure. Although the patterns and drivers of the *T. canadensis* decline have been extensively studied (Oswald and Foster, 2012; Oswald et al., 2017; Booth et al., 2012; Foster et al., 2006), the effects of the *T. canadensis* decline on Holocene climates are unknown. As the dominant evergreen conifer of cool-temperate eastern North America, the decline of *T. canadensis* at ca. 5 ka is the primary driver of the loss of ETS between 6 and 4 ka, while the gradual recovery of *T. canadensis* after 5 ka drives the corresponding increase of ETS (Fig. 4). This shift in dominance between summergreen and evergreen trees and shrubs, in turn, regulates the overall albedo of the land surface and particularly its seasonal range. During times of foliage, summergreen forests can have more than twice the albedo of evergreen forests (Hollinger et al., 2010), ~~and~~ ~~Summergreen forests also exhibit much greater seasonal variability in albedo,~~ with maximum values in full foliage being 20-50% larger than annual lows. ~~There are also differences in evapotranspiration rates between broadleaf summergreen forests and needleleaf evergreen forests, although these differences are variable based on stand age, structure, and meteorological conditions (Komatsu et al., 2007; Liu et al., 2003).~~

Second, the peak in summergreen tree cover between ca. 8.5 and 4.5 ka (Fig. 2) may indicate a seasonal-scale positive feedback loop between summer insolation, summergreen phenology, and summer temperatures. Studies have suggested that the late-Pleistocene to early-Holocene peak in summer insolation favored summergreen phenology and carbon acquisition strategies over evergreen strategies (Delcourt and Delcourt, 1994; Williams and Jackson, 2007). Studies from the Pacific Northwest and Alaska also show a peak in *Alnus* and other summergreen tree and shrub taxa during the early Holocene (Fig. 7c), coincident with the summer insolation maximum and local maxima in temperature (Gavin et al., 2013; Edwards et al., 2005; Whitlock, 1992). This climate-driven response, in turn, may have increased the seasonal range of albedo and surface temperatures, thereby further favoring summergreen strategies.

851 This summergreen feedback might have also contributed to some summer cooling, due to the higher  
852 summer albedo of summergreen trees (Hollinger et al., 2010). As an alternative hypothesis, Herzsuh  
853 (2020) suggested that the Eurasian distribution of evergreen spruce-dominated and deciduous larch-  
854 dominated evergreen forests was due to historical contingencies and alternate stable states. Although  
855 many studies have explored the effect of early to middle Holocene afforestation on vegetation feedbacks  
856 in the northern latitudes (Brovkin et al., 2009; Kutzbach et al., 1996), to our knowledge no paper has yet  
857 focused specifically on the seasonal feedback effects associated with shifting proportions of summergreen  
858 and evergreen trees and shrubs.

859

#### 860 4.3 Uncertainties and limitations in pollen-vegetation model reconstructions

861 Land cover reconstructions from pollen rely upon a variety of pollen-vegetation models (PVMs), some of  
862 which have well-understood limitations and uncertainties, and some of which are newer and are still being  
863 studied. The REVEALS PVM used here is has been widely adopted (Li et al., 2020; Githumbi et al.,  
864 2022b; Serge et al., 2023; Azuara et al., 2019; Hoevers et al., 2022) because it represents some of the  
865 taxon-level processes governing pollen production, transport, and representation (Prentice, 1985; Sugita,  
866 2007a). However, processes such as atmospheric transport are simplified (Jackson and Lyford, 1999) and  
867 key parameters such as pollen productivity estimates (PPEs) carry uncertainties (Wieczorek and  
868 Herzsuh, 2020; Broström et al., 2008; Hayashi et al., 2022) that translate to uncertainties in fractional-  
869 weighted land cover area. Most PPE estimates are generated from models fitted to spatial networks of  
870 vegetation surveys and surface pollen samples, and these PPE estimates depend strongly on models of  
871 pollen dispersal (Theuerkauf et al., 2012).

872 Such issues may explain some of the surprising aspects of the reconstructions presented here. For  
873 example, reconstructed *Acer* cover in the NEUS/SEC (Fig. 4c) is high compared to settlement-era  
874 estimates from witness trees and land surveys (Thompson et al., 2013; Paciorek et al., 2016). Similarly,  
875 reconstructed cover of *Abies* in both the NEUS/SEC and PCCS is higher than expected (Figs. 4c, 7c).  
876 *Abies* and *Acer* are notoriously underrepresented in fossil pollen assemblages (Bradshaw and Webb,  
877 1985), relative to independent surveys of tree abundance in the surrounding ecosystems, due to the low  
878 pollen productivity of maple trees relative to other taxa (Finkelstein et al., 2006; Liu et al., 2022) and high  
879 fall speeds of *Abies* (Jackson and Lyford, 1999). Hence, a key value of process-based PVMs, such as  
880 REVEALS, is the ability to correct for these known biases. However, REVEALS estimates are sensitive  
881 to parameter choices and it is possible that the estimates of *Abies* and *Acer* are too high. There are few  
882 estimates of *Abies* fall speeds compared to other taxa, and the fall speeds used here from Wieczorek and  
883 Herzsuh (2020) are similar to the values for *Larix* that initial testing indicated led to *Larix*  
884 | overrepresentation. [Larix is also challenging to model accurately in pollen-vegetation models, because its](#)

885 large pollen grains does not disperse far. Hence, local populations of larch growing at the site where a  
886 core was retrieved can be difficult to differentiate in pollen records, from those occurring across the  
887 broader source area. Moreover, the scarcity of Larix pollen grains means that the REVEALS  
888 reconstructions are highly sensitive to its parameterizations for Larix.

889 REVEALS is not spatial or temporal in nature, so there is no representation of spatiotemporal  
890 dependencies among site-level reconstructions (Supp. Fig. S2, S3). REVEALS estimates of fractional  
891 cover uncertainty are based on the total number of pollen grains counted in a sample and the error  
892 associated with the PPEs, but do not include process uncertainty or uncertainty in other model inputs.  
893 The GMRF serves as a post-hoc interpolator to generate spatially ~~temporally~~ complete vegetation  
894 reconstructions for each time period, but this approach does not mechanistically represent the underlying  
895 processes that link pollen to vegetation. Neither REVEALS nor the GMRF include soil, slope, aspect, or  
896 other edaphic controls on vegetation cover. The REVEALS-GMRF workflow as currently implemented  
897 does not account for pollen sample age uncertainty. While the Bchron age-depth modeling approach used  
898 Bchron is a Bayesian approach in nature and does result in that estimates posterior distributions of  
899 sample ages, propagating this through the REVEALS-GMRF workflow would require addition of  
900 temporal dependence in the interpolation model. Initial experiments to add temporal dependence have  
901 been computationally prohibitive; alternative model specifications that would make this possible are  
902 currently being tested. Even so, given the temporal grain of the time bins (500 years throughout the  
903 Holocene, except for the last 700 years, 100-350 years), we expect that within uncertainty the majority of  
904 sample ages will not shift among time bins. quantifies age

905 While REVEALS does quantify standard errors associated with relative abundance estimates,  
906 these standard errors are not considered by the REVEALS-GMRF approach. This means that uncertainty  
907 estimates from the REVEALS-GMRF approach quantify the uncertainty determined by the variability of  
908 REVEALS fractional land cover around the estimated spatial field (Supp. Fig. S4). Validation  
909 experiments would be useful to understand the impacts of sample unevenness; such experiments have  
910 been done using the REVEALS-GMRF approach for Europe (Pirzamanbein et al., 2018). This work  
911 generally showed little change in the predictions when data were withheld. These cross-validation  
912 experiments have a high computational burden. Future work aims to improve computational efficiency of  
913 the approach, at which point such experiments will be possible for larger domains like North America.

914 To address these limitations with the REVEALS workflow (spatial ~~and~~ temporal  
915 incompleteness, input parameter uncertainty, and uncertainty quantification), other forms of PVMs have  
916 been developed. ROPES uses pollen accumulation rates and REVEALS to estimate pollen productivity  
917 (Theuerkauf and Couwenberg, 2018). However, the dependence of ROPES on pollen accumulation rates  
918 may limit its widespread utility. At many sites, pollen accumulation rates have high uncertainties due to

variations in sedimentation rate, few radiometric dates and poor chronological controls, and spike counting uncertainties (Perrotti et al., 2022). STEPPS is a Bayesian spatio-temporal PVM (Dawson et al., 2016, 2019). While theoretically sound, STEPPS is computationally intensive and its estimates are dependent upon the density and quality of spatial forest compositional calibration datasets, limiting its applicability to regional-scale domains. Other spatial Bayesian models developed for high-resolution networks of pollen and vegetation data suggest that if the spatial site density is too low, STEPPS and similarly structured PVMs can over-estimate pollen dispersal distance (Liu et al., 2022).

#### 4.4 Future work

With North American REVEALS-based reconstructions now in place, all middle- and high-latitudes in the Northern Hemisphere have quantitative reconstructions of Holocene land cover that are well constrained by dense networks of fossil pollen records (Githumbi et al., 2022b; Li et al., 2023). The reconstructions for North America and Europe also have estimates of uncertainty from the spatiotemporal GMRF model (Pirzamanbein et al., 2018). This hemispheric coverage now enables the creation of a next generation of modeling scenarios, well-constrained by data, to explore land cover feedbacks in the Holocene climate system (Harrison et al., 2020), with relevance to both the Holocene temperature conundrum (Kaufman and Broadman, 2023; Liu et al., 2014) and the question of the missing Holocene carbon source (Brovkin et al., 2019; Indermühle et al., 1999; Ruddiman, 2003). These land cover reconstructions also help identify places where new local-scale, site-based research are needed to better understand human impacts, better constrain land cover changes in areas of sparse data and high uncertainty, and better understand how local and taxon-level dynamics scale up to affect regional- to continental-scale vegetation and climate dynamics.

Multiple opportunities exist to further enhance these reconstructions. First, this work also underscores the critical need for more work to better constrain PPEs and fall speeds, particularly for North American plant taxa. The REVEALS estimates in this analysis appear to be particularly sensitive to the parameterizations for underrepresented taxa such as *Abies* and *Acer*. Second, these land cover reconstructions could be further parsed into boreal, temperate, and subtropical components, given what is known about these individual pollen types and their climatic affinities. Note, however, that statistically modeling more finely resolved land cover groupings or taxa is complicated by the many zero-count observations in more finely resolved groupings, especially for such a large spatio-temporal domain (Haslett et al., 2021). Third, these pollen-based vegetation reconstructions would be improved by the incorporation of additional proxies, particularly in semi-arid regions, where carbon isotopic records from soils, lakes, speleothems, and other archives can provide information about the relative abundance of C3 and C4 grasslands (Braun et al., 2019; Nordt et al., 2007) and phytoliths can provide information



[about grassland composition](#) (Blinnikov et al., 2002; Evett and Bartolome, 2013), although multiple factors affect  $\delta^{13}\text{C}$  variations (Fohlmeister et al., 2020). Fourth, for some subregions where a dense network of high-resolution, well-dated records exists, such as New England (Oswald et al., 2018)

Lastly, given the maturation of pollen-vegetation modeling as a field of study and the emergence of multiple PVMs, there is now the opportunity for intercomparison studies among PVMs and, perhaps, the development of ensemble-based inferences of past vegetation cover. There is a long history of methodological comparisons in pollen-based paleoclimatic inferences and paleoclimatology more broadly (Chevalier et al., 2020). Few systematic intercomparisons of PVMs yet exist (Roberts et al., 2018), although individual papers have discussed strengths and weaknesses of different modeling approaches (Liu et al., 2022; Theuerkauf et al., 2012). Initial efforts are underway to compare land cover reconstructions from REVEALS-GMRF to those from the Bayesian spatio-temporal model STEPPS. Moreover, given that ensemble-based approaches have been shown to have higher predictive ability in fields such as climate modeling and species distribution modeling (Deser et al., 2020; Bothe et al., 2013; Rangel et al., 2009; Thuiller et al., 2009), there is value in developing approaches for ensemble-based PVMs for past land cover inference.

## 5. Conclusions

Continental-scale changes in land cover in North America during the Holocene are the outcome of multiple interacting drivers operating over a range of temporal and spatial scales. For much of the Holocene (ca. 8 to 1.5 ka), the spatial configurations of continental-scale fractional forest cover were broadly stable. Major continental-scale trends included early Holocene afforestation, a middle-Holocene peak in the areal proportion of summergreen trees and shrubs, and a last-millennium increase in open land. These trends were ~~driven~~<sup>powered</sup> by taxon-level dynamics that varied [strongly](#) within and among regions. The regional dynamics can be attributed to individualistic postglacial shifts in the range and abundance of plant taxa driven by changing climates and increased land area after deglaciation; abrupt and large species-level events, such as the mid-Holocene collapse of eastern hemlock; and a shift from localized land use during the late Holocene to massive ecosystem transformation following EuroAmerican settlement. This work rejects the ideas of both pre-industrial forest stability and widespread major conversions in land cover type.

This work contributes to the LandCover6k initiative to produce continental- to global-scale reconstructions of Holocene land cover dynamics that are well-constrained by proxy data and are quantified using consistent methods. These REVEALS-GMRF reconstructions [establish realistic benchmark scenarios constraining and assimilating with Earth system model simulations of the](#)

interactions among Holocene climate, land cover, and anthropogenic change. In addition, they can help refine Holocene models of land use and land cover change, provide a foundation for comparisons among PVMs, and indicate priority areas for future new site-level research, and establish realistic benchmark scenarios constraining and assimilating with Earth system model simulations of the interactions among Holocene climate, land cover, and anthropogenic change. Scaling up from these continental to hemispheric and global products will enable further testing of hypotheses about the drivers and feedbacks of Holocene land cover change, both biogeophysical and biogeochemical. These reconstructions also highlight several potentially important but unstudied vegetation-atmosphere feedbacks, including the collapse of eastern hemlock at ca. 5 ka and a positive feedback loop between mid-Holocene peaks in seasonality of insolation, temperature, and vegetation phenology.

Comparison of the REVEALS-GMRF land cover reconstructions with those inferred from alternate PVMs is a research priority. REVEALS-GMRF depends on a suite of input variables that are both uncertain and exhibit spatial and temporal variation not accounted for in the model. Also, the use of a post-hoc Bayesian GMRF interpolation step means that uncertainty of reconstructions associated with these assumptions is not fully characterized.

Finally, this work emphasizes the importance of quantifying ecosystem processes and feedbacks between the land surface and climate system. Holocene land cover changes in North America are non-negligible and there is a pressing need to refine ecosystem models, in particular when those models are coupled within an Earth System model. The availability of well-constrained data products has been a major barrier to this effort. Paleoecology and palynology are reaching a new level of maturity with respect to advances in data discovery and accessibility, as well as development of consistent analytical methods that permit interpretability of large-scale spatio-temporal change. Data assimilation experiments that integrate these reconstructions into Earth System models have the potential to resolve major questions in the field, such as the Holocene Temperature Conundrum and disentangling the effects of climate-vegetation-human interactions upon Holocene vegetation and climate dynamics.

## 6. Code and data availability

Code and data used in this workflow are publicly available at <https://github.com/andydawson/reveals-na>. The REVEALS-GMRF reconstructions are available at [https://github.com/BehnazP/SpatioCompo\\_entireHolocene\\_NA](https://github.com/BehnazP/SpatioCompo_entireHolocene_NA). The description and implementation of the digitization of lake area is available at [https://github.com/NeotomaDB/neotoma\\_lakes](https://github.com/NeotomaDB/neotoma_lakes).

1015 7. Author contributions

1016 AD and JWW co-designed and co-wrote the paper, with AD leading on REVEALS analyses and figure  
1017 generation. MJG helped initiate this effort, led PAGES LandCover6k, and coordinated activities with  
1018 other continental-scale mapping groups. BP and JL assisted with the GMRF modeling and analyses,  
1019 while SG assisted with Neotoma data provision and digitization of lake size. RSA, AB, DF, KG, DG, TL,  
1020 TM, WO, BS, and CW all contributed new records to the North American Pollen Database in Neotoma  
1021 and assisted in data interpretation. All authors contributed to manuscript development and revision.

1022 8. Competing interests

1023 The authors declare that they have no conflict of interest.

1024 9. Acknowledgments

1025 This work is a contribution to the Past Global Change (PAGES) project and its working group  
1026 LandCover6k (<https://pastglobalchanges.org/science/wg/former/landcover6k/intro>), which in turn  
1027 received support from the Swiss National Science Foundation, the Swiss Academy of Sciences, the US  
1028 National Science Foundation, and the Chinese Academy of Sciences. This work is also supported  
1029 through an NSERC Discovery Grant to A. Dawson. M.-J. Gaillard received support from the Swedish  
1030 Strategic Research Area (SRA) MERGE (Modelling the Regional and Global Earth system;  
1031 <http://www.merge.lu.se>). Data were obtained from the Neotoma Paleocology Database  
1032 (<http://www.neotomadb.org>) and its constituent database the North American Pollen Database. The work  
1033 of data contributors, data stewards, and the Neotoma community is gratefully acknowledged. Support for  
1034 Neotoma is from the NSF-Geoinformatics program (1948926). Digitization of basin areas was assisted  
1035 by Jenna Kilsevich, Grace Roper, Claire Rubbelke, Grace Tunski, Mathias Trachsel, and Bailey Zak.

1036 10. References

1037 Abraham, V., Oušková, V., and Kuneš, P.: Present-day vegetation helps quantifying past land cover in  
1038 selected regions of the Czech Republic, PLoS One, 9, e100117,  
1039 <https://doi.org/10.1371/journal.pone.0100117>, 2014.  
1040 Alder, J. R. and Hostetler, S. W.: Global climate simulations at 3000-year intervals for the last 21 000  
1041 years with the GENMOM coupled atmosphere–ocean model, Clim. Past, 11, 449–471,  
1042 <https://doi.org/10.5194/cp-11-449-2015>, 2015.  
1043 Alessandri, A., Catalano, F., De Felice, M., Van den Hurk, B., and Balsamo, G.: Varying snow and  
1044 vegetation signatures of surface-albedo feedback on the Northern Hemisphere land warming, Environ.  
1045 Res. Lett., 16, 034023, <https://doi.org/10.1088/1748-9326/abd65f>, 2021.

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman



1097 Booth, R. K., Brewer, S., Blaauw, M., Minckley, T. A., and Jackson, S. T.: Decomposing the mid-  
1098 Holocene *Tsuga* decline in eastern North America, *Ecology*, 93, 1841–1852, [https://doi.org/10.1890/11-](https://doi.org/10.1890/11-2062.1)  
1099 2062.1, 2012.

1100 Bothe, O., Jungclaus, J. H., Zanchettin, D., and Zorita, E.: Climate of the last millennium: ensemble  
1101 consistency of simulations and reconstructions, *Clim Past*, 9, 1089–1110, [https://doi.org/10.5194/cp-9-](https://doi.org/10.5194/cp-9-1089-2013)  
1102 1089-2013, 2013.

1103 Bradshaw, R. H. W. and Webb, I. T.: Relationships between contemporary pollen and vegetation data  
1104 from Wisconsin and Michigan, USA, *Ecology*, 66, 721–737, <https://doi.org/10.2307/1940533>, 1985.

1105 Braun, K., Bar-Matthews, M., Matthews, A., Ayalon, A., Cowling, R. M., Karkanas, P., Fisher, E. C.,  
1106 Dyez, K., Zilberman, T., and Marean, C. W.: Late Pleistocene records of speleothem stable isotopic  
1107 compositions from Pinnacle Point on the South African south coast, *Quat. Res.*, 91, 265–288,  
1108 <https://doi.org/10.1017/qua.2018.61>, 2019.

1109 Briles, C. E., Whitlock, C., Bartlein, P. J., and Higuera, P.: Regional and local controls on postglacial  
1110 vegetation and fire in the Siskiyou Mountains, northern California, USA, *Palaeogeogr. Palaeoclimatol.*  
1111 *Palaeoecol.*, 265, 159–169, <https://doi.org/10.1016/j.palaeo.2008.05.007>, 2008.

1112 Broström, A., Nielsen, A. B., Gaillard, M.-J., Hjelle, K., Mazier, F., Binney, H., Bunting, J., Fyfe, R.,  
1113 Meltsov, V., and Poska, A.: Pollen productivity estimates of key European plant taxa for quantitative  
1114 reconstruction of past vegetation: a review, *Veg. Hist. Archaeobotany*, 17, 461–478,  
1115 <https://doi.org/10.1007/s00334-008-0148-8>, 2008.

1116 Brovkin, V., Raddatz, T., Reick, C. H., Claussen, M., and Gayler, V.: Global biogeophysical interactions  
1117 between forest and climate, *Geophys. Res. Lett.*, 36, <https://doi.org/10.1029/2009GL037543>, 2009.

1118 Brovkin, V., Lorenz, S., Raddatz, T., Ilyina, T., Stemmler, I., Toohey, M., and Claussen, M.: What was  
1119 the source of the atmospheric CO<sub>2</sub> increase during the Holocene?, *Biogeosciences*, 16, 2543–2555,  
1120 <https://doi.org/10.5194/bg-16-2543-2019>, 2019.

1121 Brugam, R. B. and Swain, P.: Diatom indicators of peatland development at Pogonia Bog Pond,  
1122 Minnesota, USA, *The Holocene*, 10, 453–464, <https://doi.org/10.1191/095968300668251084>, 2000.

1123 Brugger, S. O. and Rhode, D.: Impact of Pleistocene–Holocene climate shifts on vegetation and fire  
1124 dynamics and its implications for Prearchaic humans in the central Great Basin, USA, *J. Quat. Sci.*, 35,  
1125 987–993, <https://doi.org/10.1002/jqs.3248>, 2020.

1126 Calder, W. J. and Shuman, B. N.: Extensive wildfires, climate change, and an abrupt state change in  
1127 subalpine ribbon forests, Colorado, *Ecology*, 98, 2585–2600, <https://doi.org/10.1002/ecy.1959>, 2017.

1128 Chandan, D. and Peltier, W. R.: African Humid Period precipitation sustained by robust vegetation, soil,  
1129 and lake feedbacks, *Geophys. Res. Lett.*, 47, e2020GL088728, <https://doi.org/10.1029/2020GL088728>,  
1130 2020.

1131 Chaput, M. A. and Gajewski, K.: Relative pollen productivity estimates and changes in Holocene  
1132 vegetation cover in the deciduous forest of southeastern Quebec, Canada, *Botany*, 96, 299–317,  
1133 <https://doi.org/10.1139/cjb-2017-0193>, 2018.

1134 Chen, J., Zhang, Q., Huang, W., Lu, Z., Zhang, Z., and Chen, F.: Northwestward shift of the northern  
1135 boundary of the East Asian summer monsoon during the mid-Holocene caused by orbital forcing and  
1136 vegetation feedbacks, *Quat. Sci. Rev.*, 268, 107136, <https://doi.org/10.1016/j.quascirev.2021.107136>,  
1137 2021.

1138 Chen, L. and Dirmeyer, P. A.: Adapting observationally based metrics of biogeophysical feedbacks from  
1139 land cover/land use change to climate modeling, *Environ. Res. Lett.*, 11, 034002,  
1140 <https://doi.org/10.1088/1748-9326/11/3/034002>, 2016.

1141 Chevalier, M., Davis, B. A. S., Heiri, O., Seppä, H., Chase, B. M., Gajewski, K., Lacourse, T., Telford, R.  
1142 J., Finsinger, W., Guiot, J., Kühl, N., Maezumi, S. Y., Tipton, J. R., Carter, V. A., Brussel, T., Phelps, L.  
1143 N., Dawson, A., Zanon, M., Vallé, F., Nolan, C., Mauri, A., de Vernal, A., Izumi, K., Holmström, L.,  
1144 Marsicek, J., Goring, S., Sommer, P. S., Chaput, M., and Kupriyanov, D.: Pollen-based climate  
1145 reconstruction techniques for late Quaternary studies, *Earth-Sci. Rev.*, 210, 103384,  
1146 <https://doi.org/10.1016/j.earscirev.2020.103384>, 2020.

1147 Clark, J. S., Hussey, T., and Royall, P. D.: Presettlement analogs for Quaternary fire regimes in eastern

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman



North America, *J. Paleolimnol.*, 16, 79–96, <https://doi.org/10.1007/BF00173273>, 1996.

Cruz-Silva, E., Harrison, S. P., Marinova, E., and Prentice, I. C.: A new method based on surface-sample pollen data for reconstructing palaeovegetation patterns, *J. Biogeogr.*, 49, 1381–1396, <https://doi.org/10.1111/jbi.14448>, 2022.

Cwynar, L. C. and Spear, R. W.: Paleovegetation and paleoclimatic changes in the Yukon at 6 ka BP, *Géographie Phys. Quat.*, 49, 29–35, <https://doi.org/10.7202/033027ar>, 1995.

Dallmeyer, A., Kleinen, T., Claussen, M., Weitzel, N., Cao, X., and Herzsuh, U.: The deglacial forest conundrum, *Nat. Commun.*, 13, 6035, <https://doi.org/10.1038/s41467-022-33646-6>, 2022.

Dallmeyer, A., Poska, A., Marquer, L., Seim, A., and Gaillard-Lemdhah, M.-J.: The challenge of comparing pollen-based quantitative vegetation reconstructions with outputs from vegetation models—a European perspective, *Clim. Past Discuss.*, 2023, 1–50, <https://doi.org/10.5194/cp-19-1531-2023>, 2023.

Dalton, A. S., Margold, M., Stokes, C. R., Tarasov, L., Dyke, A. S., Adams, R. S., Allard, S., Arends, H. E., Atkinson, N., Ättrig, J. W., Barnett, P. J., Barnett, R. L., Batterson, M., Bernatchez, P., Borns, H. W., Breckenridge, A., Briner, J. P., Brouard, E., Campbell, J. E., Carlson, A. E., Clague, J. J., Curry, B. B., Daigneault, R.-A., Dubé-Loubert, H., Easterbrook, D. J., Franzi, D. A., Friedrich, H. G., Funder, S., Gauthier, M. S., Gowan, A. S., Harris, K. L., Héty, B., Hooyer, T. S., Jennings, C. E., Johnson, M. D., Kehew, A. E., Kelley, S. E., Kerr, D., King, E. L., Kjeldsen, K. K., Knaeble, A. R., Lajeunesse, P., Lakeman, T. R., Lamothe, M., Larson, P., Lavoie, M., Loope, H. M., Lowell, T. V., Lusardi, B. A., Manz, L., McMartin, I., Nixon, F. C., Occhietti, S., Parkhill, M. A., Piper, D. J. W., Pronk, A. G., Richard, P. J. H., Ridge, J. C., Ross, M., Roy, M., Seaman, A., Shaw, J., Stea, R. R., Teller, J. T., Thompson, W. B., Thorleifson, L. H., Utting, D. J., Veillette, J. J., Ward, B. C., Weddle, T. K., and Wright, H. E.: An updated radiocarbon-based ice margin chronology for the last deglaciation of the North American Ice Sheet Complex, *Quat. Sci. Rev.*, 234, 106223, <https://doi.org/10.1016/j.quascirev.2020.106223>, 2020.

Davis, M. B.: Pollen evidence of changing land use around the shores of Lake Washington, Northwest Sci., 47, 133–148, <https://hdl.handle.net/11299/178248>, 1973.

Davis, M. B.: Outbreaks of forest pathogens in Quaternary history, in: Proceedings of the Fourth International Palynological Conference, Lucknow, India, 216–227, <https://hdl.handle.net/11299/178237>, 1981.

Davis, M. B.: Palynology after Y2K—understanding the source area of pollen in sediments, *Annu. Rev. Earth Planet. Sci.*, 28, 1–18, <https://doi.org/10.1146/annurev.earth.28.1.1>, 2000.

Dawson, A., Paciorek, C. J., McLachlan, J. S., Goring, S., Williams, J. W., and Jackson, S. T.: Quantifying pollen-vegetation relationships to reconstruct ancient forests using 19th-century forest composition and pollen data, *Quat. Sci. Rev.*, 137, 156–175, <https://doi.org/10.1016/j.quascirev.2016.01.012>, 2016.

Dawson, A., Cao, X., Chaput, M., Hopla, E., Li, F., Edwards, M., Fyfe, R., Gajewski, K., Goring, S., Herzsuh, U., and others: Finding the magnitude of human-induced Northern Hemisphere land-cover transformation between 6 and 0.2 ka BP, *PAGES Mag.*, 26, 34–35, <https://doi.org/10.22498/pages.26.1.34>, 2018.

Dawson, A., Paciorek, C. J., Goring, S., Jackson, S., McLachlan, J., and Williams, J. W.: Quantifying trends and uncertainty in prehistoric forest composition in the upper Midwestern United States, *Ecology*, doi: 10.1002/ecy.2856, <https://doi.org/10.1016/j.quascirev.2016.01.012>, 2019.

Dean, W. E., Bradbury, J. P., Anderson, R. Y., and Barnosky, C. W.: The variability of Holocene climate change: Evidence from varved lake sediments, *Science*, 226, 1191–1194, <https://doi.org/10.1126/science.226.4679.1191>, 1984.

Delcourt, H. R. and Delcourt, P. A.: Postglacial rise and decline of *Ostrya virginiana* (Mill.) K Koch and *Carpinus caroliniana* Walt in eastern North-America: Predictable responses of forest species to cyclic changes in seasonality of climates, *J. Biogeogr.*, 21, 137–150, <https://doi.org/10.2307/2845468>, 1994.

Delcourt, P. A., Delcourt, H. R., Criddlebaugh, P. A., and Chapman, J.: Holocene ethnobotanical and paleoecological record of human impact on vegetation in the Little Tennessee River Valley, Tennessee, *Quat. Res.*, 25, 330–349, [https://doi.org/10.1016/0033-5894\(86\)90005-0](https://doi.org/10.1016/0033-5894(86)90005-0), 1986.

Deser, C., Lehner, F., Rodgers, K. B., Ault, T., Delworth, T. L., DiNezio, P. N., Fiore, A., Frankignoul,

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman



1250 Segerström, U., von Stedingk, H., and Seppä, H.: Holocene land-cover reconstructions for studies on land  
 1251 cover-climate feedbacks, *Clim. Past*, 6, 483–499, <https://doi.org/10.5194/cp-6-483-2010>, 2010.  
 1252 Gajewski, K.: Environmental history of the northwestern Québec Treeline, *Quat. Sci. Rev.*, 206, 29–43,  
 1253 <https://doi.org/10.1016/j.quascirev.2018.12.025>, 2019.  
 1254 Gajewski, K., Kriesche, B., Chaput, M. A., Kulik, R., and Schmidt, V.: Human–vegetation interactions  
 1255 during the Holocene in North America, *Veg. Hist. Archaeobotany*, 28, 635–647,  
 1256 <https://doi.org/10.1007/s00334-019-00721-w>, 2019.  
 1257 Gajewski, K., Grenier, A., and Payette, S.: Climate, fire and vegetation history at treeline east of Hudson  
 1258 Bay, northern Québec, *Quat. Sci. Rev.*, 254, 106794, <https://doi.org/10.1016/j.quascirev.2021.106794>,  
 1259 2021.  
 1260 Gavin, D. G. and Brubaker, L. B.: Late Pleistocene and Holocene Environmental Change on the Olympic  
 1261 Peninsula, Washington, Springer, <https://doi.org/10.1007/978-3-319-11014-1>, 2014.  
 1262 Gavin, D. G., Brubaker, L. B., and Greenwald, D. N.: Postglacial climate and fire-mediated vegetation  
 1263 change on the western Olympic Peninsula, Washington (USA), *Ecol. Monogr.*, 83, 471–489,  
 1264 <https://doi.org/10.1890/12-1742.1>, 2013.  
 1265 Gavin, D. G., White, A., Sanborn, P. T., and Hebda, R. J.: Deglacial landforms and Holocene vegetation  
 1266 trajectories in the northern interior cedar-hemlock forests of British Columbia, in: *Untangling the*  
 1267 *Quaternary Period—A Legacy of Stephen C. Porter*, vol. 548, edited by: Waitt, R. B., Thackray, G. D.,  
 1268 and Gillespie, A. R., Geological Society of America, 0, [https://doi.org/10.1130/2020.2548\(05\)](https://doi.org/10.1130/2020.2548(05)), 2021.  
 1269 George, A., Widell, S., Goring, S. J., Roth, R. E., and Williams, J. W.: Range Mapper: An adaptable  
 1270 process for making and using interactive, animated web maps of late-Quaternary open paleoecological  
 1271 data, *Open Quat.*, 9, 15, <https://doi.org/10.5334/oq.114>, 2023.  
 1272 Githumbi, E., Fyfe, R., Gaillard, M.-J., Trondman, A.-K., Mazier, F., Nielsen, A.-B., Poska, A., Sugita,  
 1273 S., Woodbridge, J., Azuara, J., and others: European pollen-based REVEALS land-cover reconstructions  
 1274 for the Holocene: methodology, mapping and potentials, *Earth Syst. Sci. Data*, 14, 1581–1619,  
 1275 <https://doi.org/10.5194/essd-14-1581-2022>, 2022a.  
 1276 Githumbi, E., Pirzamanbein, B., Lindström, J., Poska, A., Fyfe, R., Mazier, F., Nielsen, A. B., Sugita, S.,  
 1277 Trondman, A.-K., Woodbridge, J., and others: Pollen-based maps of past regional vegetation cover in  
 1278 Europe over 12 Millennia—evaluation and potential, *Front. Ecol. Evol.*, 10, 795794,  
 1279 <https://doi.org/10.3389/fevo.2022.795794>, 2022b.  
 1280 Giuliano, C. and Lacourse, T.: Holocene fire regimes, fire-related plant functional types, and climate in  
 1281 south-coastal British Columbia forests, *Ecosphere*, 14, e4416, <https://doi.org/10.1002/ecs2.4416>, 2023.  
 1282 Goring, S.: Neotoma Lakes Assignment, 2024.  
 1283 Grimm, E. C., Bradshaw, R. H. W., Giesecke, T., Lézine, A.-M., Takahara, H., and Williams, J. W.:  
 1284 Pollen databases and their application, edited by: Elias, S. A., 2013.  
 1285 Gugger, P. F. and Sugita, S.: Glacial populations and postglacial migration of Douglas-fir based on fossil  
 1286 pollen and macrofossil evidence, *Quat. Sci. Rev.*, 29, 2052–2070,  
 1287 <https://doi.org/10.1016/j.quascirev.2010.04.022>, 2010.  
 1288 Guo, Z., Zhou, X., and Wu, H.: Glacial-interglacial water cycle, global monsoon and atmospheric  
 1289 methane changes, *Clim. Dyn.*, 39, 1073–1092, <https://doi.org/10.1007/s00382-011-1147-5>, 2012.  
 1290 Haas, J. N. and McAndrews, J. H.: The summer drought related hemlock (*Tsuga canadensis*) decline in  
 1291 eastern North America 5,700 to 5,100 years ago, *Symposium on Sustainable Management of Hemlock*  
 1292 *Ecosystems in Eastern North America*, <https://doi.org/10.2737/NE-GTR-267>, 1999.  
 1293 Hansen, B. C. S. and Engstrom, D. R.: Vegetation history of Pleasant Island, southeastern Alaska, since  
 1294 13,000 yr B.P., *Quat. Res.*, 46, 161–175, <https://doi.org/10.1006/qres.1996.0056>, 1996.  
 1295 Harrison, S. P., Gaillard, M. J., Stocker, B. D., Vander Linden, M., Klein Goldewijk, K., Boles, O.,  
 1296 Braconnot, P., Dawson, A., Fluet-Chouinard, E., Kaplan, J. O., Kastner, T., Pausata, F. S. R., Robinson,  
 1297 E., Whitehouse, N. J., Madella, M., and Morrison, K. D.: Development and testing scenarios for  
 1298 implementing land use and land cover changes during the Holocene in Earth system model experiments,  
 1299 *Geosci. Model Dev.*, 13, 805–824, <https://doi.org/10.5194/gmd-13-805-2020>, 2020.  
 1300 Haslett, J., Parnell, A. C., Hinde, and de Andrade Moral: Modelling Excess Zeros in Count Data: A New

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman



1301 Perspective on Modelling Approaches - Haslett - 2022 - International Statistical Review - Wiley Online  
 1302 Library, 90, 216–236, <https://doi.org/10.1111/insr.12479>, 2021.  
 1303 Hayashi, R., Sasaki, N., Takahara, H., Sugita, S., and Saito, H.: Estimation of absolute pollen productivity  
 1304 based on the flower counting approach: A review, *Quat. Int.*, 641, 122–137,  
 1305 <https://doi.org/10.1016/j.quaint.2022.04.015>, 2022.  
 1306 Hellman, S., Gaillard, M.-J., Broström, A., and Sugita, S.: The REVEALS model, a new tool to estimate  
 1307 past regional plant abundance from pollen data in large lakes: validation in southern Sweden, *J. Quat.*  
 1308 *Sci.*, 23, 21–42, <https://doi.org/10.1002/jqs.1126>, 2008a.  
 1309 Hellman, S. E., Gaillard, M., Broström, A., and Sugita, S.: Effects of the sampling design and selection of  
 1310 parameter values on pollen-based quantitative reconstructions of regional vegetation: a case study in  
 1311 southern Sweden using the REVEALS model, *Veg. Hist. Archaeobotany*, 17, 445–459,  
 1312 <https://doi.org/10.1007/s00334-008-0149-7>, 2008b.  
 1313 Herzschuh, U.: Legacy of the Last Glacial on the present-day distribution of deciduous versus evergreen  
 1314 boreal forests, *Glob. Ecol. Biogeogr.*, 29, 198–206, <https://doi.org/10.1111/geb.13018>, 2020.  
 1315 Herzschuh, U., Böhmer, T., Li, C., Chevalier, M., Hébert, R., Dallmeyer, A., Cao, X., Bigelow, N. H.,  
 1316 Nazarova, L., Novenko, E. Y., Park, J., Peyron, O., Rudaya, N. A., Schlütz, F., Shumilovskikh, L. S.,  
 1317 Tarasov, P. E., Wang, Y., Wen, R., Xu, Q., and Zheng, Z.: LegacyClimate 1.0: a dataset of pollen-based  
 1318 climate reconstructions from 2594 Northern Hemisphere sites covering the last 30 kyr and beyond, *Earth*  
 1319 *Syst Sci Data*, 15, 2235–2258, <https://doi.org/10.5194/essd-15-2235-2023>, 2023.  
 1320 Hoevers, R., Broothaerts, N., and Verstraeten, G.: The potential of REVEALS-based vegetation  
 1321 reconstructions using pollen records from alluvial floodplains, *Veg. Hist. Archaeobotany*, 31, 525–540,  
 1322 <https://doi.org/10.1007/s00334-022-00866-1>, 2022.  
 1323 Hogg, D.: High-resolution analysis of snow albedo interactions in the Arctic, MSc Thesis, University of  
 1324 Waterloo, 2022.  
 1325 Hollinger, D. Y., Ollinger, S. V., Richardson, A. D., Meyers, T. P., Dail, D. B., Martin, M. E., Scott, N.  
 1326 A., Arkebauer, T. J., Baldocchi, D. D., and Clark, K. L.: Albedo estimates for land surface models and  
 1327 support for a new paradigm based on foliage nitrogen concentration, *Glob. Change Biol.*, 16, 696–710,  
 1328 <https://doi.org/10.1111/j.1365-2486.2009.02028.x>, 2010.  
 1329 Huybers, P.: Early Pleistocene glacial cycles and the integrated summer insolation forcing, *Science*, 313,  
 1330 508–511, <https://doi.org/10.1126/science.1125249>, 2006.  
 1331 Iglesias, V. and Whitlock, C.: If the trees burn, is the forest lost? Past dynamics in temperate forests help  
 1332 inform management strategies, *Philos. Trans. R. Soc. B*, 375, 20190115,  
 1333 <https://doi.org/10.1098/rstb.2019.0115>, 2020.  
 1334 Iglesias, V., Whitlock, C., Krause, T. R., and Baker, R. G.: Past vegetation dynamics in the Yellowstone  
 1335 region highlight the vulnerability of mountain systems to climate change, *J. Biogeogr.*, 45, 1768–1780,  
 1336 <https://doi.org/10.1111/jbi.13364>, 2018.  
 1337 Indermühle, A., Stocker, T. F., Joos, F., Fischer, H., Smith, H. J., Wahlen, M., Deck, B., Mastroianni, D.,  
 1338 Tschumi, J., Blunier, T., Meyer, R., and Stauffer, B.: Holocene carbon-cycle dynamics based on CO<sub>2</sub>  
 1339 trapped in ice at Taylor Dome, Antarctica, *Nature*, 398, 121–126, <https://doi.org/10.1038/18158>, 1999.  
 1340 Ireland, A. W. and Booth, R. K.: Hydroclimatic variability drives episodic expansion of a floating peat  
 1341 mat in a North American kettlehole basin, *Ecology*, 92, 11–18, <https://doi.org/10.1890/10-0770.1>, 2010.  
 1342 Islebe, G. A., Hooghiemstra, H., Brenner, M., Curtis, J. H., and Hodell, D. A.: A Holocene vegetation  
 1343 history from lowland Guatemala, *The Holocene*, 6, 265–271,  
 1344 <https://doi.org/10.1177/095968369600600302>, 1996.  
 1345 Jackson, S. T.: Pollen source area and representation in small lakes of the northeastern United States, *Rev.*  
 1346 *Palaeobot. Palynol.*, 63, 53–76, [https://doi.org/10.1016/0034-6667\(90\)90006-5](https://doi.org/10.1016/0034-6667(90)90006-5), 1990.  
 1347 Jackson, S. T. and Lyford, M. E.: Pollen dispersal models in Quaternary plant ecology: Assumptions,  
 1348 parameters, and prescriptions, *Bot. Rev.*, 65, 39–75, <https://doi.org/10.1007/BF02856557>, 1999.  
 1349 Jackson, S. T. and Whitehead, D. R.: Holocene vegetation patterns in the Adirondack Mountains,  
 1350 *Ecology*, 72, 641–654, <https://doi.org/10.2307/2937204>, 1991.  
 1351 Jackson, S. T., Overpeck, J. T., Webb, I. T., Keatts, S. E., and Anderson, K. H.: Mapped plant-

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

1352 macrofossil and pollen records of late Quaternary vegetation change in eastern North America, *Quat. Sci.*  
1353 *Rev.*, 16, 1–70, [https://doi.org/10.1016/S0277-3791\(96\)00047-9](https://doi.org/10.1016/S0277-3791(96)00047-9), 1997.  
1354 Jackson, S. T., Betancourt, J. L., Booth, R. K., and Gray, S. T.: Ecology and the ratchet of events: Climate  
1355 variability, niche dimensions, and species distributions, *Proc. Natl. Acad. Sci.*, 106, 19685–19692,  
1356 <https://doi.org/10.1073/pnas.0901644106>, 2009.  
1357 Jensen, A. M., Fastovich, D., Watson, B. I., Gill, J. L., Jackson, S. T., Russell, J. M., Bevington, J.,  
1358 Hayes, K., Lininger, K., Rubbelke, C., Schellinger, G. C., and Williams, J. W.: More than one way to kill  
1359 a spruce forest: The role of fire and climate in the late-glacial termination of spruce woodlands across the  
1360 southern Great Lakes, *J. Ecol.*, 109, 459–477, <https://doi.org/10.1111/1365-2745.13517>, 2021.  
1361 Kaplan, J. O., Krumhardt, K. M., and Zimmermann, N.: The prehistoric and preindustrial deforestation of  
1362 Europe, *Quat. Sci. Rev.*, 28, 3016–3034, <https://doi.org/10.1016/J.Quascirev.2009.09.028>, 2009.  
1363 Kaplan, J. O., Krumhardt, K. M., Gaillard, M.-J., Sugita, S., Trondman, A.-K., Fyfe, R., Marquer, L.,  
1364 Mazier, F., and Nielsen, A. B.: Constraining the deforestation history of Europe: Evaluation of historical  
1365 land use scenarios with pollen-based land cover reconstructions, *Land*, 6, 91,  
1366 <https://doi.org/10.3390/land6040091>, 2017.  
1367 Kaufman, D. S. and Broadman, E.: Revisiting the Holocene global temperature conundrum, *Nature*, 614,  
1368 425–435, <https://doi.org/10.1038/s41586-022-05536-w>, 2023.  
1369 Kelly, R., Chipman, M. L., Higuera, P. E., Stefanova, I., Brubaker, L. B., and Hu, F. S.: Recent burning  
1370 of boreal forests exceeds fire regime limits of the past 10,000 years, *Proc. Natl. Acad. Sci.*, 110, 13055–  
1371 13060, <https://doi.org/10.1073/pnas.1305069110>, 2013.  
1372 Klein Goldewijk, K., Beusen, A., and Janssen, P.: Long-term dynamic modeling of global population and  
1373 built-up area in a spatially explicit way: HYDE 3.1, *The Holocene*, 20, 565–573,  
1374 <https://doi.org/10.1177/0959683609356587>, 2010.  
1375 Klein Goldewijk, K., Beusen, A., van Drecht, G., and de Vos, M.: The HYDE 3.1 spatially explicit  
1376 database of human-induced global land-use change over the past 12,000 years, *Glob. Ecol. Biogeogr.*, 20,  
1377 73–86, <https://doi.org/10.1111/j.1466-8238.2010.00587.x>, 2011.  
1378 Knight, C. A., Anderson, L., Bunting, M. J., Champagne, M., Clayburn, R. M., Crawford, J. N.,  
1379 Klimaszewski-Patterson, A., Knapp, E. E., Lake, F. K., Mensing, S. A., Wahl, D., Wanket, J., Watts-  
1380 Tobin, A., Potts, M. D., and Battles, J. J.: Land management explains major trends in forest structure and  
1381 composition over the last millennium in California's Klamath Mountains, *Proc. Natl. Acad. Sci.*, 119,  
1382 e2116264119, <https://doi.org/10.1073/pnas.2116264119>, 2022.  
1383 Komatsu, H., Tanaka, N., and Kume, T.: Do coniferous forests evaporate more water than broad-leaved  
1384 forests in Japan?, *J. Hydrol.*, 336, 361–375, <https://doi.org/10.1016/j.jhydrol.2007.01.009>, 2007.  
1385 Kuparinen, A., Markkanen, T., Riikonen, H., and Vesala, T.: Modeling air-mediated dispersal of spores,  
1386 pollen and seeds in forested areas, *Ecol. Model.*, 208, 177–188,  
1387 <https://doi.org/10.1016/j.ecolmodel.2007.05.023>, 2007.  
1388 Kutzbach, Bartlein, P. J., Foley, J. A., Harrison, S. P., Hostetler, S. W., Liu, Z., Prentice, I. C., and Webb  
1389 III, T.: Potential role of vegetation feedback in the climate sensitivity of high-latitude regions: A case  
1390 study at 6000 years B.P., *Glob. Biogeochem. Cycles*, 10, 727–736, <https://doi.org/10.1029/96GB02690>,  
1391 1996.  
1392 Lacourse, T.: Environmental change controls postglacial forest dynamics through interspecific differences  
1393 in life-history traits, *Ecology*, 90, 2149–2160, <https://doi.org/10.1890/08-1136.1>, 2009.  
1394 Lacourse, T. and Adeleye, M. A.: Climate and species traits drive changes in Holocene forest composition  
1395 along an elevation gradient in Pacific Canada, *Front. Ecol. Evol.*, 10,  
1396 <https://doi.org/10.3389/fevo.2022.838545>, 2022.  
1397 Lacourse, T. and Gajewski, K.: Current practices in building and reporting age-depth models, *Quat. Res.*,  
1398 96, 28–38, <https://doi.org/10.1017/qua.2020.47>, 2020.  
1399 Lacourse, T., Mathewes, R. W., and Hebda, R. J.: Paleoecological analyses of lake sediments reveal  
1400 prehistoric human impact on forests at Anthony Island UNESCO World Heritage Site, Queen Charlotte  
1401 Islands (Haida Gwaii), Canada, *Quat. Res.*, 68, 177–183, <http://dx.doi.org/10.1016/j.yqres.2007.04.005>,  
1402 2007.

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

1403 Lacourse, T., Delepine, J. M., Hoffman, E. H., and Mathewes, R. W.: A 14,000 year vegetation history of  
 1404 a hypermaritime island on the outer Pacific coast of Canada based on fossil pollen, spores and conifer  
 1405 stomata, *Quat. Res.*, 78, 572–582, <http://dx.doi.org/10.1016/j.yqres.2012.08.008>, 2012.

1406 Leopold, E. B. and Boyd, R.: An ecological history of old prairie areas, southwestern Washington, in:  
 1407 *Indians, Fire and the Land*, edited by: Boyd, R., Oregon State University Press,  
 1408 <https://dx.doi.org/10.1353/book94487>, 139–166, 1999.

1409 Lewis, S. L. and Maslin, M. A.: Defining the Anthropocene, *Nature*, 519, 171–180,  
 1410 <https://doi.org/10.1038/nature14258>, 2015.

1411 Li, F., Gaillard, M.-J., Cao, X., Herzschuh, U., Sugita, S., Tarasov, P. E., Wagner, M., Xu, Q., Ni, J., and  
 1412 Wang, W.: Towards quantification of Holocene anthropogenic land-cover change in temperate China: A  
 1413 review in the light of pollen-based REVEALS reconstructions of regional plant cover, *Earth-Sci. Rev.*,  
 1414 203, 103119, <https://doi.org/10.1016/j.earscirev.2020.103119>, 2020.

1415 Li, F., Gaillard, M.-J., Cao, X., Herzschuh, U., Sugita, S., Ni, J., Zhao, Y., An, C., Huang, X., Li, Y., Liu,  
 1416 H., Sun, A., and Yao, Y.: Gridded pollen-based Holocene regional plant cover in temperate and northern  
 1417 subtropical China suitable for climate modelling, *Earth Syst Sci Data*, 15, 95–112,  
 1418 <https://doi.org/10.5194/essd-15-95-2023>, 2023.

1419 Liefert, D. T. and Shuman, B. N.: Pervasive desiccation of North American lakes during the late  
 1420 Quaternary, *Geophys. Res. Lett.*, 47, e2019GL086412, <https://doi.org/10.1029/2019GL086412>, 2020.

1421 Lindgren, F., Rue, H., and Lindström, J.: An explicit link between Gaussian fields and Gaussian Markov  
 1422 random fields: the stochastic partial differential equation approach, *J. R. Stat. Soc. Ser. B Stat. Methodol.*,  
 1423 73, 423–498, <https://doi.org/10.1111/j.1467-9868.2011.00777.x>, 2011.

1424 Liu, J., Chen, J. M., and Cihlar, J.: Mapping evapotranspiration based on remote sensing: An application  
 1425 to Canada's landmass, *Water Resour. Res.*, 39, <https://doi.org/10.1029/2002WR001680>, 2003.

1426 Liu, Y., Ogle, K., Lichstein, J. W., and Jackson, S. T.: Estimation of pollen productivity and dispersal:  
 1427 How pollen assemblages in small lakes represent vegetation, *Ecol. Monogr.*, 92, e1513,  
 1428 <https://doi.org/10.1002/ecm.1513>, 2022.

1429 Liu, Z., Zhu, J., Rosenthal, Y., Zhang, X., Otto-Bliesner, B. L., Timmermann, A., Smith, R. S., Lohmann,  
 1430 G., Zheng, W., and Elison Timm, O.: The Holocene temperature conundrum, *Proc. Natl. Acad. Sci.*, 111,  
 1431 E3501–E3505, <https://doi.org/10.1073/pnas.1407229111>, 2014.

1432 Loisel, J., van Bellen, S., Pelletier, L., Talbot, J., Hugelius, G., Karan, D., Yu, Z., Nichols, J., and  
 1433 Holmquist, J.: Insights and issues with estimating northern peatland carbon stocks and fluxes since the  
 1434 Last Glacial Maximum, *Earth-Sci. Rev.*, 165, 59–80, <https://doi.org/10.1016/j.earscirev.2016.12.001>,  
 1435 2017.

1436 Long, C. J., Whitlock, C., Bartlein, P. J., and Millsaugh, S. H.: A 9000-year fire history from the Oregon  
 1437 Coast Range, based on a high-resolution charcoal study, *Can. J. For. Res.*, 28, 774–787,  
 1438 <https://doi.org/10.1139/x98-051>, 1998.

1439 Long, C. J., Whitlock, C., and Bartlein, P. J.: Holocene vegetation and fire history of the Coast Range,  
 1440 western Oregon, USA, *The Holocene*, 17, 917–926, <https://doi.org/10.1177/0959683607082408>, 2007.

1441 Lorant, M. M., Berner, L. T., Goetz, S. J., Jin, Y., and Randerson, J. T.: Vegetation controls on northern  
 1442 high latitude snow- albedo feedback: observations and CMIP 5 model simulations, *Glob. Change Biol.*,  
 1443 20, 594–606, <https://doi.org/10.1111/gcb.12391>, 2014.

1444 MacDonald, G. M.: Postglacial palaeoecology of the subalpine forest-grassland ecotone of southwestern  
 1445 Alberta: new insights on vegetation and climate change in the Canadian Rocky Mountains and adjacent  
 1446 foothills, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 73, 155–173, [https://doi.org/10.1016/0031-0182\(89\)90001-1](https://doi.org/10.1016/0031-0182(89)90001-1), 1989.

1447 MacDonald, G. M. and Cwynar, L. C.: Post-glacial population growth rates of *Pinus contorta* ssp. *latifolia*  
 1448 in western Canada, *J. Ecol.*, 79, 417–429, <https://doi.org/10.2307/2260723>, 1991.

1449 Marcott, S. A., Shakun, J. D., Clark, P. U., and Mix, A. C.: A reconstruction of regional and global  
 1450 temperature for the past 11,300 years, *Science*, 339, 1198–1201, <https://doi.org/10.1126/science.1228026>, 2013.

1451 Marlon, J. R., Bartlein, P. J., Gavin, D. G., Long, C. J., Anderson, R. S., Briles, C. E., Brown, K. J.,

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Colombaroli, D., Hallett, D. J., and Power, M. J.: Long-term perspective on wildfires in the western USA, Proc. Natl. Acad. Sci., 109, E535–E543, <https://doi.org/10.1073/pnas.1112839109>, 2012.

Marsicek, J., Shuman, B. N., Bartlein, P. J., Shafer, S. L., and Brewer, S.: Reconciling divergent trends and millennial variations in Holocene temperatures, Nature, 554, 92–96, <https://doi.org/10.1038/nature25464>, 2018.

McAndrews, J. H. and Turton, C. L.: Canada geese dispersed cultigen pollen grains from prehistoric Iroquoian fields to Crawford Lake, Ontario, Canada, Palynology, 31, 9–18, <https://doi.org/10.2113/gspalynol.31.1.9>, 2007.

McKay, N. P., Emile-Geay, J., and Khider, D.: geoChronR – an R package to model, analyze, and visualize age-uncertain data, Geochronology, 3, 149–169, <https://doi.org/10.5194/gchron-3-149-2021>, 2021.

McMichael, C. N. H.: Ecological legacies of past human activities in Amazonian forests, New Phytol., 229, <https://doi.org/10.1111/nph.16888>, 2020.

Mensing, S., Smith, J., Burkle Norman, K., and Allan, M.: Extended drought in the Great Basin of western North America in the last two millennia reconstructed from pollen records, 22nd Pac. Clim. Workshop, 188, 79–89, <https://doi.org/10.1016/j.quaint.2007.06.009>, 2008.

Minckley, T. A., Whitlock, C., and Bartlein, P. J.: Vegetation, fire, and climate history of the northwestern Great Basin during the last 14,000 years, Quat. Sci. Rev., 26, 2167–2184, <https://doi.org/10.1016/j.quascirev.2007.04.009>, 2007.

Mottl, O., Flantua, S. G. A., Bhatta, K. P., Felde, V. A., Giesecke, T., Goring, S., Grimm, E. C., Haberle, S., Hooghiemstra, H., Ivory, S., Kuneš, P., Wolters, S., Seddon, A. W. R., and Williams, J. W.: Global acceleration in rates of vegetation change over the last 18,000 years, Science, 321, 860–864, <https://doi.org/10.1126/science.abg1685>, 2021.

Munoz, S. E. and Gajewski, K.: Distinguishing prehistoric human influence on late-Holocene forests in southern Ontario, Canada, The Holocene, 20, 967–981, <https://doi.org/10.1177/0959683610362815>, 2010.

Munoz, S. E., Schroeder, S., Fike, D. A., and Williams, J. W.: A record of sustained prehistoric and historic land use from the Cahokia region, Illinois, USA, Geology, 42, 499–502, <https://doi.org/10.1130/G35541.1>, 2014.

Napier, J. D. and Chipman, M. L.: Emerging palaeoecological frameworks for elucidating plant dynamics in response to fire and other disturbance, Glob. Ecol. Biogeogr., n/a, <https://doi.org/10.1111/geb.13416>, 2021.

Natural Resources Canada: National Hydro Network, 2022.

Nelson, D. M., Hu, F. S., Grimm, E. C., Curry, B. B., and Slate, J. E.: The influence of aridity and fire on Holocene prairie communities in the eastern Prairie Peninsula, Ecology, 87, 2523–2536, [https://doi.org/10.1890/0012-9658\(2006\)87\[2523:TIOAAF\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[2523:TIOAAF]2.0.CO;2), 2006.

Niklas, K. J.: The motion of windborne pollen grains around conifer ovulate cones: implications on wind pollination, Am. J. Bot., 71, 356–374, <https://doi.org/10.1002/j.1537-2197.1984.tb12523.x>, 1984.

Nordt, L., von Fischer, J., and Tieszen, L.: Late Quaternary temperature record from buried soils of the North American Great Plains, Geology, 35, 159–162, <https://doi.org/10.1130/G23345A.1>, 2007.

Oswald, W. W. and Foster, D. R.: Middle-Holocene dynamics of *Tsuga canadensis* (eastern hemlock) in northern New England, USA, The Holocene, 22, 71–78, <https://doi.org/10.1177/0959683611409774>, 2012.

Oswald, W. W., Doughty, E. D., Foster, D. R., Shuman, B. N., and Wagner, D. L.: Evaluating the role of insects in the middle-Holocene *Tsuga* decline, J. Torrey Bot. Soc., 144, 35–39, <https://doi.org/10.3159/TORREY-D-15-00074.1>, 2017.

Oswald, W. W., Foster, D., Shuman, B., Doughty, E., Faison, E. K., Hall, B., Hansen, B. C. S., Lindblad, M., Marroquin, A., and Truebe, S.: Subregional variability in the response of New England vegetation to postglacial climate change, J. Biogeogr., 45, 2375–2388, <https://doi.org/10.1111/jbi.13407>, 2018.

Oswald, W. W., Foster, D. R., Shuman, B. N., Chilton, E. S., Doucette, D. L., and Duranleau, D. L.:

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman



Conservation implications of limited Native American impacts in pre-contact New England, *Nat. Sustain.*,  
 3, 241–246, <https://doi.org/10.1038/s41893-019-0466-0>, 2020.  
 Otto, J., Raddatz, T., and Claussen, M.: Strength of forest-albedo feedback in mid-Holocene climate  
 simulations, *Clim. Past*, 7, 1027–1039, <https://doi.org/10.5194/cp-7-1027-2011>, 2011.  
 Paciorek, C. J., Goring, S. J., Thurman, A., Cogbill, C. V., Williams, J. W., Mladenoff, D. J., Peters, J. A.,  
 Zhu, J., and McLachlan, J. S.: Statistically-estimated tree composition for the northeastern United States  
 at the time of Euro-American settlement, *PLoS One*, 11, e0150087,  
<http://dx.doi.org/10.1371/journal.pone.0150087>, 2016.  
 Palsea and Group), (PALeo SEA level working: The sea-level conundrum: case studies from palaeo-  
 archives, *J. Quat. Sci.*, 25, 19–25, <https://doi.org/10.1002/jqs.1270>, 2010.  
 Parnell, A. C., Haslett, J., Allen, J. R., Buck, C. E., and Huntley, B.: A flexible approach to assessing  
 synchronicity of past events using Bayesian reconstructions of sedimentation history, *Quat. Sci. Rev.*, 27,  
 1872–1885, <https://doi.org/10.1016/j.quascirev.2008.07.009>, 2008.  
 Payette, S.: A paleo-perspective on ecosystem collapse in boreal North America, edited by: Canadell, J.  
 G. and Jackson, R. B., *Ecosyst. Collapse Clim. Change*, 101–129, [https://doi.org/10.1007/978-3-030-71330-0\\_5](https://doi.org/10.1007/978-3-030-71330-0_5), 2021.  
 Payette, S., Filion, L., and Delwaide, A.: Spatially explicit fire-climate history of the boreal forest-tundra  
 (eastern Canada) over the last 2000 years, *Philos. Trans. Biol. Sci.*, 363, 2301–2316,  
<https://doi.org/10.1098/rstb.2007.2201>, 2008.  
 Payette, S., Couillard, P.-L., Frégeau, M., Laflamme, J., and Lavoie, M.: The velocity of postglacial  
 migration of fire-adapted boreal tree species in eastern North America, *Proc. Natl. Acad. Sci.*, 119,  
 e2210496119, <https://doi.org/10.1073/pnas.2210496119>, 2022.  
 Peltier, W. R.: Ice Age Paleotopography, *Science*, 265, 195–201,  
<https://doi.org/10.1126/science.265.5169.195>, 1994.  
 Peros, M. C., Gajewski, K., and Viau, A. E.: Continental-scale tree population response to rapid climate  
 change, competition and disturbance, *Glob. Ecol. Biogeogr.*, 17, 658–669, <https://doi.org/10.1111/j.1466-8238.2008.00406.x>, 2008.  
 Peros, M. C., Munoz, S. E., Gajewski, K., and Viau, A. E.: Prehistoric demography of North America  
 inferred from radiocarbon data, *J. Archaeol. Sci.*, 37, 656–664, <https://doi.org/10.1016/j.jas.2009.10.029>,  
 2010.  
 Perrotti, A. G., Ramiadantsoa, T., O’Keefe, J., and Nuñez Otaño, N.: Uncertainty in coprophilous fungal  
 spore concentration estimates, *Front. Ecol. Evol.*, 10, <https://doi.org/10.3389/fevo.2022.1086109>, 2022.  
 Pirzamanbein, B., Lindström, J., Poska, A., Sugita, S., Trondman, A.-K., Fyfe, R., Mazier, F., Nielsen, A.  
 B., Kaplan, J. O., Bjune, A., Birks, H. J. B., Giesecke, T., Kangur, M., Latalowa, M., Marquer, L., Smith,  
 B., and Gaillard, M.-J.: Creating spatially continuous maps of past land cover from point estimates: A  
 new statistical approach applied to pollen data, *Ecol. Complex.*, 20, 127–141,  
<https://doi.org/10.1016/j.ecocom.2014.09.005>, 2014.  
 Pirzamanbein, B., Lindström, J., Poska, A., and Gaillard, M.-J.: Modelling spatial compositional data:  
 Reconstructions of past land cover and uncertainties, *Spat. Stat.*, 24, 14–31,  
<https://doi.org/10.1016/j.spasta.2018.03.005>, 2018.  
 Pirzamanbein, B., Poska, A., and Lindström, J.: Bayesian reconstruction of past land cover from pollen  
 data: Model robustness and sensitivity to auxiliary variables, *Earth Space Sci.*, 7,  
<https://doi.org/10.1029/2018EA000547>, 2020.  
 Pongratz, J., Reick, C. H., Raddatz, T., and Claussen, M.: Biogeophysical versus biogeochemical climate  
 response to historical anthropogenic land cover change, *Geophys. Res. Lett.*, 37, L08702,  
<https://doi.org/10.1029/2010GL043010>, 2010.  
 Prentice, I. C.: Pollen representation, source area, and basin size: toward a unified theory of pollen  
 analysis, *Quat. Res.*, 23, 76–86, [https://doi.org/10.1016/0033-5894\(85\)90073-0](https://doi.org/10.1016/0033-5894(85)90073-0), 1985.  
 Prentice, I. C., Jolly, D., and BIOME 6000 participants: Mid-Holocene and glacial-maximum vegetation  
 geography of the northern continents and Africa, *J. Biogeogr.*, 27, 507–519,  
<https://doi.org/10.1046/j.1365-2699.2000.00425.x>, 2000.

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

Formatted: Font: (Default) TimesNew Roman

1556 Prentice, I. C., Harrison, S. P., and Bartlein, P. J.: Global vegetation and terrestrial carbon cycle changes  
 1557 after the last ice age, *New Phytol.*, 189, 988–998, <https://doi.org/10.1111/j.1469-8137.2010.03620.x>,  
 1558 2011.

1559 Raiho, A., Paciorek, C. J., Dawson, A., Jackson, S. T., Mladenoff, D. J., Williams, J. W., and McLachlan,  
 1560 J. S.: 8000-year doubling of Midwestern forest biomass driven by population- and biome-scale processes,  
 1561 *Science*, 376, 1491–1495, <https://doi.org/10.1126/science.abk3126>, 2022.

1562 Rangel, T. F. L. V. B., Diniz-Filho, J. A. F., and Araújo, M. B.: BIOENSEMBLES 1.0, Software for  
 1563 computer intensive ensemble forecasting of species distributions under climate change, 2009.

1564 Reimer, P. J., Austin, W. E., Bard, E., Bayliss, A., Blackwell, P. G., Ramsey, C. B., Butzin, M., Cheng,  
 1565 H., Edwards, R. L., and Friedrich, M.: The IntCal20 Northern Hemisphere radiocarbon age calibration  
 1566 curve (0–55 cal kBP), *Radiocarbon*, 62, 725–757, <https://doi.org/10.1017/RDC.2020.41>, 2020.

1567 Richard, P., Fréchette, B., Grondin, P., and Lavoie, M.: Histoire postglaciaire de la végétation de la forêt  
 1568 boréale du Québec et du Labrador, *Nat. Can.*, 144, 63–76, <https://doi.org/10.7202/1070086ar>, 2020.

1569 Roberts, N., Fyfe, R. M., Woodbridge, J., Gaillard, M. J., Davis, B. A. S., Kaplan, J. O., Marquer, L.,  
 1570 Mazier, F., Nielsen, A. B., Sugita, S., Trondman, A. K., and Leydet, M.: Europe’s lost forests: a pollen-  
 1571 based synthesis for the last 11,000 years, *Sci. Rep.*, 8, 716, <https://doi.org/10.1038/s41598-017-18646-7>,  
 1572 2018.

1573 Rogelj, J., Popp, A., Calvin, K. V., Luderer, G., Emmerling, J., Gernaat, D., Fujimori, S., Strefler, J.,  
 1574 Hasegawa, T., Marangoni, G., Krey, V., Kriegler, E., Riahi, K., van Vuuren, D. P., Doelman, J., Drouet,  
 1575 L., Edmonds, J., Fricko, O., Harmsen, M., Havlik, P., Humpenöder, F., Stehfest, E., and Tavoni, M.:  
 1576 Scenarios towards limiting global mean temperature increase below 1.5 °C, *Nat. Clim. Change*, 8, 325–  
 1577 332, <https://doi.org/10.1038/s41558-018-0091-3>, 2018.

1578 Roos, C. I.: Scale in the study of Indigenous burning, *Nat. Sustain.*, <https://doi.org/10.1038/s41893-020-0579-5>, 2020.

1579 Roos, C. I., Zedeño, M. N., Hollenback, K. L., and Erlick, M. M. H.: Indigenous impacts on North  
 1580 American Great Plains fire regimes of the past millennium, *Proc. Natl. Acad. Sci.*, 115, 8143–8148,  
 1581 <https://doi.org/10.1073/pnas.1805259115>, 2018.

1582 Rosenberg, S. M., Walker, I. R., and Mathewes, R. W.: Postglacial spread of hemlock (*Tsuga*) and  
 1583 vegetation history in Mount Revelstoke National Park, British Columbia, Canada, *Can. J. Bot.*, 81, 139–  
 1584 151, <https://doi.org/10.1139/b03-015>, 2003.

1585 Ruddiman, W. F.: The anthropogenic greenhouse era began thousands of years ago, *Clim. Change*, 61,  
 1586 261–293, <https://doi.org/10.1023/B:CLIM.0000004577.17928.f>, 2003.

1587 Ruddiman, W. F.: The Anthropocene, *Annu. Rev. Earth Planet. Sci.*, 41, 4.1–4.24,  
 1588 <https://doi.org/10.1146/annurev-earth-050212-123944>, 2013.

1589 Serge, M.-A., Mazier, F., Fyfe, R., Gaillard, M.-J., Klein, T., Lagnoux, A., Galop, D., Githumbi, E.,  
 1590 Mindrescu, M., and Nielsen, A. B.: Testing the effect of relative pollen productivity on the REVEALS  
 1591 model: a validated reconstruction of Europe-wide holocene vegetation, *Land*, 12, 986,  
 1592 <https://doi.org/10.3390/land12050986>, 2023.

1593 Shuman, B., Webb, T., III, Bartlein, P., and Williams, J. W.: The anatomy of a climatic oscillation:  
 1594 vegetation change in eastern North America during the Younger Dryas chronozone, *Quat. Sci. Rev.*, 21,  
 1595 1777–1791, [https://doi.org/10.1016/S0277-3791\(02\)00030-6](https://doi.org/10.1016/S0277-3791(02)00030-6), 2002.

1596 Shuman, B. N. and Marsicek, J.: The structure of Holocene climate change in mid-latitude North  
 1597 America, *Quat. Sci. Rev.*, 141, 38–51, <https://doi.org/10.1016/j.quascirev.2016.03.009>, 2016.

1598 Shuman, B. N., Newby, P. C., Huang, Y., and Webb, I. T.: Evidence for the close climatic control of  
 1599 New England vegetation history, *Ecology*, 85, 1297–1310, <https://doi.org/10.1890/02-0286>, 2004.

1600 Shuman, B. N., Marsicek, J., Oswald, W. W., and Foster, D. R.: Predictable hydrological and ecological  
 1601 responses to Holocene North Atlantic variability, *Proc. Natl. Acad. Sci.*, 116, 5985–5990,  
 1602 <https://doi.org/10.1073/pnas.1814307116>, 2019.

1603 Shuman, B. N., Stefanescu, I. C., Grigg, L. D., Foster, D. R., and Oswald, W. W.: A millennia-scale  
 1604 oscillation in latitudinal temperature gradients along the western North Atlantic during the mid-Holocene,  
 1605 *Geophys. Res. Lett.*, 50, e2022GL102556, <https://doi.org/10.1029/2022GL102556>, 2023.

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

1607 Sjögren, P., van der Knaap, W. O., Huusko, A., and van Leeuwen, J. F. N.: Pollen productivity, dispersal,  
1608 and correction factors for major tree taxa in the Swiss Alps based on pollen-trap results, *Rev. Palaeobot.*  
1609 *Palynol.*, 152, 200–210, <https://doi.org/10.1016/j.revpalbo.2008.05.003>, 2008.  
1610 Snitker, G., Roos, C. I., Sullivan, A. P., Maezumi, S. Y., Bird, D. W., Coughlan, M. R., Derr, K. M.,  
1611 Gassaway, L., Klimaszewski-Patterson, A., and Loehman, R. A.: A collaborative agenda for archaeology  
1612 and fire science, *Nat. Ecol. Evol.*, <https://doi.org/10.1038/s41559-022-01759-2>, 2022.  
1613 Spear, R. W., Davis, M. B., and Shane, L. C. K.: Late Quaternary history of low- and mid-elevation  
1614 vegetation in the White Mountains of New Hampshire, *Ecol. Monogr.*, 64, 85–109,  
1615 <https://doi.org/10.2307/2937056>, 1994.  
1616 Stegner, M. A. and Spanbauer, T. L.: North American pollen records provide evidence for macroscale  
1617 ecological changes in the Anthropocene, *Proc. Natl. Acad. Sci.*, 120, e2306815120,  
1618 <https://doi.org/10.1073/pnas.2306815120>, 2023.  
1619 Stephens, L., Fuller, D., Boivin, N., Rick, T., Gauthier, N., Kay, A., Marwick, B., Armstrong, C. G.,  
1620 Barton, C. M., Denham, T., Douglass, K., Driver, J., Janz, L., Roberts, P., Rogers, J. D., Thakar, H.,  
1621 Altaweel, M., Johnson, A. L., Sampietro Vattuone, M. M., Aldenderfer, M., Archila, S., Artioli, G., Bale,  
1622 M. T., Beach, T., Borrell, F., Braje, T., Buckland, P. I., Jiménez Cano, N. G., Capriles, J. M., Diez  
1623 Castillo, A., Çilingiroğlu, Ç., Negus Cleary, M., Conolly, J., Coutros, P. R., Covey, R. A., Cremaschi, M.,  
1624 Crowther, A., Der, L., di Lernia, S., Doershuk, J. F., Doolittle, W. E., Edwards, K. J., Erlandson, J. M.,  
1625 Evans, D., Fairbairn, A., Faulkner, P., Feinman, G., Fernandes, R., Fitzpatrick, S. M., Fyfe, R., Garcea,  
1626 E., Goldstein, S., Goodman, R. C., Dalpoim Guedes, J., Herrmann, J., Hiscock, P., Hommel, P.,  
1627 Horsburgh, K. A., Hritz, C., Ives, J. W., Junno, A., Kahn, J. G., Kaufman, B., Kearns, C., Kidder, T. R.,  
1628 Lanoë, F., Lawrence, D., Lee, G.-A., Levin, M. J., Lindsoug, H. B., López-Sáez, J. A., Macrae, S.,  
1629 Marchant, R., Marston, J. M., McClure, S., McCoy, M. D., Miller, A. V., Morrison, M., Motuzaite  
1630 Matuzeviciute, G., Müller, J., Nayak, A., Noerwidi, S., Peres, T. M., Peterson, C. E., Proctor, L., Randall,  
1631 A. R., Renette, S., Robbins Schug, G., Ryzewski, K., Saini, R., Scheinsohn, V., Schmidt, P., Sebillaud, P.,  
1632 Seitsonen, O., Simpson, I. A., Soltysiak, A., Speakman, R. J., Spengler, R. N., Steffen, M. L., et al.:  
1633 Archaeological assessment reveals Earth's early transformation through land use, *Science*, 365, 897–902,  
1634 <https://doi.org/10.1126/science.aax1192>, 2019.  
1635 Strandberg, G., Kjellström, E., Poska, A., Wagner, S., Gaillard, M. J., Trondman, A. K., Mauri, A., Davis,  
1636 B. A. S., Kaplan, J. O., Birks, H. J. B., Bjune, A. E., Fyfe, R., Giesecke, T., Kalhina, L., Kangur, M., van  
1637 der Knaap, W. O., Kokfelt, U., Kuneš, P., Lata, I owa, M., Marquer, L., Mazier, F., Nielsen, A. B., Smith,  
1638 B., Seppä, H., and Sugita, S.: Regional climate model simulations for Europe at 6 and 0.2 k BP:  
1639 sensitivity to changes in anthropogenic deforestation, *Clim Past*, 10, 661–680, [https://doi.org/10.5194/cp-](https://doi.org/10.5194/cp-10-661-2014)  
1640 [10-661-2014](https://doi.org/10.5194/cp-10-661-2014), 2014.  
1641 Strandberg, G., Lindström, J., Poska, A., Zhang, Q., Fyfe, R., Githumbi, E., Kjellström, E., Mazier, F.,  
1642 Nielsen, A. B., Sugita, S., Trondman, A.-K., Woodbridge, J., and Gaillard, M.-J.: Mid-Holocene  
1643 European climate revisited: New high-resolution regional climate model simulations using pollen-based  
1644 land-cover, *Quat. Sci. Rev.*, 281, 107431, <https://doi.org/10.1016/j.quascirev.2022.107431>, 2022.  
1645 Sugita, S.: Theory of quantitative reconstruction of vegetation I: pollen from large sites REVEALS  
1646 regional vegetation composition, *The Holocene*, 17, 229–241,  
1647 <https://doi.org/10.1177/0959683607075837>, 2007a.  
1648 Sugita, S.: Theory of quantitative reconstruction of vegetation II: all you need is LOVE, *The Holocene*,  
1649 17, 243–257, <https://doi.org/10.1177/0959683607075838>, 2007b.  
1650 Sugita, S., Parshall, T., Calcote, R., and Walker, K.: Testing the Landscape Reconstruction Algorithm for  
1651 spatially explicit reconstruction of vegetation in northern Michigan and Wisconsin, *Quat. Res.*, 74, 289–  
1652 300, <https://doi.org/10.1016/j.yqres.2010.07.008>, 2010.  
1653 Sutton, O. G.: 1953: *Micrometeorology*. New York: McGraw-Hill, 1953.  
1654 Theuerkauf, M. and Couwenberg: ROPES reveals past land cover and PPEs From single pollen records,  
1655 *Front. Earth Sci.*, 6, <https://doi.org/10.3389/feart.2018.00014>, 2018.  
1656 Theuerkauf, M., Kuparinen, A., and Joosten, H.: Pollen productivity estimates strongly depend on  
1657 assumed pollen dispersal, *The Holocene*, 23, <https://doi.org/10.1177/0959683612450194>, 2012.

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

Formatted: Font: (Default)TimesNew Roman

1658 Theuerkauf, M., Couwenberg, J., Kuparinen, A., and Liebscher, V.: A matter of dispersal: REVEALSinR  
1659 introduces state-of-the-art dispersal models to quantitative vegetation reconstruction, *Veg. Hist.*  
1660 *Archaeobotany*, 25, 541–553, <https://doi.org/10.1007/s00334-016-0572-0>, 2016.

1661 Thompson, A. J., Zhu, J., Poulsen, C. J., Tierney, J. E., and Skinner, C. B.: Northern Hemisphere  
1662 vegetation change drives a Holocene thermal maximum, *Sci. Adv.*, 8, eabj6535,  
1663 <https://doi.org/10.1126/sciadv.abj6535>, 2022.

1664 Thompson, J. R., Carpenter, D. N., Cogbill, C. V., and Foster, D. R.: Four centuries of change in  
1665 northeastern United States forests, *PLoS One*, 8, e72540, <https://doi.org/10.1371/journal.pone.0072540>,  
1666 2013.

1667 Thompson, R. S.: Late Quaternary environments in Ruby Valley, Nevada, *Quat. Res.*, 37, 1–15,  
1668 [https://doi.org/10.1016/0033-5894\(92\)90002-Z](https://doi.org/10.1016/0033-5894(92)90002-Z), 1992.

1669 Thompson, R. S., Anderson, K. H., and Bartlein, P. J.: *Atlas of Relations Between Climatic Parameters*  
1670 *and Distributions of Important Trees and Shrubs in North America—Introduction and Conifers*, U.S.  
1671 Department of the Interior, U.S. Geological Survey, <https://doi.org/10.3133/pp1650AB>, 1999.

1672 Thuiller, W., Lafourcade, B., Engler, R., and Araújo, M. B.: BIOMOD - A platform for ensemble  
1673 forecasting of species distributions, *Ecography*, 32, 369–373, <https://doi.org/10.1111/j.1600-0587.2008.05742.x>, 2009.

1675 Trachsel, M. and Telford, R. J.: All age–depth models are wrong, but are getting better, *The Holocene*,  
1676 27, 860–869, <https://doi.org/10.1177/0959683616675939>, 2017.

1677 Trachsel, M., Dawson, A., Paciorek, C. J., Williams, J. W., McLachlan, J. S., Cogbill, C. V., Foster, D.  
1678 R., Goring, S. J., Jackson, S. T., Oswald, W. W., and others: Comparison of settlement-era vegetation  
1679 reconstructions for STEPPS and REVEALS pollen–vegetation models in the northeastern United States,  
1680 *Quat. Res.*, 95, 23–42, <https://doi.org/10.1017/qua.2019.81>, 2020.

1681 Trondman, A.-K., Gaillard, M.-J., Mazier, F., Sugita, S., Fyfe, R., Nielsen, A. B., Twiddle, C., Barratt, P.,  
1682 Birks, H. J. B., Bjune, A. E., and others: Pollen-based quantitative reconstructions of Holocene regional  
1683 vegetation cover (plant-functional types and land-cover types) in Europe suitable for climate modelling,  
1684 *Glob. Change Biol.*, 21, 676–697, <https://doi.org/10.1111/gcb.12737>, 2015.

1685 Trondman, A.-K., Gaillard, M.-J., Sugita, S., Björkman, L., Greisman, A., Hultberg, T., Lagerås, P.,  
1686 Lindbladh, M., and Mazier, F.: Are pollen records from small sites appropriate for REVEALS model-  
1687 based quantitative reconstructions of past regional vegetation? An empirical test in southern Sweden,  
1688 *Veg. Hist. Archaeobotany*, 25, 131–151, <https://doi.org/10.1007/s00334-015-0536-9>, 2016.

1689 Umbanhowar, Jr., C. E., Camill, P., Geiss, C. E., and Teed, R.: Asymmetric vegetation responses to mid-  
1690 Holocene aridity at the prairie-forest ecotone in south-central Minnesota, *Quat. Res.*, 66, 53–66,  
1691 <https://doi.org/10.1016/j.yqres.2006.03.005>, 2006.

1692 United States Geological Survey: National Hydrography Dataset (NHD Model v2.3.1), 2022.

1693 van Vuuren, D. P., Stehfest, E., Gernaat, D. E. H. J., Doelman, J. C., van den Berg, M., Harmsen, M., de  
1694 Boer, H. S., Bouwman, L. F., Daioglou, V., Edelenbosch, O. Y., Girod, B., Kram, T., Lassaletta, L.,  
1695 Lucas, P. L., van Meijl, H., Müller, C., van Ruijven, B. J., van der Sluis, S., and Tabeau, A.: Energy,  
1696 land-use and greenhouse gas emissions trajectories under a green growth paradigm, *Glob. Environ.*  
1697 *Change*, 42, 237–250, <https://doi.org/10.1016/j.gloenvcha.2016.05.008>, 2017.

1698 Walsh, M. K., Whitlock, C., and Bartlein, P. J.: A 14,300-year-long record of fire–vegetation–climate  
1699 linkages at Battle Ground Lake, southwestern Washington, *Quat. Res.*, 70, 251–264,  
1700 <https://doi.org/10.1016/j.yqres.2008.05.002>, 2008.

1701 Walsh, M. K., Pearl, C. A., Whitlock, C., Bartlein, P. J., and Worona, M. A.: An 11 000-year-long record  
1702 of fire and vegetation history at Beaver Lake, Oregon, central Willamette Valley, *Quat. Sci. Rev.*, 29,  
1703 1093–1106, <https://doi.org/10.1016/j.quascirev.2010.02.011>, 2010.

1704 Walsh, M. K., Marlon, J. R., Goring, S. J., Brown, K. J., and Gavin, D. G.: A regional perspective on  
1705 Holocene fire–climate–human interactions in the Pacific Northwest of North America, *Ann. Assoc. Am.*  
1706 *Geogr.*, 105, 1135–1157, <https://doi.org/10.1080/00045608.2015.1064457>, 2015.

1707 Whitlock, C.: Vegetational and climatic history of the Pacific Northwest during the last 20,000 years:  
1708 Implications for understanding present-day biodiversity, *Northwest Environ. J.*, 8, 5–28, 1992.

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman



1709 Whitlock, C., Colombaroli, D., Conedera, M., and Tinner, W.: Land-use history as a guide for forest  
 1710 conservation and management, *Conserv. Biol.*, 32, 84–97, <https://doi.org/10.1111/cobi.12960>, 2018.  
 1711 Whitmore, J., Gajewski, K., Sawada, M., Williams, J., Shuman, B., Bartlein, P., Minckley, T., Viau, A.,  
 1712 Webb, T., Shafer, S., and others: Modern pollen data from North America and Greenland for multi-scale  
 1713 paleoenvironmental applications, *Quat. Sci. Rev.*, 24, 1828–1848,  
 1714 <https://doi.org/10.1016/j.quascirev.2005.03.005>, 2005.  
 1715 Wieczorek, M. and Herzschuh, U.: Compilation of relative pollen productivity (RPP) estimates and  
 1716 taxonomically harmonised RPP datasets for single continents and Northern Hemisphere extratropics,  
 1717 *Earth Syst. Sci. Data*, 12, 3515–3528, <https://doi.org/10.5194/essd-12-3515-2020>, 2020.  
 1718 Williams, J. W. and Jackson, S. T.: Novel climates, no-analog communities, and ecological surprises,  
 1719 *Front. Ecol. Environ.*, 5, 475–482, <https://doi.org/10.1890/070037>, 2007.  
 1720 Williams, J. W., Shuman, B. N., and Webb, I. T.: Dissimilarity analyses of late-Quaternary vegetation  
 1721 and climate in eastern North America, *Ecology*, 82, 3346–3362, [https://doi.org/10.1890/0012-](https://doi.org/10.1890/0012-9658(2001)082[3346:DAOLQV]2.0.CO;2)  
 1722 [9658\(2001\)082\[3346:DAOLQV\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2001)082[3346:DAOLQV]2.0.CO;2), 2001.  
 1723 Williams, J. W., Shuman, B. N., Webb, T., III, Bartlein, P. J., and Luduc, P. L.: Late-Quaternary  
 1724 vegetation dynamics in North America: scaling from taxa to biomes, *Ecol. Monogr.*, 74, 309–334,  
 1725 <https://doi.org/10.1890/02-4045>, 2004.  
 1726 Williams, J. W., Shuman, B., and Bartlein, P. J.: Rapid responses of the Midwestern prairie-forest ecotone  
 1727 to early Holocene aridity, *Glob. Planet. Change*, 66, 195–207,  
 1728 <https://doi.org/10.1016/j.gloplacha.2008.10.012>, 2009a.  
 1729 Williams, J. W., Shuman, B., and Bartlein, P. J.: Rapid responses of the prairie-forest ecotone to early  
 1730 Holocene aridity in mid-continental North America, *Glob. Planet. Change*, 66, 195–207,  
 1731 <https://doi.org/10.1016/j.gloplacha.2008.10.012>, 2009b.  
 1732 Williams, J. W., Tarasov, P., Brewer, S., and Notaro, M.: Late Quaternary variations in tree cover at the  
 1733 northern forest-tundra ecotone, *J. Geophys. Res. Biogeosciences*, 116, G01017,  
 1734 <https://doi.org/10.1029/2010JG001458>, 2011a.  
 1735 Williams, J. W., Tarasov, P. A., Brewer, S., and Notaro, M.: Late-Quaternary variations in tree cover at  
 1736 the northern forest-tundra ecotone, *J. Geophys. Res. - Biogeosciences*, 116, G01017, doi:  
 1737 [10.1029/2010JG001458](https://doi.org/10.1029/2010JG001458), 2011b.  
 1738 Williams, J. W., Grimm, E. G., Blois, J., Charles, D. F., Davis, E., Goring, S. J., Graham, R., Smith, A. J.,  
 1739 Anderson, M., Arroyo-Cabres, J., Ashworth, A. C., Betancourt, J. L., Bills, B. W., Booth, R. K.,  
 1740 Buckland, P., Curry, B., Giesecke, T., Hausmann, S., Jackson, S. T., Latorre, C., Nichols, J., Purdum, T.,  
 1741 Roth, R. E., Stryker, M., and Takahara, H.: The Neotoma Paleoecology Database: A multi-proxy,  
 1742 international community-curated data resource, *Quat. Res.*, 89, 156–177,  
 1743 <https://doi.org/10.1017/qua.2017.105>, 2018.  
 1744 Worona, M. A. and Whitlock, C.: Late Quaternary vegetation and climate history near Little Lake, central  
 1745 Coast Range, Oregon, *Geol. Soc. Am. Bull.*, 107, 867–876, [https://doi.org/10.1130/0016-](https://doi.org/10.1130/0016-7606(1995)107%3C0867:LQVACH%3E2.3.CO;2)  
 1746 [7606\(1995\)107%3C0867:LQVACH%3E2.3.CO;2](https://doi.org/10.1130/0016-7606(1995)107%3C0867:LQVACH%3E2.3.CO;2), 1995.  
 1747 Yu, Z. C.: Northern peatland carbon stocks and dynamics: a review, *Biogeosciences*, 9, 4071–4085,  
 1748 <https://doi.org/10.5194/bg-9-4071-2012>, 2012.  
 1749 Zapolska, A., Serge, M. A., Mazier, F., Quiquet, A., Renssen, H., Vrac, M., Fyfe, R., and Roche, D. M.:  
 1750 More than agriculture: Analysing time-cumulative human impact on European land-cover of second half  
 1751 of the Holocene, *Quat. Sci. Rev.*, 314, 108227, <https://doi.org/10.1016/j.quascirev.2023.108227>, 2023.

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman