

Environmental controls of rapid terrestrial organic matter mobilization to the western Laptev Sea since the last deglaciation

Tsai-Wen Lin^{1,2}, Tommaso Tesi³, Jens Heftner¹, Hendrik Grotheer^{1,2,4}, Jutta Wollenburg¹, Florian Adolphi^{1,2,4}, Henning A. Bauch^{1,5}, Alessio Nogarotto³, Juliane Müller^{1,2,4}, Gesine Mollenhauer^{1,2,4}

¹Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, 27570, Bremerhaven, Germany

²Department of Geosciences, University Bremen, 28334, Bremen, Germany

³Institute of Polar Sciences, National Research Council, 40129, Bologna, Italy

⁴MARUM Centre for Marine Environmental Sciences, University Bremen, 28334, Bremen, Germany

⁵GEOMAR, Helmholtz Centre for Ocean Research, 24148, Kiel, Germany

Correspondence to: Tsai-Wen Lin (tsai-wen.lin@awi.de) and Gesine Mollenhauer (gesine.mollenhauer@awi.de)

Abstract. Arctic permafrost stores vast amounts of terrestrial organic matter (terrOM). Under warming climate conditions, Arctic permafrost thaws, releasing aged carbon and potentially impacting the modern carbon cycle. We investigated the characteristics of terrestrial biomarkers, including *n*-alkanes, fatty acids, and lignin phenols, in marine sediment cores to understand how the sources of terrOM transported to the ocean change in response to varying environmental conditions such as sea-level rise, sea-ice coverage, inland climate warming, and freshwater input. We examined two sediment records from the western Laptev Sea (PS51/154 and PS51/159) covering the past 17.8 kyr. Our analyses reveal three periods with high mass accumulation rates (MARs) of terrestrial biomarkers, from 14.1 to 13.2, 11.6 to 10.9, and 10.9 to 9.5 kyr BP. These terrOM MAR peaks revealed distinct terrOM sources, likely in response to changes in shelf topography, rates of sea-level rise, and inland warming. By comparing periods of high terrOM MAR in the Laptev Sea with published records from other Arctic marginal seas, we suggest that enhanced coastal erosion driven by rapid sea-level rise during meltwater pulse 1A (mwp-1A) triggered elevated terrOM MAR across the Arctic. Additional terrOM MAR peaks varied regionally. Peaks from the Beaufort Sea during the Bølling-Allerød coincided with a freshwater flooding event, while peaks from the Laptev Sea and the Fram Strait during the Preboreal/early Holocene coincided with periods of enhanced inland warming and prolonged ice-free conditions. Our results highlight the influence of regional environmental conditions, in addition to global drivers, which can either promote or preclude regional terrOM fluxes.

Short Summary. In order to understand the mechanisms governing permafrost organic matter remobilization, we investigated organic matter composition during past intervals of rapid sea-level rise, of inland warming, and of dense sea-ice cover in the Laptev Sea. We find that sea-level rise resulted in wide-spread erosion and transport of permafrost materials to the ocean, but erosion is mitigated by regional dense sea-ice cover. Factors like inland warming or floods increase permafrost mobilization locally.

Keywords. *n*-alkanes, fatty acids, lignin phenols, permafrost thawing, organic matter mass accumulation rate, lignin phenol flux

1 Introduction

Circumarctic permafrost, the ground that has remained below 0 °C for at least two consecutive years, plays an important role in the Arctic carbon cycle and is strongly affected by the rapid warming in the Arctic, which is greater than the average warming of the Northern Hemisphere (Van Everdingen, 2005; Miller et al., 2010; Rantanen et al., 2022). It is one of the main terrestrial carbon sinks at present (Hugelius et al., 2014; Strauss et al., 2017). However, global warming could lead to permafrost thawing, potentially shifting these areas from a carbon sink to a carbon source (Winterfeld et al., 2018; Lara et al., 2019; Laurent et al., 2023). Due to the low ground temperature, organic matter stored in permafrost soils remains protected from degradation and release into the modern carbon cycle (Van Everdingen, 2005; Hugelius et al., 2014; Strauss et al., 2017). When permafrost thaws, the organic matter is transported through fluvial networks and coastal erosion to the ocean, eventually being deposited in marine basins (Schuur et al., 2008; Zhang et al., 2022). In addition to the degradation at site of thawing, during transport, previously freeze-locked organic matter can be decomposed by microbes, releasing greenhouse gases such as CO₂ and CH₄.

The extent of organic matter degradation from thawing permafrost varies between sources and transportation trajectories (Vonk and Gustafsson, 2013; Strauss et al., 2015). Permafrost deposits from different depths (and ages) are mobilized via different pathways. Initially, increasing land temperatures and precipitation mobilize surface permafrost by deepening the active layer and expanding wetland areas. In later stages, the development of thermokarst and taliks exposes and mobilizes deeper permafrost layers (Schuur et al., 2008). Accelerated coastal erosion facilitates the transport of deep permafrost from cliffs to the ocean, particularly in regions with high cliffs such as the Siberian coast (Vonk and Gustafsson, 2013). However, permafrost mobilization caused by accelerated coastal erosion can be mitigated by the presence of land-fast sea ice (Rachold et al., 2000; Overduin et al., 2016; Nielsen et al., 2020; Irrgang et al., 2022). Understanding the dominant pathways of permafrost mobilization under different environmental conditions is critical for evaluating the response of Arctic permafrost to future warming. Changes in terrestrial organic matter (terrOM) mobilization and environmental conditions can be recorded in marine sedimentary archives.

Paleoclimate records from the last deglaciation to the early Holocene (c.a., between 19 and 11 kyr BP) offer insights into Arctic permafrost changes in response to warming and rising sea levels (Clark et al., 2012). During the last deglaciation, atmospheric CO₂ concentrations increased, temperature increased globally and amplified in the Arctic, and global sea levels rose, including rapid events and meltwater pulses (Shakun et al., 2012; Lambeck et al., 2014; Marcott et al., 2014; Köhler et al., 2017). Rapid increases in atmospheric CO₂ concentration occurred at 16.4, 14.6, and 11.5 kyr BP. The 14.6 kyr BP event coincided with an accelerated global sea-level rise and a period of reduced Arctic sea-ice cover, while the 11.5 kyr BP event coincided with intensified inland warming and a drop in sea-ice cover in the Arctic (Fahl and Stein, 2012; Lambeck et al., 2014; Müller and Stein, 2014; Brosius et al., 2021; Detlef et al., 2023). Studying these periods of rapid environmental change can improve the understanding of how current abrupt warming, sea-ice loss, and sea-level rise might affect permafrost stability and the release of previously freeze-locked carbon.

Marine sedimentary archives from the Laptev Sea covering the early period of the last deglaciation (>14 ka BP) are scarce, and existing records often have low temporal resolution and are discontinuous (Tesi et al., 2016a; Keskitalo et al., 2017; Martens et al., 2019; Martens et al., 2020). Here, we present two high-resolution sediment core records that,

collectively, have continuously covered the last 17.8 kyr. We characterize the properties of organic matter exported from Siberian permafrost and relate changes in terrestrial carbon sources to respective climatic conditions. Results were then compared with published studies dealing with the terrOM released into the Arctic Ocean. Further, we explore the interplay of potential factors driving rapid terrestrial carbon translocation, including sea-level fluctuations, inland permafrost stability, and variations in sea-ice cover.




Fig 1. Map of the Arctic Ocean. The names of the marginal seas and archipelago are indicated by acronyms (BS: Beaufort Sea, LS: Laptev Sea, SV: Svalbard archipelago, FS: Fram Strait). The purple area indicates the modern Yedoma domain (Strauss et al., 2016; Strauss et al., 2021; Strauss et al., 2022). The white area shows the ice sheet cover at 18 ka BP (Dyke et al., 2003; Hughes et al., 2016). Deep blue lines indicate the main river streams in the Arctic (Lehner and Grill, 2013). The exposed continental shelf area at 18 kyr BP is labeled in grey. The black line is the -115 m contour line, which is approximately the sea level at the early period of the last deglaciation (18 kyr BP) (Klemann et al., 2015). Yellow dots show the cores from the Arctic used in this study, including cores PS51/154 and PS51/159. Red dots show the other cores from previous studies that are used for comparison, including cores ARA04C/37 (Wu et al., 2020), PC23 (Tesi et al., 2016b), PS2458 (Spielhagen et al., 2005), HH11-09 (Nogarroto et al., 2023), PS2837-5 (Birgel and Hass, 2004), and MSM05/5-712-2 (Aagaard-Sørensen et al., 2014; Müller and Stein, 2014; Zamelezky et al., 2014). The map background is from the International Bathymetric Chart of the Arctic Ocean (IBCAO) (Jakobsson et al., 2012). This map is plotted with QGIS3.22.9 (QGIS Geographic Information System. QGIS Association).

2 Study area

The Laptev Sea is a marginal sea of the Arctic Ocean located north of Siberia (Fig 1). The average depth of the Laptev Sea shelf is around 50 m, with a sharp shelf break located between 70 and 100 m water depth. Several sediment-filled paleo river channels cut through the shelf (Kleiber and Niessen, 2000). Due to its shallow water depth and flat

topography, the Laptev Sea shelf is highly sensitive to sea-level fluctuations. During the last glacial maximum, the Laptev Sea shelf was not covered by an ice sheet and was exposed during the low sea level stand (Hughes et al., 2016), allowing the accumulation of permafrost deposits. The majority of the shelf was inundated between 12 and 6.5 kyr BP, with the rate of inundation slowing down thereafter (Jakobsson et al., 2012; Klemann et al., 2015) (Fig 2h). This rapid shelf inundation led to southward shifts in both the coastline and the locations of major sediment deposition, causing a significant reduction in sediment input to the outer shelf (Bauch et al., 1999; Mueller-Lupp et al., 2000; Bauch et al., 2001). Today, in the Laptev Sea, sediments from water depths <10 m are transported further offshore, and the primary deposition center is located at around 30 m water depth (Kuptsov and Lisitsin, 1996; Are et al., 2002). The Laptev Sea plays a crucial role in the Arctic climate through extensive heat exchange with the atmosphere in summer and sea-ice formation in winter, making it a key region for studying the boundary conditions of the Arctic environment's response to climate change (Rudenko et al., 2014; Liu et al., 2022).

The Laptev Sea shelf receives substantial sediment inputs from both coastal erosion and river discharge (Lantuit et al., 2011; McClelland et al., 2016). Due to high cliffs and intensified wave-induced erosion resulting from summer sea-ice melt, the Laptev Sea receives a large proportion of terrestrial material from coastal erosion. Erosion rates along the Laptev Sea coast are higher (0.7 m yr^{-1}) compared to the Arctic coastal average (0.5 m yr^{-1}) and may even reach more than 10 m yr^{-1} (Rachold et al., 2000; Lantuit et al., 2011; Günther et al., 2015). Additionally, the Lena River serves as the primary source of riverine sediments (McClelland et al., 2016; Holmes et al., 2021), but coastal erosion contributes more than twice the amount of terrestrial export compared to riverine sources (Rachold et al., 2000).

Most of the water discharged by the Lena River is transported eastward and northward, driven by local winds, carrying suspended sediments in the same direction, and guided by the sea bottom relief (Dmitrenko et al., 1999; Guay et al., 2001). In summer, drift ice and river discharges trigger high productivity of the Laptev Sea shelf (Ovsepyan et al., 2015; Hörner et al., 2016). Large amounts of terrigenous suspended particles in the Laptev Sea are incorporated into the sea ice (Dethleff, 2005). This terrigenous-material-rich sea ice is transported to the Fram Strait by the Transpolar Drift, accounting for 20 % of the total sea-ice flow to the Fram Strait (Rigor and Colony, 1997; Stein and Macdonald, 2004).

3 Materials and methods

3.1 Materials

Marine sediment cores PS51/154-11 (77.276° N , 120.610° E , water depth 270 m, abbreviated as PS51/154) and PS51/159-10 (76.768° N , 116.032° E , water depth 60 m, abbreviated as PS51/159) were obtained with Kasten corers during the ARK-XIV/1b (PS51 Transdrift-V) expedition in August 1998 aboard the R/V *Polarstern* (Kassens, 2016).

Core PS51/154 was taken from the upper continental slope, and core PS51/159 from the outer continental shelf (Fig 1). After retrieval, the sediment cores were stored at -20° C and subsampled at 4° C . Subsamples were collected with 10 mL syringes from the cores in 5–10 cm intervals. The sampling intervals in core PS51/154 corresponded to an average temporal resolution of approximately 120 yr before 10 kyr BP, and for core PS51/159, the average temporal resolution was about 270 yr. Due to the spatial proximity and consistency of biomarker proxy records between the

two cores, we used the PS51/159 record to complement the low temporal resolution in core PS51/154 during the Holocene.

3.2 Microfossil radiocarbon dating and age-depth model

The age-depth models for cores PS51/154 and PS51/159 are based on microfossil ^{14}C ages, initially established by Bauch et al. (2001), modified by Taldenkova et al. (2010) and Hörner et al. (2016), and refined in this study. Six new microfossil radiocarbon dates from core PS51/154 and one from core PS51/159 were added to the existing data sets (Table 1). Sediments were wet-sieved through a 63 μm mesh, and bivalve shell fragments were picked from the >63 μm and oven-dried sediment fraction. Bivalve shells were examined under a microscope for species identification and photographic records (Table 1). Samples containing more than 250 μg C were converted to graphite targets, while samples containing less than that were analysed as CO_2 gas using the gas interface system (Table 1). Both types of analysis were conducted using the MIni CARbon Dating System (MICADAS) accelerator mass spectrometer at Alfred Wegener Institute (AWI) (Mollenhauer et al., 2021). We used the Marine20 curve for calibration (Heaton et al., 2020), with a time constant $\Delta R = -95 \pm 61$ yr calculated using the Marine20 database (<http://calib.org/marine/>). This ΔR value was derived from the average reservoir ages of five modern bivalve shells in the Laptev Sea (Bauch et al., 2001). The rationale for using an updated ΔR is the ~150 yr shift in the global marine reservoir age between the Marine13 curve used in the previous age model (Hörner et al., 2016) and the Marine20 curve used in this study (Heaton et al., 2020; Heaton et al., 2023). The dating results are reported as calibrated ages (cal. yr BP). Age-depth models were constructed using OxCal software 4.4 (Bronk Ramsey, 2009), with low model agreement datapoints identified as outliers and removed from the age-depth model. The outlier data are still shown in Table 1 and Fig S1.

Table 1. Radiocarbon dates of cores PS51/154 and PS51/159. The calibrated ages are shown as median ages, with 95.4 % probability ranges in brackets. The ages were calibrated against the Marine20 curve (Heaton et al., 2020), with $\Delta R = -95 \pm 61$ yr.

Lab ID	Depth (cm)	Material	Radiocarbon age (yr BP)	Calibrated age (cal. yr BP)	Reference
PS51/154-11					
KIA-27682	25	foraminifers/ <i>Yoldiella</i> sp.	3425 $\pm$ 30*	3302 (3529–3044)	Taldenkova et al. (2010)
KIA-6919	31	<i>Yoldiella</i> <i>intermedia</i>	1540 $\pm$ 45		Bauch et al. (2001)
KIA-32811	39	bivalves/gastr opods	1800 $\pm$ 35		Taldenkova et al. (2010)
KIA-32810	39	foraminifers	5040 $\pm$ 50*	5343 (5570–5065)	Taldenkova et al. (2010)
KIA-27683	51	foraminifers /ostracods/ <i>Yol</i> <i>diella</i> sp.	9570 $\pm$ 60*	10286 (10530–10024)	Taldenkova et al. (2010)
KIA-32812	73	foraminifers	9410 $\pm$ 70*	10343 (10561–10117)	Taldenkova et al. (2010)
KIA-32813	73	<i>Yoldiella</i> <i>lenticula</i>	9605 $\pm$ 45		Taldenkova et al. (2010)
KIA-27684	85	foraminifers / <i>Portlandia</i> <i>arctica</i>	9505 $\pm$ 50*	10389 (10604–10151)	Taldenkova et al. (2010)
KIA-32814	115	<i>Yoldiella</i> <i>lenticula</i>	9630 $\pm$ 50*	10566 (10864–10263)	Taldenkova et al. (2010)
KIA-32815	131	<i>Nucula tenuis</i>	10085 $\pm$ 45*	11156 (11348–10857)	Taldenkova et al. (2010)

KIA-6920	138	<i>Macoma calcarea</i>	10120 ± 55*	11182 (11383–10890)	Bauch et al. (2001)
KIA-6921	204	<i>Nucula tenuis</i>	10235 ± 45*	11381 (11641–11124)	Bauch et al. (2001)
11524.1.1	223.5	<i>Macoma calcarea</i>	10145 ± 104*	11463 (11740–11170)	This study ²
11525.1.1	247	<i>Macoma calcarea</i>	10482 ± 91*	11702 (12039–11330)	This study ²
KIA-6922	300	<i>Yoldiella intermedia</i>	10725 ± 50*	12183 (12475–11834)	Bauch et al. (2001)
	339.5	bivalves	12040 ± 55*	13479 (13712–13243)	Hörner et al. (2016)
KIA-6923	375	<i>Yoldiella lenticula</i>	12180 ± 60*	13663 (13882–13436)	Bauch et al. (2001)
11526.1.1	392.5	<i>Macoma calcarea</i>	12257 ± 35*	13741 (13963–13514)	This study ¹
11527.1.1	420.5	<i>Macoma calcarea</i>	12189 ± 97*	13850 (14090–13582)	This study ²
KIA-6924	440	<i>Yoldiella intermedia</i>	12525 ± 55*	14091 (14417–13777)	Bauch et al. (2001)
KIA-6925	518	foraminifers	13120 ± 60*	15113 (15427–14804)	Bauch et al. (2001)
		<i>/Portlandia arctica</i>			
11529.1.1	520.5	<i>Macoma calcarea</i>	12855 ± 122		This study ²
11528.1.1	530.5	<i>Yoldiella sp.</i>	14502 ± 44		This study ¹
KIA-9976	567	foraminifers	13540 ± 90*	15659 (15971–15339)	Taldenkova et al. (2010)
KIA-9977	569	foraminifers	13570 ± 110*	15687 (16006–15365)	Taldenkova et al. (2010)
PS51/159-10					
KIA-6927	11	<i>Macoma sp.</i>	845 ± 30*	380 (547–175)	Bauch et al. (2001)
11523.1.1	32.5	<i>Portlandia arctica</i>	2940 ± 24*	2633 (2841–2391)	This study ¹
KIA-6928	56	<i>Portlandia arctica</i>	4980 ± 35*	5194 (5446–4944)	Bauch et al. (2001)
	72.5	bivalve	6610 ± 40		Hörner et al. (2016)
KIA-6929	90	<i>Portlandia arctica</i>	6305 ± 35*	6629 (6865–6399)	Bauch et al. (2001)
KIA-6930	131	<i>Portlandia arctica</i>	8955 ± 40*	9526 (9790–9305)	Bauch et al. (2001)
KIA-6931	215	<i>Portlandia arctica</i>	9420 ± 50*	10160 (10416–9886)	Bauch et al. (2001)
KIA-6932	315	<i>Portlandia arctica</i>	9650 ± 45*	10501 (10773–10230)	Bauch et al. (2001)
KIA-6933	410	<i>Portlandia arctica</i>	10720 ± 55*	12012 (12376–11703)	Bauch et al. (2001)
KIA-6934	485	<i>Portlandia arctica</i>	11060 ± 70*	12517 (12732–12210)	Bauch et al. (2001)

*Dating results that were taken for the age-depth model calculation. ¹Sample analysed as graphite. ²Sample analysed as CO₂ gas.

3.3 Bulk analysis

155 Stable carbon isotope compositions of total organic carbon ($\delta^{13}\text{C}$) in cores PS51/154 and PS51/159 were analysed using 15 mg of freeze-dried, homogenized sediment acidified with 1.5 M HCl in silver boats to remove carbonate. The acidified samples were then dried in the oven at 55 °C. $\delta^{13}\text{C}$ values were measured using a Thermo Scientific FLASH 2000 CHNS Analyser coupled with a Thermo DeltaQ IRMS via CONFLO IV at the Institute of Polar Sciences,

National Research Council (CNR-ISP). $\delta^{13}\text{C}$ values were reported in parts per thousand (per mil, ‰) relative to Vienna Pee Dee Belemnite (VPDB). The standard error for replicate analyses of in-house standards was less than 0.15 ‰.

3.4 Lipid extraction and analysis

The lipid extractions were done following the procedure by Winterfeld et al. (2018). Freeze-dried, homogenized sediment samples were weighed 1–5 g and subsequently submerged in 25 mL dichloromethane (DCM) + methanol ($v/v = 9:1$) followed by ultrasonication (15 min) for three times to acquire total lipid extracts (TLE). An internal quantification standard containing 889.6 ng squalane and 1558 ng 19-methylarachidic acid was added before extraction. After drying the TLE under a stream of nitrogen, the TLE was saponified using 1 mL of 0.1 M potassium hydroxide (KOH) in methanol + H_2O ($v/v = 9:1$) at 80 °C for 2 h. After saponification, the neutral lipids (NLs) were extracted from the TLE with 3×1 mL hexane. Hydrochloric acid (HCl, 37 %) was added to acidify the remaining TLE until $\text{pH} < 2$, and the fatty acids (FAs) were subsequently extracted with 3×1 mL DCM. The neutral lipids (NLs) were further separated into three polarity fractions of hydrocarbons (containing alkanes), ketones, and polar lipids through a silica column by eluting with 4 mL of hexane, DCM + hexane ($v/v = 2:1$), and DCM + methanol ($v/v = 1:1$), respectively. The fatty acids (FAs) were methylated by adding 40 μL 37 % HCl and 1 mL methanol and heated to 50 °C for >12 h to form fatty acid methyl esters (FAMES). Prior to sealing, the head-space of the sample vials was filled with pure nitrogen gas. The FAME fraction was extracted from the methanol solution with 3×1 mL hexane after methylation was completed.

The hydrocarbon and FAME fractions were analysed with an Agilent 7890A gas chromatograph equipped with a flame ionization detector (GC-FID) using an Agilent J&W DB-5MS column (60 m $\times$ 250 μm $\times$ 0.25 μm -thick film). The oven was held at 60 °C for 1 min, heating at a ramp of 20 °C min^{-1} to 150 °C and to 320 °C at 6 °C min^{-1} , with the final holding time of 35 min. $n\text{-C}_{15-34}$ alkanes and $n\text{-C}_{14-32}$ fatty acids were identified by comparing the retention times of a $n\text{-C}_{10-40}$ alkane standard mix and a $n\text{-C}_{28:0}$ FAME standard, respectively. The uncertainty was calculated as the average of the absolute differences between the mean values from duplicate analyses; this was expressed as standard variation (cf. Grotheer et al., 2015). The uncertainty of the total contents of n -alkanes was 11.6 % ($n = 62$), and of FAMES was 12.8 % ($n = 68$).

3.5 Lignin phenol extraction and analysis

The extraction procedure for lignin-oxidation products was modified from the protocol described by Goñi and Montgomery (2000). About 200–300 mg dried and homogenized samples, around 300 mg Copper oxide (CuO), around 50 mg ferrous ammonium sulfate, and 6 mL NaOH (2 N) were added into a Teflon tube in an oxygen-free environment ($\text{O}_2 < 1.0$ %). Samples were oxidized in a microwave digestion system (MARS 6) at 150 °C for 90 min. Afterward, 9.75 μg of ethyl vanillin (Evl) was added as an internal standard; the samples were then centrifuged, and the supernatant was recovered. The solutions were acidified with 3 mL HCl (6 N). The CuO oxidation products, including lignin phenols, were recovered by liquid/liquid extraction with 2×5 mL ethyl acetate. Sodium sulfate (Na_2SO_4) was added to the ethyl acetate extract to remove any remaining water. The extracts were evaporated to dryness using a rotational vacuum concentrator and were subsequently transferred into analytical vials with 500 μL

pyridine for later analysis. Prior to analysis, 40 μL of the extracts dissolved in pyridine were silylated by adding 40 μL N,O-Bis(trimethylsilyl)trifluoroacetamide with 1 % Trimethylchlorosilane and reacted at 50 $^{\circ}\text{C}$ for 15 min. The CuO oxidation products were analysed with an Agilent Technologies 7820A gas chromatograph coupled with a 5977B MSD mass selective detector (GC-MS) equipped with a 30m $\times$ 320 μm $\times$ 0.25 μm -thick film column (Trajan SGE PB-1). The oven was set from 95 $^{\circ}\text{C}$ to 300 $^{\circ}\text{C}$ at a heating ramp of 4 $^{\circ}\text{C min}^{-1}$ with a holding time of 10 min. Here we focused on 3,5-dihydroxybenzoic acid (3,5 Bd) and three lignin phenol groups, including (1) Vanillyl phenols (V): including vanillin (VI), acetovanillone (Vn), and vanillic acid (Vd). (2) Syringyl phenols (S): including syringaldehyde (SI), acetosyringone (Sn), and syringic acid (Sd). (3) Cinnamyl phenols (C): including *p*-coumaric acid (*p*Cd) and ferulic acid (Fd). The uncertainty of the total content for lignin phenols was 6.9 % ($n = 10$).

3.6 Biomarker parameters

We used the *n*-alkane *Sphagnum* proxy defined by Vonk and Gustafsson (2009) as the ratio of the *n*-C₂₅ alkane content to the sum of *n*-C₂₅+C₂₉ alkane contents shown in Eq. (1). Previous studies have shown that *Sphagnum* mosses, which are abundant in peatlands across northern high-latitude regions, predominantly produce mid-chain *n*-alkanes, particularly *n*-C₂₅ (Van Dongen et al., 2008; Vonk and Gustafsson, 2009). In contrast, higher plants primarily synthesize long-chain *n*-alkanes (also recognized as high molecular weight, HMW *n*-alkanes) (Bianchi and Canuel, 2011). A higher C₂₅/(C₂₅+C₂₉) value therefore reflects a greater contribution from peatland-derived sources. The uncertainty of the C₂₅/(C₂₅+C₂₉) proxy was determined to be 0.006, based on the average standard variation of 31 pairs of duplicate analyses ($n = 62$).

$$\text{Sphagnum proxy} = \frac{C_{25}}{C_{25}+C_{29}} \quad (1)$$

The carbon preference index (CPI) was calculated as the ratio of odd-numbered carbon *n*-alkanes to even-numbered carbon *n*-alkanes, as shown in Eq. (2) (Bray and Evans, 1961). Lower CPI values indicate greater thermal maturity of OM, which may result from increased OM degradation (Angst et al., 2016) or contributions from petrogenic sources (Bray and Evans, 1961). The uncertainty of the CPI was 0.05 based on the average standard variation of 31 pairs of duplicate analyses ($n = 62$).

$$\text{CPI} = \frac{C_{23}+2\times C_{25}+2\times C_{27}+2\times C_{29}+2\times C_{31}+C_{33}}{2\times(C_{24}+C_{26}+C_{28}+C_{30}+C_{32})} \quad (2)$$

We used the HMW fatty acids to calculate the temporal change in mass accumulation rate (MAR) to evaluate the terrOM flux into marine basins. Fatty acids in higher plants are dominated by even-numbered HMW saturated fatty acids (Bianchi and Canuel, 2011). The MAR of HMW fatty acids was calculated from Eq. (3). *S* represents the sedimentation rate (cm yr^{-1}), ρ refers to dry bulk density (g cm^{-3}), and the *C*_{24:0}, *C*_{26:0}, *C*_{28:0}, *C*_{30:0} are the contents of the HMW *n*-fatty acids per gram of dry sediment (mg g^{-1}). The unit of MAR is $\text{mg cm}^{-2} \text{ kyr}^{-1}$ (data in Table S1 and Table S2). The average uncertainty of HMW fatty acids MAR was 14.1 % ($n = 68$). One can also use other indices that represent the terrOM source. We calculated the MARs of HMW *n*-alkanes and lignin phenols in cores PS51/154 and PS51/159 as well. The patterns between all the MARs were identical (Fig S3). Here, we only show the MAR of HMW fatty acids for a better comparison to previous studies (Winterfeld et al., 2018; Meyer et al., 2019; Alves et al., 2024).

$$\text{MAR} = S \times \rho \times (C_{24:0} + C_{26:0} + C_{28:0} + C_{30:0}) \quad (3)$$

We used lignin phenols to calculate three indices. The vanillic acid over vanillin ratio (Vd/Vl) reflects the degree of degradation of lignin in the sediments, with higher Vd/Vl indicating more degraded lignin (Hedges et al., 1988). The uncertainty of Vd/Vl was 0.01 based on the average standard variation of five pairs of duplicate analyses ($n = 10$). The syringyl phenols over vanillyl phenols ratio (S/V) indicates the relative contribution of gymnosperm and angiosperm sources, and the cinnamyl phenols over vanillyl phenols ratio (C/V) illustrates the relative contribution of woody and non-woody tissue, respectively (Hedges and Mann, 1979; Goñi and Hedges, 1992; Tesi et al., 2010). The combination of S/V and C/V ratios is a good indicator for identifying lignin phenol sources. The S/V ratio and C/V ratio were calculated with the lignin phenol contents, as shown in Eq. (4) and Eq. (5). The uncertainty of S/V was 0.003 ($n = 10$), and of C/V is 0.01 ($n = 10$).

$$\frac{S}{V} = \frac{Sl+Sn+Sc}{Vl+Vn+Vd} \quad (4)$$

$$\frac{C}{V} = \frac{pCd+Fd}{Vl+Vn+Vd} \quad (5)$$

4 Results

4.1 Chronology and organic matter mass accumulation rates in cores PS51/154 and PS51/159

An updated age-depth model for core PS51/154 was established with 15 radiocarbon dates and for core PS51/159 with 11 radiocarbon dates (Table 1, Fig S1, Fig S2). Core PS51/154 covered the period from 17.5 to 3.0 kyr BP. Three periods of peak sedimentation rates were found: during the end of Bølling-Allerød (from 14.1 to 13.2 kyr BP), the Preboreal (from 11.6 to 10.9 kyr BP), and the early Holocene (from 10.9 to 10.1 kyr BP) (Fig S3). After 10.1 kyr BP, the sedimentation rate in core PS51/154 dropped drastically. Core PS51/159 covered the period from 11.9 to 0.3 kyr BP, with a peak of sedimentation rate during the early Holocene (from 10.6 to 9.5 kyr BP), followed by a significant sedimentation rate drop afterward. The MARs of all biomarkers were largely affected by the pronounced sedimentation rate changes and thus, showed similar temporal changes in all terrestrial biomarkers, including HMW *n*-alkanes, HMW fatty acids, and lignin phenols (Fig S3, contents of each biomarker in Fig S4). Figure 2a and Figure 5g, h show the MAR of HMW fatty acids as a representation.

4.2 Bulk organic records in cores PS51/154 and PS51/159

The record before 16.2 kyr BP in core PS51/154 was characterized by high $\delta^{13}\text{C}$ (-23.8‰) and low TOC values (0.57 %) (Fig S5). A slump layer at 15.4 kyr BP (530–540 cm) was identified by increased grain size (Taldenkova et al., 2010) and a drop in total organic matter (TOC) (Hörner et al., 2016) (Fig S5). However, the $\delta^{13}\text{C}$ value did not show a significant change at this layer. The values of $\delta^{13}\text{C}$ and TOC in cores PS51/154 (-25.6‰ and 0.79 %) and PS51/159 (-25.8‰ and 0.99 %) remained rather constant between 16.2 and 9.5 kyr BP, despite the three periods of peak MAR of terrestrial biomarkers (Fig S5). After the early Holocene (9.6 kyr BP), the $\delta^{13}\text{C}$ value increased, accompanied by a

drop in sedimentation rate in both cores, indicating a decrease in terrestrial input resulting from fast marine transgression (Bauch et al., 1999; Mueller-Lupp et al., 2000; Bauch et al., 2001).

4.3 Lipid and lignin phenol records in cores PS51/154 and PS51/159

The $C_{25}/(C_{25}+C_{29})$ ratio in core PS51/154 dropped to 0.27 between 17.0 and 16.0 kyr BP (Fig 2b), remained stable afterward, began to increase after 12.6 kyr BP, and reached its highest value of 0.41 at 10.8 kyr BP. In core PS51/159, the $C_{25}/(C_{25}+C_{29})$ ratio remained high (0.38) before the early Holocene but decreased subsequently (Fig 2b). The CPI index in core PS51/154 remained low before 16.0 kyr BP, averaging 5.3. The CPI value then increased and remained high until 13.0 kyr BP, with an average of 6.3. Afterward, the CPI value declined, reaching a low value of 5.3 at 10.8 kyr BP, and remained low thereafter (Fig 2c). In core PS51/159, the CPI value decreased from 6.0 during the Preboreal to 5.3 during the mid-Holocene, remained low between 5.5 and 2 kyr BP, and increased slightly to 5.6 in the late Holocene (Fig 2c). Despite variability, the CPI values in cores PS51/154 and PS51/159 (4.5–6.5) range within the typical CPI value for surface and deep permafrost (Sánchez-García et al., 2014; Wild et al., 2022), and remain significantly higher than CPI values from petrogenic sources (~1) (Bray and Evans, 1961).

In the record from core PS51/154, the Vd/Vl ratio was high (1.05) before 16.2 kyr BP (Fig 2d). The Vd/Vl ratio remained stable in both cores before 10 kyr BP, of 0.68 in PS51/154 and 0.77 in PS51/159. In core PS51/159, the Vd/Vl ratio increased since 8.8 kyr BP and reached a stable high value of 1.00 after 6.0 kyr BP. The S/V ratio in core PS51/154 was low (0.64) before 16.0 kyr BP; the ratio increased afterward, peaked at 11.5 kyr BP with a ratio of 0.82, and decreased subsequently (Fig 3). The core PS51/159 record exhibited the same trend. The S/V ratio reached its highest of 0.94 at 11.9 kyr BP, decreased afterward, and reached a stable low value of 0.70 after 9.5 kyr BP (Fig 3). The C/V ratio remained constant (~0.51) in both records, except for one drop at the depth of the slump layer to 0.30 at 15.4 kyr BP in core PS51/154 (Fig 3).




Fig 2. (a–f) The biomarker proxies from cores PS51/154 (dark blue) and PS51/159 (light green) and (g–i) environmental changes in the western Laptev Sea since the last deglaciation. (a) High molecular weight (HMW) fatty acid ($n\text{-C}_{24:0} + n\text{-C}_{26:0} + n\text{-C}_{28:0} + n\text{-C}_{30:0}$) mass accumulation rate (MAR) of cores PS51/154 and PS51/159 (this study), the HMW fatty acid MAR peaks are labeled with numbers. (b) C₂₅/(C₂₅+C₂₉) ratio of cores PS51/154 and PS51/159 (this study). (c) Carbon preference index (CPI) of cores PS51/154 and PS51/159 (this study). (d) Vanillic acid/vanillin ratio (Vd/Vl) of cores PS51/154 and PS51/159 (this study). (e) IP₂₅ contents of cores PS51/154 and PS51/159 (Hörner et al., 2016). (f) Age-depth model controlling points from radiocarbon dating measurements of cores PS51/154 and PS51/159. (g) Rate of sea-level rise in the western Laptev Sea (Klemann et al., 2015). (h) Area of land inundation per kyr in the western Laptev Sea, calculated from the sea-level reconstruction from Klemann et al. (2015) and the bathymetric data from the international bathymetric chart of the Arctic Ocean (IBCAO) (Jakobsson et al., 2012). (i) Counts of newly developed thermokarst lakes, categorized by the basal ages of the reported thermokarst lakes (number within 500-yr bins) in Siberia (Brosius et al., 2021). The color bars highlight the periods with HMW fatty acid MAR peaks from 14.1 to 13.2 kyr BP (blue, terrOM MAR peak I), from 11.6 to 10.9 kyr BP (red, terrOM MAR peak II), and from 10.9 to 9.5 kyr BP (green, terrOM MAR peak III). The names of different

paleoclimate periods are indicated by acronyms (HS1: Heinrich Stadial 1, B/A: Bølling-Allerød, YD: Younger Dryas, PB: Preboreal).




Fig 3. Vegetation source parameters derived from ratios of syringyl/vanillyl phenols (S/V) and cinnamyl/vanillyl phenols (C/V) of core records in the Laptev Sea and terrestrial records from Siberia. The core records include cores PS51/154 (dots, this study), PS51/159 (crosses, this study), and PC23 (Tesi et al., 2016b). For data from PS51/154 and PS51/159, symbol colors mark data from different terrOM MAR peaks: blue: terrOM MAR peak I period (from 14.1 to 13.2 kyr BP in core PS51/154); red: terrOM MAR peak II in core PS51/154 (from 11.6 to 10.9 kyr BP) and core bottom in core PS51/159 (from 12 to 10.6 kyr BP); green: terrOM MAR peak III (from 10.9 to 10.1 kyr BP in core PS51/154 and 10.6 to 9.5 kyr BP in core PS51/159). Diamond data points indicate Siberian terrestrial records, including Holocene peat permafrost collected from the Lena Delta (peat PF) (Winterfeld et al., 2015), suspended particulate matter from the Lena Delta (Lena SPM) (Winterfeld et al., 2015), Yedoma (Tesi et al., 2014), active layer from the tundra region (AL tundra) (Tesi et al., 2014), and active layer from the taiga region (AL taiga) (Tesi et al., 2014).

5 Discussion

5.1 Temporal changes of terrOM characteristics in the western Laptev Sea since the last deglaciation

The record from core PS51/154 before 16 kyr BP suggests that land-to-ocean terrOM transport was low, which could have resulted from permanent sea-ice cover (Hörner et al., 2016) (Fig 2e) and the low terrestrial supply due to cold and dry hinterland condition (Andreev et al., 2003). Specifically, high $\delta^{13}\text{C}$ and low OC-normalized terrestrial biomarker contents suggest limited terrOM input from land (Fig S5). The high Vd/Vl ratio and low CPI value during this period indicate either enhanced degradation of the small amount of terrOM reaching the core site due to longer transport times from land to the shelf caused by sea-ice blockage or, alternatively, the terrOM originated from an already degraded pool (Fig 2c, d).

Three distinct peaks of HMW fatty acid MAR were recorded in core PS51/154, occurring from 14.1 to 13.2 kyr BP (terrOM MAR peak I), from 11.6 to 10.9 kyr BP (terrOM MAR peak II), and from 10.9 to 10.1 kyr BP (terrOM MAR peak III) (Fig 2a). Core PS51/159 recorded one HMW fatty acid MAR peak, from 10.6 to 9.5 kyr BP (terrOM MAR peak III) (Fig 2a). The absence of terrOM MAR peaks before 10.6 kyr BP in core PS51/159 may be because the water depth at this location was too shallow, and the main sediment depocenter of the Laptev Sea was located further north at that time (Stein and Fahl, 2000; Bauch et al., 2001). The lack of terrOM MAR peak II could also be due to the lack of available chronology tie points in core PS51/159 during this period (Fig 2f). In addition to reflecting elevated terrestrial input, these elevated HMW fatty acid MAR peaks could also indicate enhanced organic matter preservation in marine sediments. However, disentangling these two factors is challenging, primarily because marine primary production likely increased during periods of elevated terrOM export. Terrestrial nutrients serve as a critical source that fuels marine primary production in the Arctic Ocean (Terhaar et al., 2021). As a result, increased terrOM input could stimulate marine organic matter production, leading to concurrent variations in the contents of marine-source and terrestrial-source organic matters (Fig S6). Notably, from the Bølling-Allerød to the early Holocene, the Vd/Vl ratio remained relatively consistent while the CPI value showed a gradual decrease (Fig 2c, d). Neither the Vd/Vl ratio nor the CPI value exhibited noticeable differences between periods of high and low HMW fatty acid MAR. This lack of variation hinted that terrOM preservation did not significantly increase during the HMW fatty acid MAR peak periods.

Our records suggest that different mechanisms of terrOM mobilization led to varying terrOM characteristics. During the period of terrOM MAR peak I, rapid sea-level rise resulted in enhanced coastal erosion on the steep coast near the shelf break (Fig 4a). The intruding warm Atlantic seawater inhibited sea-ice formation, further aggravating coastal erosion (Taldenkova et al., 2010; Hörner et al., 2016) (Fig 2a, e, Fig 4a). As observed in modern environments, coastal areas with high cliffs experience accelerated erosion when rapid sea-level rise causes waves to reach higher steep cliffs (Limber et al., 2018; Shadrick et al., 2022). The relatively cold and dry Siberian inland hindered peatland development (Fig 2i) (Hubberten et al., 2004; Brosius et al., 2021). The combination of enhanced coastal erosion and limited peat sources is reflected in low $C_{25}/(C_{25}+C_{29})$ proxy and low S/V ratios during terrOM MAR peak I compared to the later terrOM MAR peak periods (Fig 2b, Fig 4a). The relatively high CPI value and low Vd/Vl ratio also implied that the terrOM was sourced from less degraded terrestrial permafrost.

During the periods of terrOM MAR peak II and terrOM MAR peak III, increasing temperature and humidity in the Siberian hinterland facilitated contemporary peat development, as reflected by the increasing tundra and shrub vegetation (Andreev et al., 2003; Hubberten et al., 2004; Andreev et al., 2011) and peat cover (Smith et al., 2004; Yu et al., 2010) in northern Siberia. The increasing temperature also facilitates thermokarst slump formation, which exposed deep old peatland (Schuur et al., 2015) (Fig 2i). The peak $C_{25}/(C_{25}+C_{29})$ ratio values during this period indicate that terrestrial peat deposits, either developed contemporarily or exposed from deep permafrost, were transported by river or/and widely flooded shelf and mobilized (Fig 2b). The elevated peat input during terrOM MAR peak II was also reflected by an increased soft angiosperm tissue contribution, originating from grass-like material abundant in Holocene peats and Holocene permafrost (Winterfeld et al., 2015) (Fig 3). The S/V ratio decreased during terrOM MAR peak III, indicating an elevated contribution of gymnosperm plants, likely reflecting the northward expansion

of the conifer tree line due to rising temperature and humidity during this period (Hubberten et al., 2004; Wild et al., 2022) (Fig 2b, Fig 3). During this period, the Laptev Sea shelf was rapidly inundated due to its flat topography, and the inundation rate remained high even after the sea-level rise slowed down (Mueller-Lupp et al., 2000; Bauch et al., 2001; Klemann et al., 2015) (Fig 2g, Fig 4b, c). In coastal areas with gentle slopes, rapid sea-level rise results in accelerated marine transgression and inundates terrestrial permafrost. Seawater also increases the temperature of the inundated permafrost and facilitates thawing (Overduin et al., 2016). The rapid inundation could be a major driver of elevated terrOM mobilization. Increased inland warming and shelf inundation rates likely facilitate terrOM degradation both on land and during cross-shelf transport (Bröder et al., 2018; Brosius et al., 2021), as evidenced by the decreasing CPI value in core PS51/154 (Fig 2c). However, the Vd/Vl ratio during this period remained consistent with that of the previous period of terrOM MAR peak I (Fig 2d). Since *n*-alkanes predominately source from deep permafrost, whereas lignin phenols primarily reflect surface transport (Feng et al., 2013; Feng et al., 2015), the divergence between these two indices may indicate a shift in terrOM source. Our records highlight that variations in terrOM reflect changes in inland permafrost stability and marine transgression dynamics.

An additional source of brassicasterol, typically ascribed to marine diatoms, during the period of terrOM MAR peak II in core PS51/154 was attributed to increased river runoff during this period (Hörner et al., 2016). This additional river runoff was likely not originating from the Lena River, as the lignin phenol assemblages in core PS51/154 differed significantly from that of the nearby sediment core records (PC23) (Tesi et al., 2016b) and Lena River suspended particulate matter (Lena SPM) (Winterfeld et al., 2015) (Fig 3). Instead, the riverine terrOM deposited in the western Laptev Sea likely originated from the Olenyok or Khatanga Rivers rather than the Lena River (Fig 4b). This is supported by modern observations that most freshwater and suspended sediment discharged from the Lena River are transported eastward to the eastern Laptev Sea (Dmitrenko et al., 1999; Guay et al., 2001), and the mineral assemblages in western Laptev Sea surface sediments resemble those from the Khatanga River rather than the Lena River (Dethleff et al., 2000).

Overall, the S/V and C/V ratios in cores PS51/154 and PS51/159 are higher than those in the Lena SPM (Fig 3) (Winterfeld et al., 2015). The low S/V and C/V ratios in the Lena SPM suggest a significant contribution of woody gymnosperm tissues transported from boreal forests in the southern part of the Lena River catchment (Tesi et al., 2016b; Wild et al., 2022). In contrast, the higher S/V and C/V ratios in PS51/154 and PS51/159 indicate that the vegetation source of the western Laptev Sea primarily originated from higher latitudes, reflecting a regional signal. This is further supported by the less degraded lignin phenol signals (lower Vd/Vl ratio) in cores PS51/154 and PS51/159 compared to those in the Lena SPM, which could be due to either a shorter transport distance or a better-preserved organic matter that was freeze-locked in permafrost (Wild et al., 2022).




Fig 4. Flooded shelf area during the periods of high HMW fatty acid MAR recorded in cores PS51/154 and PS51/159. (a) terrOM MAR peak I, from 14.1 to 13.2 kyr BP. (b) terrOM MAR peak II, from 11.6 to 10.9 kyr BP. (c) terrOM MAR peak III, from 10.9 to 9.5 kyr BP. The grey area shows the exposed continental shelf. The red areas show the flooded area during the periods. The flooded and exposed areas are calculated using the modern bathymetry map from IBCAO (Jakobsson et al., 2012). The river data are from Lehner and Grill (2013). The dark grey areas denote the paleo river channel (Kleiber and Niessen, 2000). The thick black line denotes the contour of -115 m sea level, which approximates the sea level at 18 kyr BP. Red dots are the core locations in the Laptev Sea, including cores PC23 (Tesi et al., 2016b), PS2458 (Spielhagen et al., 2005), PS51/154 (this study), and PS51/159 (this study). The orange arrow indicates the period that the Laptev Sea shelf experienced a strong impact from Atlantic water (Taldenkova et al., 2010). The maps are plotted with QGIS3.22.9 (QGIS Geographic Information System. QGIS Association).

5.2 Environmental change affecting rapid organic matter transport to the ocean on cross Arctic and regional scales

In order to evaluate whether the described terrOM MAR peaks reflected cross Arctic scale climate dynamics, we compared our HMW fatty acid MAR results with other terrOM MAR records from the Arctic marginal seas. Studies have used different biomarkers to trace terrestrial inputs, including lignin phenols, HMW *n*-alkanes, and campesterol and β -sitosterol. Campesterol and β -sitosterol are biosynthesized in vascular plants and thus reflect terrestrial signals, in contrast to dinosterol and brassicasterol, which are found abundant in marine plankton (Bianchi and Canuel, 2011). Data were collected from sites in the Beaufort Sea (ARA04C/37) (Wu et al., 2020), the Laptev Sea (PC23) (Tesi et al., 2016b), the northern Svalbard continental margin (HH11-09) (Nogarotto et al., 2023), and the Fram Strait (PS2837-5 and MSM05/5-712-2) (Birgel and Hass, 2004; Aagaard-Sørensen et al., 2014; Müller and Stein, 2014; Zamelczyk et al., 2014) (Fig 1, Fig 5).

TerrOM MAR largely depends on changes in sedimentation rate (Fig 5, Fig S7), which can vary significantly between age models. Using age models with dense chronological control points is crucial. To ensure confidence in the timing of terrOM MAR peaks, we selected records with multiple chronological tie-points both below and above the identified terrOM MAR peaks (Fig 5d, f, g, h) and multiple records from the same region to provide a more comprehensive regional pattern (Fig 5f–h, i–k).

Age-depth models for these records were recalibrated against the Marine20 calibration curve (Heaton et al., 2020) or a combination of the IntCal20 (Reimer et al., 2020) and Marine20 curves, depending on the original studies. For cores PC23 and HH11-09, which already have published age models updated to the Marine20 or Marine20 + IntCal20 curves, we adopted the existing age models (Nogarotto et al., 2023; Sabino et al., 2024). For records from the Fram Strait (PS2837-5, MSM05/5-712-2), updated ΔR values were derived from the Marine20 database. We applied a constant updated ΔR for calibration, following their previously published age models (Birgel and Hass, 2004; Aagaard-Sørensen et al., 2014; Müller and Stein, 2014; Zamelczyk et al., 2014). Since no ΔR values were available for the Beaufort Sea region in the Marine20 database, new ΔR values for each dating point were calculated by subtracting 150 yr from the previous ΔR values for each point (Keigwin et al., 2018), following the guidelines from Heaton et al. (2023). Detailed information on the updated ΔR values is provided in Table S3. We further calculated the age uncertainty of each terrOM MAR peak by including ± 1 -sigma uncertainty from the age models. The possible age ranges of these terrOM MAR peaks are shown in Fig 5.




Fig 5. Environmental changes since the last deglaciation and terrestrial biomarker mass accumulation rates (MARs) of core records in the Arctic Ocean and higher-latitude northern hemisphere. (a) Atmospheric CO₂ concentration (Marcott et al., 2014; Köhler et al., 2017). (b) Rate of global sea-level change (Lambeck et al., 2014). (c) Compilation of basal ages of thermokarst lakes (number within 500-yr bins) in North America (Brosius et al., 2021). (d) Campesterol + β-sitosterol MAR from core ARA04C/37, Beaufort Sea (Wu et al., 2020). (e) Compilation of basal ages of thermokarst lakes (number within 500-yr bins) in Siberia (Brosius et al., 2021). (f) Lignin phenol MAR from core PC23, Laptev Sea (Tesi et al., 2016b). (g) HMW fatty acid MAR from core PS51/154, Laptev Sea (this study). (h) HMW fatty acid MAR from core PS51/159, Laptev Sea (this study). (i) Lignin phenol MAR from core HH11-09, northern Svalbard continental margin (Nogarotto et al., 2023). (j) HMW *n*-alkane MAR from core PS2837-5, Fram Strait (Birgel and Hass, 2004). (k) β-sitosterol MAR from core MSM05/5-712-2, Fram Strait (Aagaard-Sørensen et al., 2014; Müller and Stein, 2014; Zamelczyk et al., 2014). The blue

bars highlight the period of rapid sea-level rise. The black intervals under each terrOM MAR peak indicate the age uncertainty range of the terrOM MAR peaks, calculated by the ± 1 -sigma age models. Meltwater pulses are denoted as mwp-1A and mwp-1B. The names of different paleoclimate periods are indicated by acronyms (HS1: Heinrich Stadial 1, B/A: Bølling-Allerød, YD: Younger Dryas, PB: Preboreal).

5.2.1 Effects of global sea-level rise

The rapid global sea-level rise during meltwater pulse 1A (mwp-1A) was an important process in terrOM mobilization across different regions in the Arctic. TerrOM MAR peaks during this period are observed widely in records from the Eurasian Arctic (Fig 5b, d, g, i–k) (Birgel and Hass, 2004; Lambeck et al., 2014; Nogarotto et al., 2023). Inland temperatures in both North America and Siberia remained low during this time (Fig 5c, e). These concurrent terrOM MAR peaks suggest that rapid sea-level rise was the primary cause for circumarctic permafrost mobilization, possibly contributing to the rapid increase in atmospheric CO₂ concentration (Marcott et al., 2014; Winterfeld et al., 2018). The record from core ARA04C/37 did extend to the mwp-1A period. However, increased transport of old terrOM was observed at the terrOM MAR peak in ARA04C/37 during the pedios between 13.6 and 13 kyr BP, suggesting contributions from deep permafrost mobilized from coastal erosion (Wu et al., 2022). This terrOM peak may have begun earlier than the time period covered by the record and is likely associated with the rapid sea-level rise during the mwp-1A period (Fig 5b).

During mwp-1B, terrOM MAR also increased in records from the Beaufort Sea and the Laptev Sea (Fig 5b, d, f, g). However, this peak was absent in records from Svalbard/Fram Strait (Fig 5i, j, k). One of the possible reasons could be the lack of radiocarbon age controls in these cores during this period. Also, the amplitude of sea-level rise and even the existence of mwp-1B remain debated (Lambeck et al., 2014). The lack of widespread terrOM MAR during mwp-1B suggests that this event might not affect the pan-Arctic region as extensively as mwp-1A did.

5.2.2 Regional processes: inland warming, sea-ice loss, and freshwater pulses

Outside the periods of mwp-1A and mwp-1B (indicated by blue bars in Fig 5), several regional terrOM MAR peaks were observed in the Arctic marginal seas. These peaks were likely driven by regional factors such as inland warming, decreased sea-ice cover, and freshwater pulses. Warming facilitates permafrost thawing and thermokarst lake development (Schuur et al., 2015). Lake basal ages indicate the time of lake formation, and a higher count of thermokarst lake formation in a certain period thus implies intense inland warming (Brosius et al., 2021). In the Canadian Arctic, terrOM MAR peaks appeared during the interval between mwp-1A and mwp-1B (Fig 5d). Inland warming in North America began at approximately 13.5 kyr BP, while Siberia remained cold (Brosius et al., 2021). TerrOM MAR peaks appeared during the early Holocene across the Siberian Arctic and Fram Strait records (Fig 5g, h, k). During the early Holocene, enhanced warming in Siberian hinterlands led to elevated terrOM mobilization (Fig 5e). These terrOM MAR peaks occurred during periods of little sea-level variation, indicating that intensified inland warming could destabilize permafrost and result in regional terrOM mobilization even in the absence of rapid sea-level rise.

Notably, the observed terrOM MAR peaks in the Arctic marginal seas coincided with regionally reduced seasonal sea-ice cover, as indicated by reduced IP₂₅ (Müller et al., 2009; Hörner et al., 2016; Wu et al., 2020). IP₂₅ refers to an alkane that is produced by sea-ice associated diatoms (Volkman et al., 1994). The existence of IP₂₅ indicates the

appearance of seasonal sea ice, while the lack of IP₂₅ implies either the lack of seasonal sea-ice or permanent sea-ice coverage in the area (Belt and Müller, 2013). Studies have shown that accelerated coastal erosion can be mitigated by the presence of land-fast sea ice, which prevents thermal denudation by sea water, direct wave erosion, and saltwater intrusion into permafrost coasts (Rachold et al., 2000; Overduin et al., 2016; Nielsen et al., 2020; Irrgang et al., 2022). Modern observations have associated elevated coastal erosion rates in the Laptev Sea with reduced winter sea-ice cover (Nielsen et al., 2020). Modeling studies also indicate a positive correlation between inland permafrost stability and winter sea-ice extent under modern conditions (Vandenberghe et al., 2012). The terrOM MAR peak III in core PS51/154 and PS51/159 reflected that even without rapid sea-level rise, the mobilized material caused by inland warming was more easily transported to marine basins because of the lack of sea-ice protection on the coast (Fig 2a, e). A similar relation was shown in the core from the Beaufort Sea (ARA04C/37) as the two terrOM MAR peaks registered in the core both happened during an ice-free period (Wu et al., 2020). The Holocene record of core PS51/159 shows the opposite condition. The seasonal sea-ice cover started to increase since 7 kyr BP (Hörner et al., 2016). Even though several warming periods in Siberia were recorded since then, no terrOM MAR peak was recorded in core PS51/159 (Fig 2a, e, i). The growing sea ice might play a role in protecting the warming land from being eroded and keeping the terrestrial material from being transported into the Laptev Sea shelf.

Freshwater floods triggered by ice-dammed lake breakings could be another regional factor in terrOM mobilization. These flooding events can be recorded by the decreasing stable oxygen isotope ratio ($\delta^{18}\text{O}$) in planktic foraminiferas *Neogloboquadrina pachyderma* (Spielhagen et al., 2005; Keigwin et al., 2018). In the Canadian Arctic record (ARA04C/37), the terrOM MAR peak occurring during the Younger Dryas was linked to a well-described meltwater flood event through the Mackenzie River, based on the drop of both $\delta^{18}\text{O}$ value in *Neogloboquadrina pachyderma* and TOC (Keigwin et al., 2018; Klotsko et al., 2019; Wu et al., 2020) (Fig S8). However, freshwater events were less likely to be the cause for terrOM MAR in the western Laptev Sea. While freshwater flooding events were recorded in an ice-dammed lake upstream of the Lena River (14.9 ± 2.0 kyr BP) and in a sediment record from the Laptev Sea (PS2458, at 12 kyr BP) (Spielhagen et al., 2005; Margold et al., 2018), the timing of these events did not correspond with any of the terrOM MAR peaks in cores PS51/154 and PS51/159 (Fig S8), nor causing any grain size change in cores PS51/154 and PS51/159 (Taldenkova et al., 2010). This temporal mismatch suggests that Siberian freshwater pulses had little impact on the increase in terrestrial biomarker MAR in the western Laptev Sea.

6 Summary and Conclusions

This study explores temporal changes in the composition and rate of terrOM mobilization to the western Laptev Sea, as indicated by lipid and lignin phenol records. Three rapid terrigenous organic matter supply events were identified from 14.1 to 13.2, from 11.6 to 10.9, and from 10.9 to 9.5 kyr BP. Each peak showed different compositional characteristics, suggesting distinct terrOM sources derived from different mechanisms. The first terrOM MAR peak was likely driven by enhanced coastal erosion, while the latter two peaks were associated with inland warming and rapid shelf inundation. The source shift was characterized by increased peat input, as evidenced by both $C_{25}/(C_{25}+C_{29})$ proxy and S/V ratios. Comparing our findings with records from across the Arctic indicates that the enhanced terrOM mobilization during mwp-1A period was primarily driven by enhanced coastal erosion resulting from rapid global

sea-level rise. However, terrOM MAR peaks did not always align with periods of rapid sea-level rise, suggesting that other regional factors, such as inland warming, lack of sea-ice protection, and freshwater floods, also played significant roles.

Overall, the study highlights that the topography of the western Laptev Sea shelf strongly influenced erosion scenarios linked to sea-level rise, leading to different terrOM sources mobilized from land to ocean. Our results suggest that while rapid sea-level rise contributed to elevated terrOM mobilization on a cross Arctic scale, regional factors such as inland warming, freshwater floods, and sea-ice cover decrease were responsible for regional terrOM mobilization.

According to the IPCC report, the projected global sea-level rise by the end of the 21st century could reach up to 8 m kyr⁻¹ (Church et al., 2013). While this projected rate is significantly slower than the rapid sea-level rise during the mwp-1A (~35 m kyr⁻¹) and mwp-1B (~20 m kyr⁻¹) periods (Lambeck et al., 2014), our results from the last deglaciation indicate that even without such rapid sea-level rise, the combination of reduced sea-ice protection and increased hinterland warming can still mobilize substantial terrOM through regional coastal erosion. Given that the Arctic is experiencing rapid warming compared to the global average, and destabilization of permafrost and sea-ice decrease is already underway (Meredith et al., 2019), an increase in terrOM input from coastal erosion is likely if the warming persists.

Author contributions

GM designed the study; TWL, JH, HG, JW and AN performed the measurements; TWL, TT, JH, HG, FA, and GM analysed the data; FA and JW verified the age models; TWL wrote the manuscript draft with the support from TT and GM; all the coauthors reviewed and edited the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgment

This study was supported by the German-Italian partnership project between Alfred Wegener Institute and CNR-ISP on Chronologies for Polar Paleoclimate Archives (PAIGE), funded by the Helmholtz Association. FA was supported through the Helmholtz association (VH-NG 1501). We thank the captain, the chief scientist, the crew, and the scientific party of the *Polarstern* Expedition ARK-XIV/1b (PS51 Transdrift-V) for providing the studied material. We thank Kirsten Fahl and Rüdiger Stein for offering coring information. We thank the MICADAS radiocarbon laboratory group members at Alfred Wegener Institute, namely Elizabeth Bonk, Torben Gentz, and Lea Philips. We appreciate help from Arnaud Nicolas in core subsampling. Katarzyna Zamełczyk and Steffen Aagaard Sørensen are acknowledged for dry bulk density data of core MSM05/5-712-2. We appreciate constructive suggestions from the editor Bjørg Risebrobakken and two anonymous reviewers. ChatGPT (version GPT-4) was used in this manuscript to edit grammar and improve sentence fluency.

540 Data availability

Data generated in this study will be uploaded to PANGAEA database.

References

- Aagaard-Sørensen, S., Husum, K., Werner, K., Spielhagen, R. F., Hald, M., and Marchitto, T. M.: A Late Glacial–Early Holocene multiproxy record from the eastern Fram Strait, Polar North Atlantic, *Marine Geology*, 355, 15-26, 10.1016/j.margeo.2014.05.009, 2014.
- Alves, E. Q., Wong, W., Hefter, J., Grotheer, H., Tesi, T., Gentz, T., Zonneveld, K., and Mollenhauer, G.: Deglacial export of pre-aged terrigenous carbon to the Bay of Biscay, *Climate of the Past*, 20, 121-136, 10.5194/cp-20-121-2024, 2024.
- Andreev, A. A., Schirrmeister, L., Tarasov, P. E., Ganopolski, A., Brovkin, V., Siebert, C., Wetterich, S., and Hubberten, H.-W.: Vegetation and climate history in the Laptev Sea region (Arctic Siberia) during Late Quaternary inferred from pollen records, *Quaternary Science Reviews*, 30, 2182-2199, 10.1016/j.quascirev.2010.12.026, 2011.
- Andreev, A. A., Tarasov, P. E., Siebert, C., Ebel, T., Klimanov, V. A., Melles, M., Bobrov, A. A., Dereviagin, A. Y., Lubinski, D. J., and Hubberten, H. W.: Late Pleistocene and Holocene vegetation and climate on the northern Taymyr Peninsula, Arctic Russia, *Boreas*, 32, 484-505, 2003.
- Angst, G., John, S., Mueller, C. W., Kogel-Knabner, I., and Rethemeyer, J.: Tracing the sources and spatial distribution of organic carbon in subsoils using a multi-biomarker approach, *Sci Rep*, 6, 29478, 10.1038/srep29478, 2016.
- Are, F., Grigoriev, M. N., Hubberten, H.-W., Rachold, V., Razumov, S., and Schneider, W.: Comparative shoreface evolution along the Laptev Sea coast, *Polarforschung*, 135-150, 2002.
- Bauch, H. A., Kassens, H., Erlenkeuser, H., Grootes, P. M., and Thiede, J.: Depositional environment of the Laptev Sea (Arctic Siberia) during the Holocene, *Boreas*, 28, 194-204, 1999.
- Bauch, H. A., Mueller-Lupp, T., Taldenkova, E., Spielhagen, R. F., Kassens, H., Grootes, P. M., Thiede, J., Heinemeier, J., and Petryashov, V.: Chronology of the Holocene transgression at the North Siberian margin, *Global Planetary Change*, 31, 125-139, 2001.
- Belt, S. T. and Müller, J.: The Arctic sea ice biomarker IP25: a review of current understanding, recommendations for future research and applications in palaeo sea ice reconstructions, *Quaternary Science Reviews*, 79, 9-25, 2013.
- Bianchi, T. S. and Canuel, E. A.: Chemical biomarkers in aquatic ecosystems, in: *Chemical Biomarkers in Aquatic Ecosystems*, Princeton University Press, 2011.
- Birgel, D. and Hass, H.: Oceanic and atmospheric variations during the last deglaciation in the Fram Strait (Arctic Ocean): a coupled high-resolution organic-geochemical and sedimentological study, *Quaternary Science Reviews*, 23, 29-47, 10.1016/j.quascirev.2003.10.001, 2004.
- Bray, E. and Evans, E.: Distribution of n-paraffins as a clue to recognition of source beds, *Geochimica et Cosmochimica Acta*, 22, 2-15, 1961.

Bröder, L., Tesi, T., Andersson, A., Semiletov, I., and Gustafsson, O.: Bounding cross-shelf transport time and degradation in Siberian-Arctic land-ocean carbon transfer, *Nature Communications*, 9, 806, 10.1038/s41467-018-03192-1, 2018.

575 Bronk Ramsey, C.: Bayesian Analysis of Radiocarbon Dates, *Radiocarbon*, 51, 337-360, 10.1017/s0033822200033865, 2009.

Brosius, L. S., Anthony, K. M. W., Treat, C. C., Lenz, J., Jones, M. C., Bret-Harte, M. S., and Grosse, G.: Spatiotemporal patterns of northern lake formation since the Last Glacial Maximum, *Quaternary Science Reviews*, 580 253, 10.1016/j.quascirev.2020.106773, 2021.

Church, J. A., Clark, P. U., Cazenave, A., Gregory, J. M., Jevrejeva, S., Levermann, A., Merrifield, M. A., Milne, G. A., Nerem, R. S., and Nunn, P. D.: *Sea level change*, PM Cambridge University Press, 2013.

Clark, P. U., Shakun, J. D., Baker, P. A., Bartlein, P. J., Brewer, S., Brook, E., Carlson, A. E., Cheng, H., Kaufman, D. S., Liu, Z., Marchitto, T. M., Mix, A. C., Morrill, C., Otto-Bliesner, B. L., Pahnke, K., Russell, J. M., Whitlock, C., 585 Adkins, J. F., Blois, J. L., Clark, J., Colman, S. M., Curry, W. B., Flower, B. P., He, F., Johnson, T. C., Lynch-Stieglitz, J., Markgraf, V., McManus, J., Mitrovica, J. X., Moreno, P. I., and Williams, J. W.: Global climate evolution during the last deglaciation, *Proceedings of the National Academy of Sciences of the United States of America*, 109, E1134-1142, 10.1073/pnas.1116619109, 2012.

Dethleff, D.: Entrainment and export of Laptev Sea ice sediments, Siberian Arctic, *Journal of Geophysical Research: Oceans*, 110, 10.1029/2004jc002740, 2005.

590 Dethleff, D., Rachold, V., Tintelnot, M., and Antonow, M.: Sea-ice transport of riverine particles from the Laptev Sea to Fram Strait based on clay mineral studies, *International Journal of Earth Sciences*, 89, 496-502, 2000.

Detlef, H., O'Regan, M., Stranne, C., Jensen, M. M., Glasius, M., Cronin, T. M., Jakobsson, M., and Pearce, C.: Seasonal sea-ice in the Arctic's last ice area during the Early Holocene, *Communications Earth & Environment*, 4, 595 10.1038/s43247-023-00720-w, 2023.

Dmitrenko, I., Golovin, P., Griбанov, V., and Kassens, H.: Oceanographic causes for transarctic ice transport of river discharge, in: *Land-Ocean Systems in the Siberian Arctic*, edited by: Kassens, H., Bauch, H. A., Dmitrenko, I. A., Eicken, H., Hubberten, H.-W., Melles, M., Thiede, J., and Timokhov, L. A., Springer Berlin, Heidelberg, 73-92, 1999.

Dyke, A. S., Moore, A., and Robertson, L.: Deglaciation of North America [dataset], <https://doi.org/10.4095/214399>, 600 2003.

Fahl, K. and Stein, R.: Modern seasonal variability and deglacial/Holocene change of central Arctic Ocean sea-ice cover: New insights from biomarker proxy records, *Earth and Planetary Science Letters*, 351-352, 123-133, 10.1016/j.epsl.2012.07.009, 2012.

Feng, X., Vonk, J. E., van Dongen, B. E., Gustafsson, O., Semiletov, I. P., Dudarev, O. V., Wang, Z., Montluçon, D. 605 B., Wacker, L., and Eglinton, T. I.: Differential mobilization of terrestrial carbon pools in Eurasian Arctic river basins, *Proceedings of the National Academy of Sciences of the United States of America*, 110, 14168-14173, 10.1073/pnas.1307031110, 2013.

Feng, X., Gustafsson, Ö., Holmes, R. M., Vonk, J. E., van Dongen, B. E., Semiletov, I. P., Dudarev, O. V., Yunker, M. B., Macdonald, R. W., Wacker, L., Montluçon, D. B., and Eglinton, T. I.: Multimolecular tracers of terrestrial

610 carbon transfer across the pan-Arctic: ^{14}C characteristics of sedimentary carbon components and their environmental controls, *Global Biogeochemical Cycles*, 29, 1855-1873, 10.1002/2015gb005204, 2015.

Ficken, K. J., Li, B., Swain, D., and Eglinton, G.: An n-alkane proxy for the sedimentary input of submerged/floating freshwater aquatic macrophytes, *Organic Geochemistry*, 31, 745-749, 2000.

Goñi, M. A. and Hedges, J. I.: Lignin dimers: Structures, distribution, and potential geochemical applications, 615 *Geochimica et Cosmochimica Acta*, 56, 4025-4043, 1992.

Goñi, M. A. and Montgomery, S.: Alkaline CuO oxidation with a microwave digestion system: Lignin analyses of geochemical samples, *Analytical Chemistry*, 72, 3116-3121, 2000.

Grotheer, H., Robert, A. M., Greenwood, P. F., and Grice, K.: Stability and hydrogenation of polycyclic aromatic hydrocarbons during hydrothermal degradation (HyPy) – Relevance for high maturity organic matter, *Organic Geochemistry*, 86, 620 45-54, 10.1016/j.orggeochem.2015.06.007, 2015.

Guay, C. K., Falkner, K. K., Muench, R. D., Mensch, M., Frank, M., and Bayer, R.: Wind-driven transport pathways for Eurasian Arctic river discharge, *Journal of Geophysical Research*, 106, 11469-11480, 2001.

Günther, F., Overduin, P. P., Yakshina, I. A., Opel, T., Baranskaya, A. V., and Grigoriev, M. N.: Observing Muostakh disappear: permafrost thaw subsidence and erosion of a ground-ice-rich island in response to arctic summer warming and sea ice reduction, *The Cryosphere*, 9, 151-178, 2015.

625 Heaton, T. J., Bard, E., Bronk Ramsey, C., Butzin, M., Hatté, C., Hughen, K. A., Köhler, P., and Reimer, P. J.: A Response to Community Questions on the Marine20 Radiocarbon Age Calibration Curve: Marine Reservoir Ages and the Calibration of ^{14}C Samples from the Oceans, *Radiocarbon*, 65, 247-273, 10.1017/rdc.2022.66, 2023.

Heaton, T. J., Köhler, P., Butzin, M., Bard, E., Reimer, R. W., Austin, W. E. N., Bronk Ramsey, C., Grootes, P. M., 630 Hughen, K. A., Kromer, B., Reimer, P. J., Adkins, J., Burke, A., Cook, M. S., Olsen, J., and Skinner, L. C.: Marine20—The Marine Radiocarbon Age Calibration Curve (0–55,000 cal BP), *Radiocarbon*, 62, 779-820, 10.1017/rdc.2020.68, 2020.

Hedges, J. I. and Mann, D. C.: The characterization of plant tissues by their lignin oxidation products, *Geochimica et Cosmochimica Acta*, 43, 1803-1807, 1979.

635 Hedges, J. I., Blanchette, R. A., Weliky, K., and Devol, A. H.: Effects of fungal degradation on the CuO oxidation products of lignin: a controlled laboratory study, *Geochimica et Cosmochimica Acta*, 52, 2717-2726, 1988.

Holmes, R., Shiklomanov, A. I., Suslova, A., Tretiakov, M., McClelland, J., Scott, L., Spencer, R., and Tank, S.: River discharge, United States. National Oceanic and Atmospheric Administration. Office of Oceanic and Atmospheric Research ;Global Ocean Monitoring and Observing (GOMO) Program, <https://doi.org/10.25923/zevf-ar65>, 2021.

640 Hörner, T., Stein, R., Fahl, K., and Birgel, D.: Post-glacial variability of sea ice cover, river run-off and biological production in the western Laptev Sea (Arctic Ocean)—A high-resolution biomarker study, *Quaternary Science Reviews*, 143, 133-149, 2016.

Hubberten, H. W., Andreev, A., Astakhov, V. I., Demidov, I., Dowdeswell, J. A., Henriksen, M., Hjort, C., Houmark-Nielsen, M., Jakobsson, M., Kuzmina, S., Larsen, E., Lunkka, J. P., Lysa, A., Mangerud, J., Möller, P., Saarnisto, M., 645 Schirmer, L., Sher, A. V., Siegert, C., Siegert, M. J., and Svendsen, J. I.: The periglacial climate and environment

in northern Eurasia during the Last Glaciation, *Quaternary Science Reviews*, 23, 1333-1357, 10.1016/j.quascirev.2003.12.012, 2004.

Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C. L., Schirmer, L., Grosse, G., Michaelson, G. J., Koven, C. D., O'Donnell, J. A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J., and Kuhry, P.: Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps, *Biogeosciences*, 11, 6573-6593, 10.5194/bg-11-6573-2014, 2014.

Hughes, A. L., Gyllencreutz, R., Lohne, Ø. S., Mangerud, J., and Svendsen, J. I.: The last Eurasian ice sheets—a chronological database and time-slice reconstruction, *DATED-1, Boreas*, 45, 1-45, 2016.

Irrgang, A. M., Bendixen, M., Farquharson, L. M., Baranskaya, A. V., Erikson, L. H., Gibbs, A. E., Ogorodov, S. A., Overduin, P. P., Lantuit, H., Grigoriev, M. N., and Jones, B. M.: Drivers, dynamics and impacts of changing Arctic coasts, *Nature Reviews Earth & Environment*, 3, 39-54, 10.1038/s43017-021-00232-1, 2022.

Jakobsson, M., Mayer, L., Coakley, B., Dowdeswell, J. A., Forbes, S., Fridman, B., Hodnesdal, H., Noormets, R., Pedersen, R., and Rebecq, M.: The international bathymetric chart of the Arctic Ocean (IBCAO) version 3.0, *Geophysical Research Letters*, 39, 2012.

Jakobsson, M., Pearce, C., Cronin, T. M., Backman, J., Anderson, L. G., Barrientos, N., Björk, G., Coxall, H., de Boer, A., Mayer, L. A., Mörrth, C.-M., Nilsson, J., Rattray, J. E., Stranne, C., Semiletov, I., and O'Regan, M.: Post-glacial flooding of the Bering Land Bridge dated to 11 cal ka BP based on new geophysical and sediment records, *Climate of the Past*, 13, 991-1005, 10.5194/cp-13-991-2017, 2017.

Kassens, H.: Station list and links to master tracks in different resolutions of POLARSTERN cruise ARK-XIV/1b, Tiksi - Tromsø, 1998-07-28 - 1998-08-28, PANGAEA [dataset], <https://doi.org/10.1594/PANGAEA.858867>, 2016.

Keigwin, L. D., Klotsko, S., Zhao, N., Reilly, B., Giosan, L., and Driscoll, N. W.: Deglacial floods in the Beaufort Sea preceded Younger Dryas cooling, *Nature Geoscience*, 11, 599-604, 10.1038/s41561-018-0169-6, 2018.

Keskitalo, K., Tesi, T., Bröder, L., Andersson, A., Pearce, C., Sköld, M., Semiletov, I. P., Dudarev, O. V., and Gustafsson, Ö.: Sources and characteristics of terrestrial carbon in Holocene-scale sediments of the East Siberian Sea, *Climate of the Past*, 13, 1213-1226, 10.5194/cp-13-1213-2017, 2017.

Kleiber, H. P. and Niessen, F.: Variations of continental discharge pattern in space and time: implications from the Laptev Sea continental margin, Arctic Siberia, *International Journal of Earth Sciences*, 89, 605-616, 10.1007/s005310000130, 2000.

Klemann, V., Heim, B., Bauch, H. A., Wetterich, S., and Opel, T.: Sea-level evolution of the Laptev Sea and the East Siberian Sea since the last glacial maximum, *Arktos*, 1, 10.1007/s41063-015-0004-x, 2015.

Klotsko, S., Driscoll, N., and Keigwin, L.: Multiple meltwater discharge and ice rafting events recorded in the deglacial sediments along the Beaufort Margin, Arctic Ocean, *Quaternary Science Reviews*, 203, 185-208, 10.1016/j.quascirev.2018.11.014, 2019.

Köhler, P., Nehrbass-Ahles, C., Schmitt, J., Stocker, T. F., and Fischer, H.: A 156 kyr smoothed history of the atmospheric greenhouse gases CO₂, CH₄, and N₂O and their radiative forcing, *Earth System Science Data*, 9, 363-387, 10.5194/essd-9-363-2017, 2017.

- Kuehn, H., Lembke-Jene, L., Gersonde, R., Esper, O., Lamy, F., Arz, H., Kuhn, G., and Tiedemann, R.: Laminated sediments in the Bering Sea reveal atmospheric teleconnections to Greenland climate on millennial to decadal timescales during the last deglaciation, *Climate of the Past*, 10, 2215–2236, 10.5194/cp-10-2215-2014, 2014.
- 685 Kuptsov, V. and Lisitsin, A.: Radiocarbon of Quaternary along shore and bottom deposits of the Lena and the Laptev Sea sediments, *Marine Chemistry*, 53, 301–311, 1996.
- Lambeck, K., Rouby, H., Purcell, A., Sun, Y., and Sambridge, M.: Sea level and global ice volumes from the Last Glacial Maximum to the Holocene, *Proceedings of the National Academy of Sciences of the United States of America*, 111, 15296–15303, 10.1073/pnas.1411762111, 2014.
- 690 Lantuit, H., Atkinson, D., Paul Overduin, P., Grigoriev, M., Rachold, V., Grosse, G., and Hubberten, H.-W.: Coastal erosion dynamics on the permafrost-dominated Bykovsky Peninsula, north Siberia, 1951–2006, *Polar Research*, 30, 10.3402/polar.v30i0.7341, 2011.
- Lara, M. J., Lin, D. H., Andresen, C., Lougheed, V. L., and Tweedie, C. E.: Nutrient release from permafrost thaw enhances CH₄ emissions from arctic tundra wetlands, *Journal of Geophysical Research: Biogeosciences*, 124, 1560–1573, 10.1029/2018jg004641, 2019.
- 695 Laurent, M., Fuchs, M., Herbst, T., Runge, A., Liebner, S., and Treat, C.: Relationships between greenhouse gas production and landscape position during short-term permafrost thaw under anaerobic conditions in the Lena Delta., *Biogeosciences*, 20, 2049–2064, 10.5194/bg-2022-122, 2023.
- Lehner, B. and Grill, G.: Global river hydrography and network routing: baseline data and new approaches to study the world's large river systems, *Hydrological Processes*, 27, 2171–2186, 2013.
- 700 Limber, P. W., Barnard, P. L., Vitousek, S., and Erikson, L. H.: A Model Ensemble for Projecting Multidecadal Coastal Cliff Retreat During the 21st Century, *Journal of Geophysical Research: Earth Surface*, 123, 1566–1589, 10.1029/2017jg004401, 2018.
- Liu, S., Wang, P., Huang, Q., Yu, J., Pozdniakov, S. P., and Kazak, E. S.: Seasonal and spatial variations in riverine DOC exports in permafrost-dominated Arctic river basins, *Journal of Hydrology*, 612, 10.1016/j.jhydrol.2022.128060, 2022.
- 705 Marcott, S. A., Bauska, T. K., Buizert, C., Steig, E. J., Rosen, J. L., Cuffey, K. M., Fudge, T. J., Severinghaus, J. P., Ahn, J., Kalk, M. L., McConnell, J. R., Sowers, T., Taylor, K. C., White, J. W., and Brook, E. J.: Centennial-scale changes in the global carbon cycle during the last deglaciation, *Nature*, 514, 616–619, 10.1038/nature13799, 2014.
- 710 Margold, M., Jansen, J. D., Codilean, A. T., Preusser, F., Gurinov, A. L., Fujioka, T., and Fink, D.: Repeated megafloods from glacial Lake Vitim, Siberia, to the Arctic Ocean over the past 60,000 years, *Quaternary Science Reviews*, 187, 41–61, 10.1016/j.quascirev.2018.03.005, 2018.
- Martens, J., Wild, B., Muschitiello, F., O'Regan, M., Jakobsson, M., Semiletov, I., Dudarev, O. V., and Gustafsson, Ö.: Remobilization of dormant carbon from Siberian-Arctic permafrost during three past warming events, *Science Advances*, 6, eabb6546, 2020.
- 715 Martens, J., Wild, B., Pearce, C., Tesi, T., Andersson, A., Broder, L., O'Regan, M., Jakobsson, M., Skold, M., Gemery, L., Cronin, T. M., Semiletov, I., Dudarev, O. V., and Gustafsson, O.: Remobilization of Old Permafrost Carbon to

- Chukchi Sea Sediments During the End of the Last Deglaciation, *Global Biogeochemical Cycles*, 33, 2-14, 10.1029/2018GB005969, 2019.
- 720 McClelland, J. W., Holmes, R. M., Peterson, B. J., Raymond, P. A., Striegl, R. G., Zhulidov, A. V., Zimov, S. A., Zimov, N., Tank, S. E., Spencer, R. G. M., Staples, R., Gurtovaya, T. Y., and Griffin, C. G.: Particulate organic carbon and nitrogen export from major Arctic rivers, *Global Biogeochemical Cycles*, 30, 629-643, 10.1002/2015gb005351, 2016.
- 725 Meredith, M., Sommerkorn, M., Cassotta, S., Derksen, C., Ekaykin, A., Hollowed, A., Kofinas, G., Mackintosh, A., Melbourne-Thomas, J., Muelbert, M. M. C., Ottersen, G., Pritchard, H., and E.A.G., S.: Polar Regions, in: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate, edited by: H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, and Weyer, N. M., Cambridge University Press, Cambridge, UK and New York, NY, USA, 203–320, <https://doi.org/10.1017/9781009157964.005>, 2019.
- 730 Meyer, V. D., Hefter, J., Köhler, P., Tiedemann, R., Gersonde, R., Wacker, L., and Mollenhauer, G.: Permafrost-carbon mobilization in Beringia caused by deglacial meltwater runoff, sea-level rise and warming, *Environmental Research Letters*, 14, 10.1088/1748-9326/ab2653, 2019.
- 735 Miller, G. H., Alley, R. B., Brigham-Grette, J., Fitzpatrick, J. J., Polyak, L., Serreze, M. C., and White, J. W. C.: Arctic amplification: can the past constrain the future?, *Quaternary Science Reviews*, 29, 1779-1790, 10.1016/j.quascirev.2010.02.008, 2010.
- Mollenhauer, G., Grotheer, H., Gentz, T., Bonk, E., and Hefter, J.: Standard operation procedures and performance of the MICADAS radiocarbon laboratory at Alfred Wegener Institute (AWI), Germany, *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 496, 45-51, 10.1016/j.nimb.2021.03.016, 2021.
- 740 Mueller-Lupp, T., Bauch, H. A., Erlenkeuser, H., Hefter, J., Kassens, H., and Thiede, J.: Changes in the deposition of terrestrial organic matter on the Laptev Sea shelf during the Holocene: evidence from stable carbon isotopes, *International Journal of Earth Sciences*, 89, 563-568, 10.1007/s005310000128, 2000.
- 745 Müller, J. and Stein, R.: High-resolution record of late glacial and deglacial sea ice changes in Fram Strait corroborates ice–ocean interactions during abrupt climate shifts, *Earth and Planetary Science Letters*, 403, 446-455, 10.1016/j.epsl.2014.07.016, 2014.
- Müller, J., Massé, G., Stein, R., and Belt, S. T.: Variability of sea-ice conditions in the Fram Strait over the past 30,000 years, *Nature Geoscience*, 2, 772-776, 10.1038/ngeo665, 2009.
- 750 Nielsen, D. M., Dobrynin, M., Baehr, J., Razumov, S., and Grigoriev, M.: Coastal Erosion Variability at the Southern Laptev Sea Linked to Winter Sea Ice and the Arctic Oscillation, *Geophysical Research Letters*, 47, 10.1029/2019gl086876, 2020.
- Nogarotto, A., Noormets, R., Chauhan, T., Mollenhauer, G., Hefter, J., Grotheer, H., Belt, S., Colleoni, F., Muschitiello, F., Capotondi, L., Pellegrini, C., and Tesi, T.: Coastal permafrost was massively eroded during the Bølling-Allerød warm period, *Communications Earth & Environment*, 4, 350, <https://doi.org/10.1038/s43247-023-01013-y>, 2023.

755 Overduin, P. P., Wetterich, S., Günther, F., Grigoriev, M. N., Grosse, G., Schirrmeister, L., Hubberten, H.-W., and Makarov, A.: Coastal dynamics and submarine permafrost in shallow water of the central Laptev Sea, East Siberia, *The Cryosphere*, 10, 1449-1462, 10.5194/tc-10-1449-2016, 2016.

Rachold, V., Grigoriev, M. N., Are, F. E., Solomon, S., Reimnitz, E., Kassens, H., and Antonow, M.: Coastal erosion vs riverine sediment discharge in the Arctic Shelf seas, *International Journal of Earth Sciences*, 89, 450-460, 10.1007/s005310000113, 2000.

760 Rantanen, M., Karpechko, A. Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Communications Earth & Environment*, 3, 10.1038/s43247-022-00498-3, 2022.

Reimer, P. J., Austin, W. E. N., Bard, E., Bayliss, A., Blackwell, P. G., Bronk Ramsey, C., Butzin, M., Cheng, H., 765 Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kromer, B., Manning, S. W., Muscheler, R., Palmer, J. G., Pearson, C., van der Plicht, J., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Turney, C. S. M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S. M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., and Talamo, S.: The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP), 770 *Radiocarbon*, 62, 725-757, 10.1017/rdc.2020.41, 2020.

Rigor, I. and Colony, R.: Sea-ice production and transport of pollutants in the Laptev Sea, 1979–1993, *Science of the Total Environment*, 202, 89-110, 1997.

Rudenko, O., Tarasov, P. E., Bauch, H. A., and Taldenkova, E.: A Holocene palynological record from the northeastern Laptev Sea and its implications for palaeoenvironmental research, *Quaternary International*, 348, 82-92, 775 10.1016/j.quaint.2014.04.032, 2014.

Sabino, M., Gustafsson, Ö., Wild, B., Semiletov, I. P., Dudarev, O. V., Ingrosso, G., and Tesi, T.: Feedbacks From Young Permafrost Carbon Remobilization to the Deglacial Methane Rise, *Global Biogeochemical Cycles*, 38, 10.1029/2024gb008164, 2024.

Sánchez-García, L., Vonk, J. E., Charkin, A. N., Kosmach, D., Dudarev, O. V., Semiletov, I. P., and Gustafsson, Ö.: 780 Characterisation of Three Regimes of Collapsing Arctic Ice Complex Deposits on the SE Laptev Sea Coast using Biomarkers and Dual Carbon Isotopes, *Permafrost and Periglacial Processes*, 25, 172-183, 10.1002/ppp.1815, 2014.

Schuur, E. A., Bockheim, J., Canadell, J. G., Euskirchen, E., Field, C. B., Goryachkin, S. V., Hagemann, S., Kuhry, P., Lafleur, P. M., and Lee, H.: Vulnerability of permafrost carbon to climate change: Implications for the global carbon cycle, *BioScience*, 58, 701-714, 2008.

785 Schuur, E. A., McGuire, A. D., Schadel, C., Grosse, G., Harden, J. W., Hayes, D. J., Hugelius, G., Koven, C. D., Kuhry, P., Lawrence, D. M., Natali, S. M., Olefeldt, D., Romanovsky, V. E., Schaefer, K., Turetsky, M. R., Treat, C. C., and Vonk, J. E.: Climate change and the permafrost carbon feedback, *Nature*, 520, 171-179, 10.1038/nature14338, 2015.

Shadrick, J. R., Rood, D. H., Hurst, M. D., Piggott, M. D., Hebditch, B. G., Seal, A. J., and Wilcken, K. M.: Sea-level 790 rise will likely accelerate rock coast cliff retreat rates, *Nature Communications*, 13, 7005, 10.1038/s41467-022-34386-3, 2022.

- Shakun, J. D., Clark, P. U., He, F., Marcott, S. A., Mix, A. C., Liu, Z., Otto-Bliesner, B., Schmittner, A., and Bard, E.: Global warming preceded by increasing carbon dioxide concentrations during the last deglaciation, *Nature*, 484, 49-54, 10.1038/nature10915, 2012.
- 795 Smith, L., MacDonald, G., Velichko, A., Beilman, D., Borisova, O., Frey, K., Kremenetski, K., and Sheng, Y.: Siberian peatlands a net carbon sink and global methane source since the early Holocene, *Science*, 303, 353-356, 2004.
- Spielhagen, R., Erlenkeuser, H., and Siebert, C.: History of freshwater runoff across the Laptev Sea (Arctic) during the last deglaciation, *Global and Planetary Change*, 48, 187-207, 10.1016/j.gloplacha.2004.12.013, 2005.
- Stein, R. and Fahl, K.: Holocene accumulation of organic carbon at the Laptev Sea continental margin (Arctic Ocean):
800 sources, pathways, and sinks, *Geo-Marine Letters*, 20, 27-36, 2000.
- Stein, R. and Macdonald, R. W., Stein, R., and MacDonald, R. W. (Eds.): The organic carbon cycle in the Arctic Ocean, Springer Berlin, Heidelberg, <https://doi.org/10.1007/978-3-642-18912-8>, 2004.
- Strauss, J., Schirrmeister, L., Mangelsdorf, K., Eichhorn, L., Wetterich, S., and Herzsuh, U.: Organic-matter quality of deep permafrost carbon – a study from Arctic Siberia, *Biogeosciences*, 12, 2227-2245, 10.5194/bg-12-2227-2015,
805 2015.
- Strauss, J., Schirrmeister, L., Grosse, G., Fortier, D., Hugelius, G., Knoblauch, C., Romanovsky, V., Schädel, C., Schneider von Deimling, T., Schuur, E. A. G., Shmelev, D., Ulrich, M., and Veremeeva, A.: Deep Yedoma permafrost: A synthesis of depositional characteristics and carbon vulnerability, *Earth-Science Reviews*, 172, 75-86, 10.1016/j.earscirev.2017.07.007, 2017.
- 810 Strauss, J., Laboor, S., Fedorov, A. N., Fortier, D., Froese, D. G., Fuchs, M., Grosse, G., Günther, F., Harden, J. W., Hugelius, G., Kanevskiy, M. Z., Kholodov, A. L., Kunitsky, V. V., Kraev, G., Lapointe-Elmrabti, L., Lozhkin, A. V., Rivkina, E., Robinson, J., Schirrmeister, L., Shmelev, D., Shur, Y., Siebert, C., Spektor, V., Ulrich, M., Vartanyan, S. L., Veremeeva, A., Walter Anthony, K. M., and Zimov, S. A.: Database of Ice-Rich Yedoma Permafrost (IRYP) [dataset], <https://doi.org/10.1594/PANGAEA.861733>, 2016.
- 815 Strauss, J., Laboor, S., Schirrmeister, L., Fedorov, A. N., Fortier, D., Froese, D., Fuchs, M., Günther, F., Grigoriev, M., Harden, J., Hugelius, G., Jongejans, L. L., Kanevskiy, M., Kholodov, A., Kunitsky, V., Kraev, G., Lozhkin, A., Rivkina, E., Shur, Y., Siebert, C., Spektor, V., Streletskaia, I., Ulrich, M., Vartanyan, S., Veremeeva, A., Anthony, K. W., Wetterich, S., Zimov, N., and Grosse, G.: Circum-Arctic Map of the Yedoma Permafrost Domain, *Frontiers in Earth Science*, 9, 10.3389/feart.2021.758360, 2021.
- 820 Strauss, J., Laboor, S., Schirrmeister, L., Fedorov, A. N., Fortier, D., Froese, D. G., Fuchs, M., Günther, F., Grigoriev, M. N., Harden, J. W., Hugelius, G., Jongejans, L. L., Kanevskiy, M. Z., Kholodov, A. L., Kunitsky, V., Kraev, G., Lozhkin, A. V., Rivkina, E., Shur, Y., Siebert, C., Spektor, V., Streletskaia, I., Ulrich, M., Vartanyan, S. L., Veremeeva, A., Walter Anthony, K. M., Wetterich, S., Zimov, N. S., and Grosse, G.: Database of Ice-Rich Yedoma Permafrost Version 2 (IRYP v2), PANGAEA [dataset], 10.1594/PANGAEA.940078, 2022.
- 825 Taldenkova, E., Bauch, H. A., Gottschalk, J., Nikolaev, S., Rostovtseva, Y., Pogodina, I., Ovsepyan, Y., and Kandiano, E.: History of ice-rafting and water mass evolution at the northern Siberian continental margin (Laptev Sea) during Late Glacial and Holocene times, *Quaternary Science Reviews*, 29, 3919-3935, 2010.

- Terhaar, J., Lauerwald, R., Regnier, P., Gruber, N., and Bopp, L.: Around one third of current Arctic Ocean primary production sustained by rivers and coastal erosion, *Nature Communications*, 12, 169, 10.1038/s41467-020-20470-z, 2021.
- 830 Tesi, T., Puig, P., Palanques, A., and Goñi, M. A.: Lateral advection of organic matter in cascading-dominated submarine canyons, *Progress in Oceanography*, 84, 185-203, 10.1016/j.pocean.2009.10.004, 2010.
- Tesi, T., Semiletov, I., Dudarev, O., Andersson, A., and Gustafsson, Ö.: Matrix association effects on hydrodynamic sorting and degradation of terrestrial organic matter during cross-shelf transport in the Laptev and East Siberian shelf
- 835 seas, *Journal of Geophysical Research: Biogeosciences*, 121, 731-752, 10.1002/2015jg003067, 2016a.
- Tesi, T., Semiletov, I., Hugelius, G., Dudarev, O., Kuhry, P., and Gustafsson, Ö.: Composition and fate of terrigenous organic matter along the Arctic land–ocean continuum in East Siberia: Insights from biomarkers and carbon isotopes, *Geochimica et Cosmochimica Acta*, 133, 235-256, 10.1016/j.gca.2014.02.045, 2014.
- Tesi, T., Muschitiello, F., Smittenberg, R. H., Jakobsson, M., Vonk, J. E., Hill, P., Andersson, A., Kirchner, N.,
- 840 Noormets, R., Dudarev, O., Semiletov, I., and Gustafsson, O.: Massive remobilization of permafrost carbon during post-glacial warming, *Nature Communications*, 7, 13653, 10.1038/ncomms13653, 2016b.
- van Dongen, B. E., Semiletov, I., Weijers, J. W. H., and Gustafsson, Ö.: Contrasting lipid biomarker composition of terrestrial organic matter exported from across the Eurasian Arctic by the five great Russian Arctic rivers, *Global Biogeochemical Cycles*, 22, 10.1029/2007gb002974, 2008.
- 845 van Everdingen, R. O., van Everdingen, R. O. (Ed.): Multi-language glossary of permafrost and related ground-ice terms, International Permafrost Association, Calgary, Alberta, Canada, 159 pp.2005.
- Vandenberghe, J., Renssen, H., Roche, D. M., Goosse, H., Velichko, A. A., Gorbunov, A., and Levvasseur, G.: Eurasian permafrost instability constrained by reduced sea-ice cover, *Quaternary Science Reviews*, 34, 16-23, 10.1016/j.quascirev.2011.12.001, 2012.
- 850 Volkman, J. K., Barrett, S. M., and Dunstan, G. A.: C25 and C30 highly branched isoprenoid alkenes in laboratory cultures of two marine diatoms, *Organic Geochemistry*, 21, 407-414, 1994.
- Vonk, J. E. and Gustafsson, Ö.: Calibrating n-alkane Sphagnum proxies in sub-Arctic Scandinavia, *Organic Geochemistry*, 40, 1085-1090, 10.1016/j.orggeochem.2009.07.002, 2009.
- Vonk, J. E. and Gustafsson, Ö.: Permafrost-carbon complexities, *Nature Geoscience*, 6, 675-676, 2013.
- 855 Wild, B., Shakhova, N., Dudarev, O., Ruban, A., Kosmach, D., Tumskey, V., Tesi, T., Grimm, H., Nybom, I., Matsubara, F., Alexanderson, H., Jakobsson, M., Mazurov, A., Semiletov, I., and Gustafsson, O.: Organic matter composition and greenhouse gas production of thawing subsea permafrost in the Laptev Sea, *Nature Communications*, 13, 5057, 10.1038/s41467-022-32696-0, 2022.
- Winterfeld, M., Goñi, M. A., Just, J., Heffer, J., and Mollenhauer, G.: Characterization of particulate organic matter
- 860 in the Lena River delta and adjacent nearshore zone, NE Siberia – Part 2: Lignin-derived phenol compositions, *Biogeosciences*, 12, 2261-2283, 10.5194/bg-12-2261-2015, 2015.
- Winterfeld, M., Mollenhauer, G., Dumann, W., Kohler, P., Lembke-Jene, L., Meyer, V. D., Heffer, J., McIntyre, C., Wacker, L., Kokfelt, U., and Tiedemann, R.: Deglacial mobilization of pre-aged terrestrial carbon from degrading permafrost, *Nature Communications*, 9, 3666, 10.1038/s41467-018-06080-w, 2018.

- 865 Wu, J., Stein, R., Fahl, K., Syring, N., Nam, S.-I., Hefter, J., Mollenhauer, G., and Geibert, W.: Deglacial to Holocene variability in surface water characteristics and major floods in the Beaufort Sea, *Communications Earth & Environment*, 1, 10.1038/s43247-020-00028-z, 2020.
- Wu, J., Mollenhauer, G., Stein, R., Kohler, P., Hefter, J., Fahl, K., Grotheer, H., Wei, B., and Nam, S. I.: Deglacial release of petrogenic and permafrost carbon from the Canadian Arctic impacting the carbon cycle, *Nature*
- 870 *Communications*, 13, 7172, 10.1038/s41467-022-34725-4, 2022.
- Yu, Z., Loisel, J., Brosseau, D. P., Beilman, D. W., and Hunt, S. J.: Global peatland dynamics since the Last Glacial Maximum, *Geophysical Research Letters*, 37, 10.1029/2010gl043584, 2010.
- Zamelczyk, K., Rasmussen, T. L., Husum, K., Godtliebsen, F., and Hald, M.: Surface water conditions and calcium carbonate preservation in the Fram Strait during marine isotope stage 2, 28.8–15.4 kyr, *Paleoceanography*, 29, 1-12,
- 875 2014.
- Zhang, T., Li, D., East, A. E., Walling, D. E., Lane, S., Overeem, I., Beylich, A. A., Koppes, M., and Lu, X.: Warming-driven erosion and sediment transport in cold regions, *Nature Reviews Earth & Environment*, 3, 832-851, 10.1038/s43017-022-00362-0, 2022.

880